

Population dynamics of the red bigeye (*Priacanthus macracanthus* Cuvier, 1829) (Fish: Family Priacanthidae)

by Meuthia A. Jabbar, mohamma M. Kamal, Mennofatria B Ali Suman, i
Nyoman Suyasa, Erfind Nurdin


Submission date: 31-May-2022 04:33PM (UTC+0700)

Submission ID: 1847715984

File name: Pop_dyn_red_bigeye_Plbratu_MAJ_Oct_2017.pdf (279.26K)

Word count: 4933

Character count: 25255



Population dynamics of the red bigeye (*Priacanthus macracanthus* Cuvier, 1829) (Fish: Family Priacanthidae) in Palabuhanratu Bay, Indonesia

^{1,2}Meuthia A. Jabbar, ²Mohammad M. Kamal, ²Mennofatria Boer, ³Ali Suman, ¹I Nyoman Suyasa, ³Erfind Nurdin

23

¹ Department of Aquatic Resources Management, Jakarta Fisheries University, Indonesia; ² Department of Aquatic Resources Management, Faculty of Fisheries and Marine Science, Bogor Agricultural University, Darmaga-Bogor, Indonesia;

³ Research Institute for Marine Fisheries, Ministry of Marine Affairs and Fisheries, Jakarta, Indonesia. Corresponding author: M. A. Jabbar, meuthia.aula@gmail.com

Abstract. The red bigeye (*Priacanthus macracanthus*) is an economically important demersal fish species which is intensively exploited in Indonesia. A study on population dynamics of *P. macracanthus* has been conducted in Palabuhanratu Bay from January to December 2016. Fish sample was collected at Palabuhanratu Fish Landing Station of which the catch was mainly derived from bottom gillnet. Total length data was presented and calculated by Excel spreadsheet and analyzed further by ELEFAN I (FISAT II). The results showed that the size of *P. macracanthus* ranged between 5-34 cmTL, with a mode between 21-22 cmTL. The growth coefficient (K) was equal to 1.38 year⁻¹ which asymptotic length (L_∞) up to 35.7 cm TL, theoretical age (t₀) -0.21 years and growth performance index (Ø') 3.25. The rate of natural mortality (M), fishing mortality (F), and accordingly total mortality (Z) was 2.14 year⁻¹, 2.51 year⁻¹, and 4.64 year⁻¹, respectively. The recruitment occurs twice a year in February/March and August/September. The rate of exploitation (E_{cur}) 0.54 on Y/R_{cur} 0.04378 is closer to the value (E₁₀) 0.553 than (E₅₀) 0.353, indicating at fully exploited condition of the fishery. It is suggested to adopt E₁₀ (economic yield) as reference point of red bigeyes's fisheries management in Bay of Palabuhanratu.

Key Words: growth parameters, mortality rate, exploitation rate, recruitment pattern, fisheries management.

Introduction. The red bigeye (*Priacanthus macracanthus*) belongs to family Priacanthidae (Froese & Pauly 2017), among an important demersal target species in capture fishery of Palabuhanratu Bay, southern Java Sea, part of Indian Ocean. This species is mainly caught by bottom gillnet, and to the lesser extent by hand line as well as incidentally collected by lift net. They inhabit coral reefs environment (Starnes 1999; Ramachandran & Varghese 2009; White et al 2013), which plays a major role in the trophic structure (mainly in coral reef ecosystem) (Powell 2000).

In Asian countries, member of *Priacanthidae* family are mainly economic important species for its high protein content. They are major elements for a variety of typical Asiatic cuisines including noodles, fish balls, sauces and surimi products. For instance, the moontail bullseye, *Priacanthus hamrur* with 20.8% protein content is used for fish powder (Sivakami et al 2003; Safitri et al 2016). In the following of high demand and increasing price, hence, exploitation rate of those priacanthids species is considered to be high leading to the importance of studying its population dynamic.

During the last decade, studies on *P. macracanthus* population dynamics has been reported in the Southeast Asia including Java Sea, the West and East Coast of Malaysia Peninsular, Taiwan Waters, the South and East China Sea (Nugroho & Rustam 1983; Lester & Watson 1985; Joung & Chen 1992; Liu et al 1992; Liu & Cheng 1999; Oki & Tabeta 1999; Silvestre & Garces 2004). Some authors concludes that *P. macracanthus*

population has already over exploited such as in Java Sea (Dwiponggo et al 1986), Malaysia Peninsular (Ahmad et al 2003), the Northern part of South China Sea (Sun & Qiu 2004), Northern Taiwan (Ju et al 2016), and Chinese Beibu Gulf (Zhang et al 2016). However, *P. macracanthus* population in Palabuhanratu Bay has never been studied for such important aspects. In fact, this area is one of the important fishing ports in Indonesia where the *P. macracanthus* and other demersal species such as moonfish (*Mene maculata*), slipmouths (*Leiognathus* spp.), trasih prawn (*Mysis* sp.), and hairtails (*Trichiurus* spp.) were landed (PPNP 2015).

In order to understand the current status and for better management of the *P. macracanthus*, *P. macracanthus* in Palabuhanratu Bay, the present study has been focused on the population dynamic of this species. In this case the present paper deals with growth, mortality, capture probability, recruitment pattern, exploitation rate, and relative yield per recruit.

Material and Method. Study area was situated in Palabuhanratu Bay, where fishing ground has been defined participatorily by local fishermen on map, where the catch are transported to the nearest fish landing site (Figure 1). Data collection was done from January to December 2016 with monthly interval. Fish sample was collected from the catch using bottom gillnet, hand line and lift net, of which selected sample was total length measured (TL) with 0.1 cm accuracy. Length data is the main prerequisite for every analysis used in this study.




Figure 1. Fishing ground and fish landing site for *Priacanthus macracanthus* in Palabuhanratu Bay, West Java, Indian Ocean.

Growth parameters estimation. Growth coefficient (K) and asymptotic length (L_{∞}) was estimated using ELEFAN (Electronic Length Frequency Analysis) I model which made by FISAT (FAO-ICLARM Stock Assessment Tools) II (Gayanilo et al 2005). First of all, the theoretical age (t_0) is determined by Pauly's formula (1983):

$$\text{Log}(-t_0) = -0.3922 - 0.2752 \text{Log} L_{\infty} - 1.03 \text{Log} K$$

The values of length, K, and t_0 are then calculated to build von Bertalanffy's growth model (Sparre & Venema 1998) as follows:

$$L_t = L_{\infty} (1 - e^{-K(t-t_0)})$$

where L_t is the length at age t (cm), L_{∞} is the asymptotic length (cm), K is the growth coefficient (year^{-1}) and t_0 is the theoretical age (year).

In order to explore the growth rate in *P. macracanthus* of several different stock, a

growth performance index (ϕ') is calculated (Pauly & Munro 1984):

$$\phi' = \text{Log } K + 2 \text{ Log } L_{\infty}$$

Mortality. Total mortality (Z) is predicted using Length-Converted Catch Curve made by FISAT II (Gayanilo et al 2005). In estimating natural mortality rate (M), the formula from Pauly's (1983) is applied:

$$\text{Log } (M) = -0.0066 - 0.2795 \text{ Log } (L_{\infty}) + 0.6543 \text{ Log } (K) + 0.4634 \text{ Log } (T)$$

where M is the natural mortality (year^{-1}) and T is the average annual water temperature ($^{\circ}\text{C}$).

Having know Z and M values, fishing mortality (F) is accordingly estimated after Pauly (1983):

$$F = Z - M$$

Recruitment pattern. By projecting the length-frequency data backward onto the time axis down to zero length, using the von Bertalanffy growth equation and the estimated growth parameters, the recruitment pattern of the *P. macracanthus* was estimated. It used program package FISAT II (Gayanilo et al 2005).

Exploitation rate. Based on a year data collection, the exploitation rate (E) is used as proxy to estimate current utilization of *P. macracanthus* (Gulland 1971):

$$E = \frac{F}{F+M} = \frac{F}{Z}$$

Relative yield per recruit (Y'/R). Relative yield per recruit (Y'/R) is estimated using Beverton and Holt Model (Sparre & Venema 1998) in FISAT II program packages and also in Excel worksheet. From this, the values of exploitation rate at maximum relative yield per recruit (E_{max}) could be estimated. Also E_{10} , the value of E at which the marginal increase in Y'/R is 10% of its value at $E=0$; and E_{50} , the value of E corresponding to 50% of the relative biomass per-recruit (B'/R), were estimated. The formulae of Y'/R as follows:

$$\frac{Y'}{R} = EU^{\frac{M}{K}} \left\{ 1 - \left(\frac{3U}{1+m} \right) + \left(\frac{3U^2}{1+2m} \right) - \left(\frac{U^3}{1+3m} \right) \right\}$$

where $U = 1 - (L_c / L_{\infty})$, $m = (1-E) / (M/K)$ and $E = F/Z$

Results and Discussion

Growth parameters estimation. A total of 3,420 *P. macracanthus* individuals were collected from the catch of bottom gillnet, handline, and liftnet Palabuhanratu Bay during the year. These three fishing gears contribute to 87.7%, 11.9% and 0.3 % of total catch, respectively. Based on total length data, catch composition distributed skewedly where modus values were 21-22 cm, 23-24 cm, 19-20 cm, and 7-8 cm in range size (Figure 2).

Analyzed using TL data, growth coefficient (K) was 1.38 year^{-1} , with asymptotic length (L_{∞}) 35.7 cm, and theoretical age (t_0) -0.21 years (Figure 3). Based on these, growth model for *P. macracanthus* in Palabuhanratu could be formulated as $L_t = 35.7(1 - e^{-1.38(t+0.21)})$ (Figure 4). Fish age of 1, 2, and 3 years old coincides with total length of 29, 34 and 35.3 cm, consecutively. A highest growth rate was found at 0-1 year with $1.82 \text{ cm month}^{-1}$ in average, followed by 1-2 years with $0.46 \text{ cm month}^{-1}$. The asymptotic length was 35.7 cm, i.e. size at maximum length is reached in 4.6 years.

In comparing growth from the different stock, it was found that growth performance index (ϕ') ranged between 2.31 and 3.25, of which the highest ϕ' value was obtained in Palabuhanratu Bay, indicating that *P. macracanthus* grow at fastest rate in this area (Table 1).




Figure 2. Length frequency distribution of *Priacanthus macracanthus* composed by catch from 3 fishing gears during research in Palabuhanratu Bay.




Figure 3. The growth model of *Priacanthus macracanthus* in Palabuhanratu Bay.




Figure 4. Length and ages relationship curve of *Priacanthus macracanthus* in Palabuhanratu Bay.

Table 1 shows the spatial variability of K ranged between 0.08 - 1.38 year^{-1} , of which K 's *P. macracanthus* population in Palabuhanratu Bay was the highest. A slight differ in growth coefficient occurred with the stock in similar larger sea ecosystem. The previous reports (Sparre & Venema 1998; Pardo et al 2013) pointed out that $K \geq 0.5$ of *P. macracanthus* was categorized into rapid growth species. This was the case for population in Palabuhanratu Bay, Java Sea, South China Sea, and Indian Ocean. A slower growth was found in Taiwan waters. Accordingly, fastest growth rate is associated with smaller L_{∞} (length infinity), meaning that *P. macracanthus* belongs to short live fish, of which fish reaches its maximum size of 35.7 cm for 4.6 years which constituted by 20 cm , 5 cm and 1.3 cm of annual growth rate. In comparison, in Taiwan water, the same species reached the length infinity at the age of 6 with the estimated combined growth rate of male and female fish during the first, second and third year respectively 10.7 cm , 4.2 cm , and 3.75 (Joung & Chen 1992). In general, in addition to its dependence on behavior, fish growth is highly dependent on environmental factors, especially the food availability, optimum temperatures and genetic factors (Magnussen 2007). The implication is that there may be major differences between waters and interestingly, some areas (ecosystems) are more productive than others. Gjedrem (2000) found that 70 - 80% of fish growth in temperate regions is determined by environmental factors, while the rests are thought to be influenced by genetic factors.

P. macracanthus in waters adjacent to the Indian Ocean (including Palabuhanratu Bay) have a higher \emptyset' value than those found in other areas, as presented in Table 1. *P. macracanthus* in the Java Sea and South China Sea have a lower \emptyset' value than those observed in the Indian Ocean but higher than in the waters of Taiwan. Specific ecological factors in different areas such as population density, water environmental temperature and prey density may cause variations in the value of growth parameters (Ju et al 2016). Differences in growth parameters and growth performance indexes between different regions can be attributed to differences in species-to-species composition and method of analysis (Mehanna et al 2013).

Mortality. Natural mortality rate (M) of *P. macracanthus* in Palabuhanratu Bay was 0.26 . 14 year^{-1} at annual average temperature 29°C . Fishing mortality rate (F) was 2.51 year^{-1} and accordingly total mortality rate (Z) 4.64 year^{-1} (Figure 5).

Table 1

Estimation of growth parameter of the *Priacanthus macracanthus* stock in some areas

Age Character	Location	Year of sampling	K	L _∞	t ₀	φ'	Source
TL-F	Palabuhanratu Bay (Indian Ocean)	2016	1.38	35.7	-0.21	3.25*	Present study
LF	Waters off the west coast of Peninsular Malaysia (Indian Ocean)	1997	1.30	37.0	-	3.25*	(Ahmad et al 2003)
FL-F	North Coast of Java	1980s	1.36	26.7	-	2.99	(Nugroho & Rustam 1983)
TL-F	Java Sea	1978-1979 1977-1978	1.15 1.30	23 23.75	-	2.78* 2.67*	(Dwiponggo et al 1986)
SL-F, O	North-central Taiwan Strait	2006	0.4647	27.379	-0.9140	2.54*	(Ju et al 2016)
LF	Waters off Northeastern Taiwan	1997	0.118	46.8	-1.26	2.41*	(Liu & Cheng 1999)
		1996	0.119	45.4	-1.41	2.39*	
		1995	0.119	46.5	-1.37	2.41*	
		1994	0.139	42.4	-1.17	2.40*	
S, O, V	Surrounding water of Guei-Shan Island, Taiwan	1981-1982	0.113 (M) 0.088 (F)	48.2 (M) 62 (F)	-1.335 (M) -1.05 (F)	2.42* 2.53*	(Joung & Chen 1992)
S	Tungfang waters. Southwestern Taiwan	1981-1982	0.128 (M) 0.136 (F)	40.168 (M) 44.107 (F)	-2.632 (M) -2.900 (F)	2.31* 2.42*	(Liu et al 1992)
LF	Beibu Gulf, (South China Sea)	1997-1999 and 2011-2015	0.70	29.1	-	2.77*	(Zhang et al 2016)
LF	Waters off the east coast of Peninsular Malaysia (South China Sea)	1998	0.56	39.0	-	2.93*	(Ahmad et al 2003)
M	East China Sea	1995-1998	-	40.22 (M) 52.88 (F)	-	-	(Oki & Tabeta 1999)
LF	Brunei Darussalam (South China Sea)	1989-1990	0.95	29.0	-	2.90	(Silvestre & Garces 2004)
LF	North continental shelf of South China Sea	1990s and 1960s	0.60	31.5	-0.44	2.77*	(Sun & Qiu 2004)
FL-F	South China Sea	1965-1967 (all)	0.7	32	1.0	2.86*	(Lester & Watson 1985)
		Spring	1.0	28	1.1	2.89*	
		Autumn	0.7	34	2.1	2.91*	

Note: TL-F – total length frequency; SL-F – standard length frequency; FL-F – fork length frequency; LF – length frequency; M – mesopterygoid; S, O, V – scale, otolith, vertebrae; (M) – male; (F) – female; * – growth performance index (φ') were estimated in this study.




Figure 5. Length converted catch curve of *Priacanthus macracanthus* in Palabuhanratu Bay. The black circles depicted an exploited group suspected to be >1 year old.

The total mortality rate (Z) of *P. macracanthus* in the Indian Ocean, Java Sea, Taiwan Waters, and China Sea ranged from 0.88 to 8.45 year⁻¹ (Table 2).

25

Table 2
Estimation of exploitation and mortality rate of *Priacanthus macracanthus* in some areas

Location	Year of sampling	M	F	Z	E	Source
Palabuhanratu Bay (Indian Ocean)	2016	2.14	2.51	4.64	0.54	Present study
Waters of the west coast of Peninsular Malaysia (Indian Ocean)	1997	2.06	6.39	8.45	0.76	(Ahmad et al 2003)
Java Sea	1978-1979	2.13	2.07	4.20	0.49	(Dwiponggo et al 1986)
	1977-1978	2.28	4.10	6.38	0.64	
North-central Taiwan Strait	2006	1.0218	1.0875	2.1093	0.5156	(Ju et al 2016)
Waters of Northeastern Taiwan	1997	0.33	1.24	1.56		(Liu & Cheng 1999)
	1996	0.33	1.64	1.97		
	1995	0.33	1.03	1.36		
	1994	0.37	0.51	0.88		
Beibu Gulf (South China Sea)	1997-1999 and 2011-2015	1.38	-	4.57	0.70	(Zhang et al 2016)
Waters of the east coast of Peninsular Malaysia (South China Sea)	1998	1.17	5.43	6.60	0.82	(Ahmad et al 2003)
Brunei Darussalam (South China Sea)	1989-1990	1.74	0.34	2.08	0.16	(Silvestre & Garces 2004)
North continental shelf of South China Sea	1990s and 1960s	1.07	1.87	2.94	0.64	(Sun & Qiu 2004)
South China Sea	1967	1.27	0.65	1.92	0.34	(Lester & Watson 1985)
	1966	1.27	0.37	1.64	0.23	
	1965	1.27	1.05	2.32	0.45	

Natural mortality rate (M) was around >0.5 in all areas except in the waters off northeastern Taiwan. The high value of M is due to the rapid growth and short life of the *P. macracanthus*. Differences in M values closely related to life history parameters such as growth rate, length at first maturity, reproduction and maximum age (Vetter 1988). Sparre & Venema (1998) suggested that predation, disease, environmental conditions, stress associated with ecosystems and others can also be determinant factors. The fishing mortality rate (F) during the study period was higher than the natural mortality rate (M). This indicates that mortality of *P. macracanthus* in Palabuhanratu Bay is more influenced by catch, factor of the fisherman. On the other hand, fishing mortality is directly proportional to fishing effort and catch ability, implying that the increase in deaths from capture will be followed by an increase in fishing effort (Atmaja & Nugroho 2004).

Recruitment pattern. Recruitment pattern was apparently occurred throughout the year with two peaks in February-March and August-September. The recruitment level was presumably higher in September compared to March, i.e. 14.66% compared to 11.56% (Figure 6).




Figure 6. Recruitment pattern of *Priacanthus macracanthus* in Palabuhanratu Bay.

P. macracanthus in Palabuhanratu Bay has presumably two recruitment modes (peak and lean recruitment). The major peak occurs during east monsoon season in August/September and the minor ones during west monsoon season in February/March. It was characterized by the finding of many small samples on the peaks and it was assessed that spawning season occurs in the months before the peaks a year. The presence of two recruitment peaks in this study is similar to those reported in the Java Sea (Dwiponggo et al 1986). The occurrence of bimodal recruitment peaks may be related to the rainy season in a region (Pauly & Navaluna 1983; Ingles & Pauly 1984).

Exploitation rate (E) and relative yield per recruit (Y'/R). The Beverton and Holt's Model finding (Figure 7) shows that sustainable yields is achieved at which exploitation rate 0.35, 0.68, and 0.53 for sustainable yield optimum target (E_{50}), maximum sustainable yield (E_{max}), and economic yield (E_{10}), successively. After mortality rate being calculated, the exploitation rate (E_{cur}) was 0.54. As it is shown by Figure 7 below, the current relative yield per recruit (Y'/R_{cur}) 0.04378 which is just 0.0033 less than Y'/R_{max} . The relative yield per recruit (Y'/R) increases steadily until the exploitation rate reaches 0.68 and subsequently decreases with increasing of exploitation.




Figure 7. E values of the *Priacanthus macracanthus* by Y'/R Beverton and Holt $E_{10}=0.55$ (green), $E_{50}=0.35$ (red), $E_{max}=0.68$ (yellow).

The exploitation rate (E) in several areas in Table 2 generally indicates that the species has been over exploited. The current exploitation rate (E_{cur}) of *P. macracanthus* in Palabuhanratu Bay is 0.54 between E_{50} (0.353) with E_{10} (0,553), but closer to E_{10} (Figure 7). The E_{cur} score 0.54 is higher than the optimum ($E_{opt}\approx 0.5$ as in Gulland (1971) and $E_{opt}\approx 0.4$ as in Pauly(1987)). This indicates that the stock of *P. macracanthus* in Palabuhanratu Bay is fully exploited, that means the utilization must be carefully managed where the unit of fishing effort should not be added.

The current exploitation rate (E_{cur}) 0.54 results in Y'/R_{cur} 0.04378, only 0.0033 less than Y'/R_{max} (Figure 7). By increasing the current exploitation from 0.54 to $E_{10} = 0.553$ (2.4% increase in fishing effort) then Y'/R_{cur} will increase from 0.04378 to 0.04426 (1.1% increase). However, if we consider the reference point option $E_{50} = 0.353$ (the exploitation rate) is in which the biomass per recruit is 50% of the virgin stock biomass value at $E = 0$, then the current exploitation should be reduced by 34.6%, leading to a decrease in Y'/R by 23.5%. According to the three options of the biological reference point, it is more optimal to adopt the value obtained from E_{10} (0,553) as the target reference point of *P. macracanthus*'s fisheries management in Palabuhanratu Bay.

Conclusions. The *P. macracanthus* in Palabuhanratu Bay can grow up to asymptotic length (L_{∞}) = 35.7 cm, with growth coefficient (K) of 1.38 year^{-1} and the estimated maximum age of 4.6 years. The natural (M), fishing (F) and total (Z) mortality rates of the *P. macracanthus* were 2.14 year^{-1} , 2.51 year^{-1} , and 4.64 year^{-1} , respectively. The recruitment pattern showed two peaks, in February/March and August/September. The exploitation rate (E) of *P. macracanthus* in Palabuhanratu Bay has reached 0.54. Therefore, it might recommended that the value obtained from E_{10} (0,553) can be used as the target reference point for its management purposes.

Acknowledgements. The authors would like to thank the Head Indonesia Research Institute for Marine Fisheries, Dr. Fayakun Satria, M.App.Sc and person in charge (PIC) for Fisheries Management Area (FMA) 573 in 2016, Anthony Sisco Panggabean, M.Si., for providing the facilities mainly in the beginning of the research. Also, to Ilham, Ph.D., Kadarusman, Ph.D., Mr. Thomas Hidayat and Mr. Muhammad Albab, for their technical help. Our sincere thanks for the valuable discussions to all of them.

References

Ahmad A. T., Isa M. M., Ismail M. S., Yusof S., 2003 Status of demersal fishery resources of Malaysia. In: Assessment, management and future directions for coastal fisheries in Asian Countries. Silvestre G., Garces L., Stobutzki I., Ahmed M.,

- Valmonte-Santos R. A., Luna C., Lachica-Alino L., Munro P., Christensen V., Pauly D. (eds), pp. 83-135, WorldFish Center.
- Atmaja S. B., Nugroho D., 2004 [Characteristics of population parameters of *Amblygaster sirm* and applied models of Beverton and Holt in the Natuna Sea and surrounding areas]. *Journal of Indonesian Fisheries Research* 10:9-15. [In Indonesian, with English abstract].
- Dwiponggo A., Hariati T., Banon S., Palomares M. L., Pauly D., 1986 Growth, mortality and recruitment of commercially important fishes and penaeid shrimps in Indonesian Waters. ICLARM Technical Reports 17. L. J. and Mary C. Skaggs Foundation, Oakland, CA.
- Froese R., Pauly D., 2017 Family Priacanthidae. www.fishbase.org [Internet].
- Gayanilo F. C. J., Sparre P., Pauly D., 2005 FAO-ICLARM Stock Assessment Tools II (FISAT II). Revised Version, User's Guide. Computerized Information Series (Fisheries). Rome: FAO, p. 168.
- Gjedrem T., 2000 Genetic improvement of cold-water fish species. *Aquaculture Research* 31:25-33.
- Gulland J. A., 1971 The fish resources of the oceans. FAO/Fishing News Books Ltd, Surrey, England.
- Ingles J., Pauly D., 1984 An atlas of the growth, mortality and recruitment of Philippine fishes. ICLARM Tech. Rep. 13.
- Joung S. J., Chen C. T., 1992 Age and growth of the big eye *Priacanthus macracanthus* from the surrounding Water of Guei-Shan Island, Taiwan. *Nippon Suisan Gakkaishi* 58:481-488.
- Ju P. L., Yang L., Lu Z. B., Yang S. Y., Du J. G., Zhong H. Q., Chen J., Xiao J. M., Chen M. R., Zhang C. Y., 2016 Age, growth, mortality and population structure of silver croaker *Pennahia argentata* (Houttuyn, 1782) and red bigeye *Priacanthus macracanthus* Cuvier, 1829 in the north-central Taiwan Strait. *Journal of Applied Ichthyology* 32:652-660.
- Lester R. J. G., Watson R. A., 1985 Growth, mortality, parasitism, and potential yields of two Priacanthus species in the South China Sea. *Journal of Fish Biology* 27:307-318.
- Liu K. M., Cheng Y. L., 1999 Virtual population analysis of the big eye *Priacanthus macracanthus* in the waters off northeastern Taiwan. *Fisheries Research* 41:243-254.
- Liu K. M., Chen C. T., Joung S. J., 1992 Some aspects on the fishery biology of big eye *Priacanthus macracanthus* in the Tungkang Waters, Southwestern Taiwan. *Journal of the Fisheries Society of Taiwan* 19:251-262.
- Magnussen E., 2007 Interpopulation comparison of growth patterns of 14 fish species on Faroe Bank: are all fishes on the bank fast-growing? *Journal of Fish Biology* 71:453-475.
- Mehanna S. F., Al-Marzouqi A., El-Siabi B., 2013 Stock characteristics and population dynamics of the spiny cheek grouper *Epinephelus diacanthus* (Valenciennes, 1828) from the Arabian Sea, Oman. *Turkish Journal of Fisheries and Aquatic Sciences* 13:127-132.
- Nugroho D., Rustam R., 1983 [Research on growth and some parameters of fish biology of red bigeye (*Priacanthus macracanthus*) in the North Coast of Java]. Report of Marine Fisheries Research 27:9-14. [In Indonesian, with English abstract].
- Oki D., Tabeta O., 1999 Age and growth of big eye *Priacanthus macracanthus* in the East China Sea. *Fisheries Science* 65:436-440.
- Pardo S. A., Cooper A. B., Dulvy N. K., 2013 Avoiding fishy growth curves. *Methods in Ecology and Evolution* 4:353-360.
- Pauly D., 1983 Some simple methods for the assessment of tropical fish stocks. FAO Fisheries Technical Paper 234, FAO, Rome.
- Pauly D., 1987 A review of the ELEFAN system for analysis of length frequency data in fish and invertebrates. In: Length-based Methods in Fisheries Research. ICLARM Conference Proceedings 13. Pauly D., Morgan G. R. (eds), pp. 7-34. ICLARM, Manila.

- Pauly D., Navaluna N. A., 1983 Monsoon-induced seasonality in the recruitment of philippine fishes. FAO Fisheries Report No. 291, 3:823-833.
- Pauly D., Munro J. L., 1984 Once more on the comparison of growth in fish and invertebrates. Fishbyte. ICLARM Manila.
- Powell A. B., 2000 Preliminary identification of the early life history stages of priacanthid fishes of the Western Central Atlantic. NOAA Technical Memorandum NMFS-SEFSC-439. NOAA Fisheries, Beaufort, NC.
- Ramachandran S., Varghese B. C., 2009 New record of arrowfin bigeye *Priacanthus sagittarius* Starnes, 1988 (Pisces: Priacanthidae) from Indian waters with taxonomic account. Journal of the Marine Biological Association of India 51:217-222.
- Safitri F. E., Arief M., Lamid M., 2016 [Red bigeye (*Priacanthus macracanthus*) surimi solid waste chemical utilization of nutrition content as alternative fish feeding materials]. Journal of Aquaculture and Fish Health 5:1-10. [In Indonesian, with English Abstract].
- Silvestre G. T., Garces L. R., 2004 Population parameters and exploitation rate of demersal fishes in Brunei Darussalam (1989-1990). Fisheries Research 69:73-90.
- Sivakami S., Vivekanandan E., Raje S. G., Shobha J. K., Rajkumar U., 2003 Lizardfishes, pomfrets and bullseye. In: Status of exploited marine fishery resources of India. Joseph M. M., Jayaprakash A. A. (eds), pp. 141-157, Indian Council of Agricultural Research, India.
- Sparre P., Venema S. C., 1998 Introduction to tropical fish stock assessment. Manual FAO Fisheries Technical Paper 306/1, Rev 2. FAO, Rome.
- Starnes W. C., 1999 Priacanthidae. Bigeyes. In: FAO species identification guide for fishery purposes. The living marine resources of the Western Central Pacific. Ref No 40512, Key No. 238. Carpenter K. E., Niem V. H. (eds), pp. 2590-2610, FAO, Rome.
- Sun D. R., Qiu Y. S., 2004 [Estimation of growth and mortality parameters of *Priacanthus macracanthus* in the north continental shelf of South China Sea]. Journal of Zhanjiang Ocean University 24:28-34. [In Chinese, with English abstract].
- Vetter E. F., 1988 Estimation of natural mortality in fish stocks: a review. Fishery Bulletin 86:25-43.
- White W. T., Last P. R., Dharmadi, Faizah R., Chodrijah U., Prisantoso B. I., Pogonoski J. J., Puckridge M., Blaber S. J. M., 2013 Market fishes of Indonesia. ACIAR Monograph No. 155. Australian Centre for International Agricultural Research, Canberra.
- Zhang K., Chen Z., Wang Y., Sun D., Qiu Y., 2016 Population structure of *Priacanthus macracanthus* in the Beibu Gulf, and parameters for its growth, mortality and maturity. Journal of Tropical Oceanography 35:20-28. [In Chinese, with English abstract].
- *** PPNP, 2015 The Annual Statistical Report of Palabuhanratu National Fishing Port 2014. Pelabuhan Perikanan Nusantara (PPN) Palabuhanratu (ID).

Received: 02 September 2017. Accepted: 03 October 2017. Published online: 14 October 2017.

Authors:

Meuthia Aula Jabbar, Jakarta Fisheries University, Department of Aquatic Resources Management, Indonesia, Jakarta 12520, Pasar Minggu-Jakarta Selatan, Jl. AUP No. 1; Bogor Agricultural University, Faculty of Fisheries and Marine Science, Department of Aquatic Resources Management, Indonesia, Bogor 16680, Jl. Lingkar Kampus IPB Dramaga, e-mail: meuthia.aula@gmail.com

Mohammad Mukhlis Kamal, Bogor Agricultural University, Faculty of Fisheries and Marine Science, Department of Aquatic Resources Management, Indonesia, Bogor 16680, Jl. Lingkar Kampus IPB Dramaga, e-mail: m_mukhliskamal@yahoo.com

Menofatria Boer, Bogor Agricultural University, Faculty of Fisheries and Marine Science, Department of Aquatic Resources Management, Indonesia, Bogor 16680, Jl. Lingkar Kampus IPB Dramaga, e-mail: mboer@ymail.com

Ali Suman, Research Institute for Marine Fisheries, Ministry of Marine Affairs and Fisheries, Indonesia, Bogor, Cibinong 16912, Komplek Raiser Ikan Hias Cibinong, Jl. Raya Bogor KM 47 Nanggewer Mekar, e-mail: alisuman_62@yahoo.com

I Nyoman Suyasa, Jakarta Fisheries University, Department of Aquatic Resources Management, Indonesia, Jakarta 12520, Pasar Minggu-Jakarta Selatan, Jl. AUP No. 1, e-mail: soeyasa_stp@hotmail.com

Erfind Nurdin, Research Institute for Marine Fisheries, Ministry of Marine Affairs and Fisheries, Indonesia, Bogor, Cibinong 16912, Komplek Raiser Ikan Hias Cibinong, Jl. Raya Bogor KM 47 Nanggewer Mekar, e-mail: erfind_nurdin@yahoo.co.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Jabbar M. A., Kamal M. M., Boer M., Suman A., Suyasa I. N., Nurdin E., 2017 Population dynamics of the red bigeye (*Priacanthus macracanthus* Cuvier, 1829) (Fish: Family Priacanthidae) in Palabuhanratu Bay, Indonesia. *AAFL Bioflux* 10(5):1198-1209.

Population dynamics of the red bigeye (*Priacanthus macracanthus* Cuvier, 1829) (Fish: Family Priacanthidae)

ORIGINALITY REPORT

13%

SIMILARITY INDEX

9%

INTERNET SOURCES

9%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Institut Pertanian Bogor Student Paper	2%
2	Submitted to CSU, Long Beach Student Paper	<1%
3	Submitted to Vietnam Commercial University Student Paper	<1%
4	myrepositori.pnm.gov.my Internet Source	<1%
5	psasir.upm.edu.my Internet Source	<1%
6	www.fisheries.ubc.ca Internet Source	<1%
7	www.ejfa.me Internet Source	<1%
8	Siddeek, M.S.M.. "Estimation of natural mortality of Kuwait's grooved tiger prawn <i>Penaeus semisulcatus</i> (de Haan) using tag-	<1%

recapture and commercial fisheries data",
Fisheries Research, 199104

Publication

9

Duan, L.J.. "Modeling changes in the coastal ecosystem of the Pearl River Estuary from 1981 to 1998", Ecological Modelling, 20091024

Publication

10

www.scribd.com

Internet Source

11

Ferhat Kabakli, Deniz Erguden. "Age, Growth, and Mortality of the Redcoat Sargocentron rubrum (Forsskal, 1775), in Iskenderun Bay, Northeastern Mediterranean", Thalassas: An International Journal of Marine Sciences, 2021

Publication

12

J.G. Pajuelo, J.M. Lorenzo. "Life history of the red porgy Pagrus pagrus (Teleostei: Sparidae) off the Canary Islands, central east Atlantic", Fisheries Research, 1996

Publication

13

www.uwu.ac.lk

Internet Source

14

Contreras, H.. "Life History of Emerita analoga (Stimpson) (Anomura, Hippidae) in a Sandy Beach of South Central Chile", Estuarine, Coastal and Shelf Science, 199901

Publication

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

15 Silvestre, G.T.. "Population parameters and exploitation rate of demersal fishes in Brunei Darussalam (1989-1990)", Fisheries Research, 200408
Publication

<1 %

16 Ajang, RO, CB Ndome, A Ekwu, EC Uttah, and CI Iboh. "Population dynamics and gillnets selectivity of *Chrysichthys nigrodigitatus* (Lalepede 1803) in lower reaches of the Cross River estuary, Nigeria", Ethiopian Journal of Environmental Studies and Management, 2013.
Publication

<1 %

17 nepadsanbiofishnode.org
Internet Source

<1 %

18 www.alr-journal.org
Internet Source

<1 %

19 www.tandfonline.com
Internet Source

<1 %

20 Ali Suman, Ap'idatul Hasanah, Khairul Amri, Andina Ramadhani Putri Pane, Pratiwi Lestari. "POPULATION CHARACTERISTICS OF MUD CRAB (*Scylla serrata*) IN THE WATERS OF KENDARI BAY AND SURROUNDING AREAS", Indonesian Fisheries Research Journal, 2018
Publication

<1 %

21 E.V. Radhakrishnan, V.D. Deshmukh, Mary K. Manisseri, M. Rajamani, Joe K. Kizhakudan, R. Thangaraja. "Status of the major lobster fisheries in India", New Zealand Journal of Marine and Freshwater Research, 2010
Publication <1 %

22 Eyðfinn Magnussen, Marita D. Magnussen. " Ecology of poor-cod () on the Faroe Bank ", Marine Biology Research, 2009
Publication <1 %

23 Muhammad Abu Bakar Siddik, Md Reaz Chaklader, Md Abu Hanif, Ashfaqun Nahar, Ilham Ilham, Anthony Cole, Ravi Fotedar. "Variation in the life-history traits of a Schilbid catfish, Clupisoma garua (Hamilton, 1822) in the coastal waters of southern Bangladesh", Chinese Journal of Oceanology and Limnology, 2016
Publication <1 %

24 Vladlena V. Gertseva. "The population dynamics of the longnose skate, Raja rhina, in the northeast Pacific Ocean", Fisheries Research, 2009
Publication <1 %

25 citeseerx.ist.psu.edu
Internet Source <1 %

26 dergipark.ulakbim.gov.tr
Internet Source

<1 %

27

drs.cift.res.in

Internet Source

<1 %

28

tdr.lib.ntu.edu.tw

Internet Source

<1 %

29

text-id.123dok.com

Internet Source

<1 %

30

www.fao.org

Internet Source

<1 %

31

www.faoadriamed.org

Internet Source

<1 %

32

www.mdpi.com

Internet Source

<1 %

33

T. Jutagate. "Yield, growth and mortality rate of the Thai river sprat, *Clupeichthys aesarnensis*, in Sirinthorn Reservoir, Thailand", *Fisheries Management and Ecology*, 8/2003

Publication

<1 %

34

Andina Ramadhani Pane. "DINAMIKA POPULASI DAN TINGKAT PEMANFAATAN KEPITING BAKAU (*Scylla serrata* Forskal, 1775) DI PERAIRAN KEPULAUAN ARU DAN SEKITARNYA, MALUKU", BAWAL Widya Riset Perikanan Tangkap, 2019

<1 %

35

Eglė Jakubavičiūtė. "Perch (*Perca fluviatilis*) Growth, Mortality and Stock Exploitation by 40-45 mm Mesh-Sized Gillnet Fishery in the Curonian Lagoon", *Acta Zoologica Lituanica*, 09/01/2011

Publication

<1 %

36

Hyung Kee Cha. "Reproduction and population dynamics of *Penaeus chinensis* (Decapoda: Penaeidae) on the western coast of Korea, Yellow Sea", *Fisheries Research*, 200204

Publication

<1 %

37

M. Njiru. "Are fisheries management, measures in Lake Victoria successful? The case of Nile perch and Nile tilapia fishery", *African Journal of Ecology*, 9/2007

Publication

<1 %

38

Radhakrishnan, Edakkepravan, Mary Manisseri, and Vinay Deshmukh. "Biology and Fishery of the Slipper Lobster, *Thenus orientalis*, in India", *Crustacean Issues*, 2007.

Publication

<1 %