


Review 1



~~The effect of moon phases to the pelagic fish catches purse seine in fisheries management area (FMA) 716, Indonesia~~ The Effect of Moon Phases Upon Purse Seine Pelagic Fish Catches in Fisheries Management Area (FMA) 716, Indonesia

¹Suharyanto, ¹Mohammad K. Arifin, ¹Jerry Hutajulu, ¹Ali Syamsudin Waluyo, ¹Yusrizal, ¹Muhammad Handri, ¹Aman Saputra, ¹Abdul Basith, ¹Erick Nugraha, ²Sepri

¹ Faculty of Fishing Technology, ~~Jakarta Fisheries Institute~~ Jakarta Technical University of Fisheries, South Jakarta, Indonesia; ² Faculty of Fishing Technology, Sorong Fisheries Institute, West Papua, Indonesia. Corresponding author: E. Nugraha, nugraha_eri1@yahoo.co.id

Abstract. The purse seine is a fishing tool whose main part is a net whose target catch is pelagic fish. Many factors influence the amount of catch and one of them is the Moon Phases. Changes in the moon phases can identify best times for fishing operations. The purpose of this study was to analyze the effect of the moon phases on the total catch of *K. pelamis*. The data was collected by following the fishing operation directly. The data analysis method used was descriptive to understand how the purse seine operation process and to classify the number of catches based on 4 moon phases. The results of the analysis of differences in the number of catches in each moon phases showed that the highest number of catches occurred in the first crescent phases 8.575 kg (35%) and the smallest number of catches was during the dark moon phases 1.877 kg (8%). For the results of the analysis of the influence of the moon carried out by the ANOVA test, the value was $0.577 > 0.05$, which means that the moon phases did not have a significant effect on the number of catches.

Key Words: Sulawesi sea, *Pelagic Schooling Species*, Hunter's Moon, Skipjack tuna

Introduction. Indonesia is an archipelagic country that has natural wealth and high fishery resources, both in capture fisheries, marine cultivation, public waters and others (KKP 2017). Fishery resources in North Gorontalo Regency are estimated to have the potential for capture fisheries of 590,970 tons consisting of 175,260 tons of large pelagic fish, 384,750 tons of small pelagic fish, and other types of fish of 30,960 tons. Concerning the utilization rate, it is estimated that capture fishery it has only reached 46% of its potential (Department of Fisheries and Marine Affairs of Gorontalo Province, 2012; 2010). To take advantage of the potential of capture fisheries in North Gorontalo, there are several ways and one of them is using purse seine fishing (Center for Marine and Fisheries Education 2012).

Fishing boats are boats or other floating means that are used for fishing (Fachrussyah 2017). The purse seine boat used for fishing activities in Gorontalo waters uses a "two boat system". The purse seine is a net fishing gear from that is operated by circling a group of fish to a bowl-shaped tool at the end of the fishing process (Salencer 2018). The operation of this fishing gear basically consists of 4 stages of activities which include setting, pursing, hauling and brailing (Santoso & Bawole 2014).

In the operation of a purse seine, there are several factors that affect its operation, one of which is the moon phases. Changes in the moon phases can indicate a good time in fishing operations because there is a difference in light intensity in each moon phases and will affect fish that have positive or negative phototaxis properties of

Comment [A1]: Title suggestion: The effect of moon phases upon purse seine pelagic fish catches in fisheries management area (FMA) 716, Indonesia

Comment [EN2]: Corrected, thank you

Comment [A3]: Please display full name.

Comment [EN4]: Corrected thank you.

Comment [EN5]: I changed our college name. thank you.

Comment [A6]: 590 tones or 590 thousand tones?

Comment [EN7]: 590,970 tons and 384,750 tons

Comment [A8]: Department of Marine Affairs and Fisheries, 2010?

Comment [EN9]: Corrected thank you

Review 1

light so that differences in intensity will affect the volume of the catch when fishermen operate (Jatmiko 2015).

The principle of catching fish with a purse seine is to purse a school of fish with a net. After that the lower net is drained like a bowl, so that the fish is collected in the codend and cannot escape (Syamsuddin et al 2014). The net is operated in the morning starting at 05.00 AM, setting time lasts half an hour. In one trip, the purse seine is operated 1 to 2 times (setting), depending on the catch (Rahmat & Witdiarso 2017).

Fish that are the main purpose of catching from purse seine are fish that are "Pelagic Schooling Species" (~~Gatut Bintoro, 2011~~). (~~Gatut and Sukandar, 2011~~). According to Telussa (2006), fishing operations with purse seine around FADs with small pelagic fish as target species include *Decapterus* spp., *Selaroides leptolepis*, large pelagic fish such as *K. pelamis*), *Auxis* ~~rochei~~ *rochei* and *Thunnus* spp.. A fishing ground is where fish that is the target of fishing are caught in maximum amount and fishing gear can be operated economically as well (Nusantara et al 2014).

According to ~~Gatut Bintoro (2011)~~ *Gatut and Sukandar (2011)* the first step in operating this fishing gear is to find a fishing ground. Because the fish that are targeted by the purse seine are clustered fish living in palagics, generally the catching area is in the form of seas in offshore areas with water depths of about 50 meters or more.

The distribution of demersal fish resources in FMA-716 is relatively narrow covering the coastal areas of Tarakan, Belinyu and Nunukan in East Kalimantan and Likupang Bay and around the Sangihe Talaud islands in North Sulawesi (Suman et al 2014).

Fish catches are multispecies in nature comprising demersal and pelagic species. The Indonesian fisheries administration records the catch divided to eleven statistical areas also called "management areas" (<http://www.fao.org/fishery/facp/IDN/en>). These are shown in Figure 1.




Figure 1. Fisheries Management Areas (FMA) in Indonesia (<http://www.fao.org/fishery/facp/IDN/en>).

Comment [A10]: Gatut & Sukandar 2011?

Comment [EN11]: Corrected. thank you

Comment [A12]: rochei <https://www.fishbase.se/summary/Auxis-rochei.html>

Comment [EN13]: corrected. Thank you

Comment [A14]: Figure must be cited in the paper before it is displayed.

Comment [EN15]: Corrected thank you

Review 1

Moon phases. Optimization of fishing will work well if fishermen can find out the factors that influence it. These factors include suitability in using fishing gear. The fishing gear used should be adjusted to the fishing ground and the type of fish that is the main target. In addition to the suitability of using fishing gear, fish resources will affect the catch obtained. The factor of the phases of the day of the moon will indirectly have an impact on the availability of fish resources, so fishermen need to know the changes in each phases of the day of the moon (Jatmiko 2015).

The catch of fish is strongly influenced by natural factors, one of which is the moon phases. The catch is also affected by changes in the intensity level of the moonlight. Changes in the amount and type of fishermen's catch in each moon phases (dark moon, dark to first crescent, first crescent to light moon, bright moon to last crescent) greatly impact the amount of catch and also the income level of fishermen (Nurlindah et al 2017).

The changing conditions of the moon phases are divided into four phases. New or dark moon phase (new moon), moon phase quadrant 1 (first quarter), full moon phase (full moon), and moon phase quadrant 3 (third quarter) (Figure 2). The phases of change in the conditions of the moon on average occurs every seven days. This division is based on the time or phases of the appearance of the moon. The condition of a bright moon occurs when the appearance of the moon is more than 8 hours in one day, while the moon bright occurs when the appearance of the moon is between 4 hours-7.5 hours, and the dark moon phases occurs when the appearance of the moon only appears between 0 hours - 3.5 hours (Lee 2010).



Figure 2. ~~Siklus bulan wilayah Gorontalo~~ Gorontalo region of moon phase (timeanddate.com/moon/phase/).

Comment [A16]: Please translate into English.

Comment [EN17]: Corrected thank you

The use of light as a fishing aid is closely related to the behavior of fish towards light. In the lift net fishing, the light sources are natural and artificial. Natural light sources come from the sun and moon, when the moon is full, the moonlight will spread over the surface of the water so that the fish will also spread on the surface of the water. This makes it very difficult for fishermen to carry out fishing operations with a purse seine, because it is difficult for fishermen to collect fish into one catchable area. The catch of the boat chart is a group of small pelagic fish that are reactive to light. There are patterns of fish arrival around the light source that go directly to the light source and some are only around the light source. The moon phases are an indication for determining fishing time for fishermen (Siahainenia 2017).

Material and Method. The present research was conducted from November 2019 to May 2020 by participating in the fishing operations of the purse seine vessels operating in North Gorontalo waters. In Figure 3 below we can shown the purse seine vessel used during the study.

Review 1



Figure 3. The purse seine fishing boat used during the study.

The tools and materials needed in this research were camera, calculator, laptop, stationery and Moon Phase Calendar application.

Data collection method. ~~The author-~~ We uses several methods in data collection, including by way of: observation, interviews and documentation.

1. Primary data were obtained by making direct observations on the ship by participating in ship fishing operations, the data includes; (1) preparation, (2) setting, (3) pursing, (4) hauling, (5) brailing, (6) fish handling, (7) date of each moon phases (Table 3).
2. Secondary data were data from fishing owners or companies. The data collection method used was a survey method.

Comment [A18]: Figure must be cited in the paper before it is displayed.

Comment [EN19]: I have added, thank you.

Comment [A20]: Which author? Or the authors?

Comment [EN21]: We/writer. Thank you

Table 3

The emergence of moon phases

Third Quarter		New Moon	
Date	Time	Date	Time
20/11/2019	05:10	26/11/2019	23:05
19/12/2019	07:14	26/12/2019	13:13
17/01/2020	20:58	25/01/2020	05:42
First Quarter		Full Moon	
Date	Time	Date	Time
04/12/2019	14:58	12/12/2019	13:02
03/1/2020	12:45	11/1/2020	03:21
02/2/2020	09:41	09/2/2020	15:33

Data processing. The data that have been obtained during the study were grouped and classified using tables. Data for each trip of fishing activities were grouped according to the catch, type of fish captured, income and the moon phases.

Data analysis. Using descriptive analysis that describes the process of operating fishing gear and to determine the composition of the catch related to the moon phases, a quantitative research was carried out. Furthermore, to observe the effect of the moon phases, the composition of the catch was grouped into four moon phases. To find out the percentage comparison of fishing results between the four moon phases, the approach of Susaniati et al (2013) was applied:

$$p = \frac{n_i}{n} \times 100$$

Where:

p = percentage

n_i = value of catch per species

Review 1

n = value of total catch

$$\text{Percentage of dark moons} = \frac{\text{total new moon catch}}{\text{total moon phases catches}} \times 100$$

$$\text{Percentage of dark moons} = \frac{\text{total first quarter catch}}{\text{total moon phases catches}} \times 100$$

$$\text{Percentage of dark moons} = \frac{\text{total full moon catch}}{\text{total moon phases catches}} \times 100$$

$$\text{Percentage of dark moons} = \frac{\text{total third quarter catch}}{\text{total moon phases catches}} \times 100$$

To find out whether the moon phases affects the number of catches, the authors conducted a One Way Anova test using SPSS 22 software. Before carrying out the One Way Anova test, the normality and homogeneity tests must be passed first. The basis for decision making is as follows:

- Normality test
If the value is Sig. > 0.05, then the data is normally distributed
If the value is Sig. < 0.05, then the data is not normally distributed
- Homogeneity Test
If the value is Sig. > 0.05, then the data is the same or homogeneous
If the value is Sig. < 0.05, then the data is not the same or not homogeneous
- Anova test
If the value is Sig. > 0.05, then the average is equal or has no effect
If the value is Sig. < 0.05, then the average is different or influential

Fishing ground. The fishing operation area at the time of the research was in the Fisheries Management Area of the Republic of Indonesia (FMA-RI) 716, namely in the Sulawesi Sea region (Figure 4).




Figure 4. Fishing ground (Tegoeh et al 2018).

The setting (catching) positions were as it is presented in Table 4.

Table 4

Position at the time of the catching operation		
No	Latitude	Longitude
1	01° 19' 54" U	123° 19' 36" T
2	01° 15' 54" U	122° 28' 18" T

Review 1

3	01°11' 18" U	122°34' 36" T
4	01°22' 42" U	122°39' 12" T
5	01°21' 36" U	122°36' 00" T
6	01°15' 54" U	122°28' 18" T
7	01°21' 48" U	122°36' 42" T
8	01°20' 36" U	122°13' 54" T
9	01°12' 00" U	122°14' 30" T
10	01°12' 48" U	122°35' 48" T

Results

Catched fish. The dominant target fish caught in the waters of North Gorontalo were skipjack tuna (*Katsuwonus pelamis*) *K. pelamis*, *Euthynnus affinis*, *Thunnus albacares*, *Decapterus* sp..

Amount of fish caught during 16 trips. The moon phases are divided into 4 phases, namely the *first quarter*, the *new moon*, the *third quarter* and the *full moon*. The grouping of the amount of catches according to the moon phases is based on the number of trips as many as 16 trips which were then subdivided into 4 moon phases. Figure 5 preents the amount of catch of 16 trips.




Figure 5. Amount of fish catching during 16 trips.

It can be seen that based on the results of the research, the most fish caught was on the 8th trip with the amount of fish of 4,500 kg consisting of *Katsuwonus pelamis* with 1,500 kg, *Decapterus* sp. with 2,000 kg and *Euthynnus affinis* with 1,000 kg.

The total amount of catches was 24,552 kg, with 10,852 kg of *Decapterus* sp., *Euthynnus affinis* 7,850 kg and *K. pelamis* 5,850 kg. It can be seen that the highest catch consisted of *Decapterus* sp. 44% (10,852 kg) and the lowest was recorded for *K. pelamis* 24% (5,850 kg) (Figure 6).

Comment [A22]: After the first mention only the abbreviated Latin name should be displayed.

Comment [EN23]: Corrected thank you

Comment [A24]: Five thousand kg (5 tons) should be displayed as: 5,000 etc.

Comment [EN25]:

Comment [A26]: 4.5 tones or 4.5 kg?

Comment [EN27]: Tones. Thank you

Review 1




Figure 6. Composition of caught fish.

The total number per type of fish catches based on four moon phases can be seen in the graph 7 below:




Figure 7. Amount of catch according species based on four moon phases.

Effect of the moon phases upon the catches amount. Grouping based on moon phases revealed the best results for the *first quarter* of 35% (8.575 kg) and the weakest results 8% (1.877 kg) in the *new moon* phases as shown in Figure 8.




Figure 8. Percentage of fish caught based on four moon phases.

Comment [A28]: Figure must be cited in the paper before it is displayed.

Comment [EN29]: I have added. Thank you

Comment [EN30]: 8 ton, and 1 ton

Review 1

To determine whether or not the moon phases affects the number of catches SPSS software for the *One Way Anova* test was used (Tables 8-10).

- Normality test

Table 8

Normality test

Phase	Kolmogorov-Smirnov ^a			Shapiro-Wilk		
	Statistic	df	Sig.	Statistic	df	Sig.
Amount Third Quarter	.224	3	.	.984	3	.759
New Moon	.318	3	.	.887	3	.344
First Quarter	.268	3	.	.950	3	.571
Full Moon	.276	3	.	.942	3	.537

From the normality test above, it can be concluded that the Sig. >0.05 so that the data is normally distributed and can be continued to the homogeneity test

- Homogeneity Test

Table 9

Homogeneity test

Levene Statistic	df1	Df2	Sig.
1.093	3	8	.406

For the homogeneity test, the Sig. 0.406>0.05 so that it can be said to be homogeneous and it can be continued to the last stage, namely the anova test.

- Anova test

Table 10

Anova test

	Sum of Squares	df	Mean Square	F	Sig.
Between Groups	8601192.667	3	2867064.222	.701	.577
Within Groups	32720469.33	8	4090058.667		
Total	41321662.00	11			

From the ANOVA test results above, it can be seen that the Sig. 0.577> 0.05, which means that the moon phases has a weak or insignificant effect on the amount of catch.

Conclusions

1. From the results of the study it can be concluded that the total number of fish caught was **24,552 kg** with the highest catch consisting in *Decapterus* sp. (**10,852 kg**).
2. The highest amount of catch was obtained during the *first quarter phase*, namely as much as **8,575 kg** (35%).
3. The lowest amount of catch was recorded during the *new moon phases* (1,877 kg; 8%).
4. For the one way ANOVA test results, the Sig. 0.577>0.05, revealed that the moon phases has no significant effect on the amount of catch.

Acknowledgements. We would like to thank to all **skipper** Captain and crew of FB. DOA RESTU 01 who has allowed us to do research on purse seine fishing boat, provided tremendous opportunities, attention and assistance during the research. Thanks also go to all related parties, so that this research can be carried out and published.

References

Comment [A31]: What is a skipper? Please explain.

Comment [EN32]: Skipper, is a captain on fishing boat. Captain/Master is a captain on merchant marine. (IMO rule)

Review 1

- Department of Marine Affairs and Fisheries, 2010. Annual Report. Gorontalo: Department of Marine and Fisheries, North Gorontalo District. 51 p.
- Fachrusyiah Zhulmaydin Chairil. 2017. Textbook of Basics of Fishing. Faculty of Fisheries and Marine Sciences, Gorontalo State University.
- Gatut B., Sukandar. 2011. Fishing Method: bag Net Fishing Equipment. FPIK Universitas Brawijaya.
- Jatmiko, G. G. (2015). Analysis of the Effect of the Moon Day phases on Catch Results and Mini Purse Seine Business Income in PPP Morodemak, Demak [Thesis], Department of Fisheries Resource Utilization, Faculty of Fisheries and Marine Sciences. Bogor Agricultural Institute. Bogor.
- KKP 2017. Decree of the Minister of Marine Affairs and Fisheries No. 50 of 2017. Estimation of Potential, Allowed Catches and Utilization Level of Fish Resources in WPP NRI.
- Lee JW. 2010. The Effect of Day and Moon phases on Catch Yield and Income Level of Lift net Fishermen in Serang District [thesis]. Bogor: Faculty of Fisheries and Marine Sciences, Bogor Agricultural University.
- Nurlindah, A., Kurnia, M., & Nelwan, A. F. 2017. Differences in Boat Chart Production Based on the Moon phases in Barru Regency Waters. Journal of Science and Technology of Fisheries Resource Utilization, 4 (8).
- Nusantara, R. A., Rosyid, A., & Boesono, H. 2014. Analysis of Differences in the Depth of Fishing Areas on Composition of Catches in Cantrang Fishing gear (Boat Seine) in Rembang Waters. Journal of Fisheries Resources Utilization Management and Technology, 3 (3), 96-104.
- Center for Marine and Fisheries Education. 2012. The Purse Seine Fishing Teaching Factory Module.
- Rahmat E., Witdiarso B., 2017 Operation of Mini Purse Seine in Tomini Bay by Fishermen in Gorontalo. Technique Bulletin Resources and Capture Engineering 15(1):31-34.
- Salencer O. C., 2018 Feasibility Study on Fishing with Purse Seine Fishing Equipment at the Port Technical Implementation Unit and Management of Marine and Fisheries Resources (P2SKP) Tambakrejo, Wonotirto District, Blitar Regency, East Java (Doctoral Dissertation, University of Muhammadiyah Malang).
- Santoso H., Bawole F., 2014 Operational Techniques for Purse Seine Fishing Equipment on Northeast Ships 00. Bitung Fishery Academy. Bitung.
- Siahainen R. S., 2017 Capture Chart Results Based on Moon Age in the Waters of Leihtu District, Central Maluku Regency.
- Suman A., Wudianto, Sumiono B., Irianto H. I., Badrudin, Khairul A., 2014 Potential and Level of Fish Resource Utilization in Fisheries Management Areas of the Republic of Indonesia (WPP RI). Publisher: Ref Graphika Jakarta Hal. 149.
- Susaniati W., Nelwan A. F., Kurnia M., 2013 Productivity of Chart Tancap Fishing Areas with Different Distance from Beaches in Jeneponto Regency Waters. Aquatic Journal, 4 (1) ~~xxx~~ p. 68-79.
- Syamsuddin., Mallawa A., Salam A., Koniyo K., 2014 Sustainable and Environmentally Friendly Capture Fisheries Development Strategy in Gorontalo Province.
- Tegoeh N., Thomas H., Umi C., Mufti. P.P. 2018 Reproductive Biology of Spanish Mackerel (*Scomberomorus commerson Lacepede*, 1800) At Kwandang Bay Waters, Sulawesi Sea.
- Telussa R. F., 2006 Effectiveness of Floating Charts in Waai Waters, Ambon Island. [Thesis]. Bogor: Postgraduate School, Bogor Agricultural Institute.
- <http://www.fao.org/fishery/facp/IDN/en>.
- <http://www.timeanddate.com/moon/phase>

Received: September 2020. Accepted: xxxx 2020. Published online: xxxxxxx.

Authors:

Suharyanto, ~~Jakarta Institute of Fisheries~~ Jakarta Technical University of Fisheries, Faculty of Fishing Technology, Jl. AUP Pasar Minggu, South Jakarta, Indonesia, e-mail: suharyanto.jfu@gmail.com

Comment [A33]: Please display pages.

Comment [EN34]: Added. Thank you

Review 1

Mohammad Khaidar Arifin, (Student). ~~Jakarta Institute of Fisheries~~ [Jakarta Technical University of Fisheries, Faculty of Fishing Technology](#). Jl. AUP Pasar Minggu, South Jakarta, Indonesia, e-mail: arifinkhaydar@gmail.com

Jerry Hutajulu, ~~Jakarta Institute of Fisheries~~ [Jakarta Technical University of Fisheries, Faculty of Fishing Technology](#), Jl. AUP Pasar Minggu, South Jakarta, Indonesia, e-mail: jerryhutajulu15@gmail.com

Ali Syamsudin Waluyo, ~~Jakarta Institute of Fisheries~~ [Jakarta Technical University of Fisheries, Faculty of Fishing Technology](#), Jl. AUP Pasar Minggu, South Jakarta, Indonesia, e-mail: aliswaluyo6@gmail.com

Yusrizal, ~~Jakarta Institute of Fisheries~~ [Jakarta Technical University of Fisheries, Faculty of Fishing Technology](#), Jl. AUP Pasar Minggu, South Jakarta, Indonesia, e-mail: buyung_trc@yahoo.co.id

[Muhammad Handri](#), ~~Jakarta Institute of Fisheries~~ [Jakarta Technical University of Fisheries, Faculty of Fishing Technology](#), Jl. AUP Pasar Minggu, South Jakarta, Indonesia, e-mail: handrimuhammad@gmail.com

Aman Saputra, ~~Jakarta Institute of Fisheries~~ [Jakarta Technical University of Fisheries, Faculty of Fishing Technology](#), Jl. AUP Pasar Minggu, South Jakarta, Indonesia, e-mail: amansaputra@yahoo.com

Abdul Basith, ~~Jakarta Institute of Fisheries~~ [Jakarta Technical University of Fisheries, Faculty of Fishing Technology](#), Jl. AUP Pasar Minggu, South Jakarta, Indonesia, e-mail: abdulbasith_ppsstp63@gmail.com

Erick Nugraha, ~~Jakarta Institute of Fisheries~~ [Jakarta Technical University of Fisheries, Faculty of Fishing Technology](#), Jl. AUP Pasar Minggu, South Jakarta, Indonesia, e-mail: nugraha_eriq1@yahoo.co.id

Sepri, Sorong Fisheries Institute, Faculty of Fishing Technology, Jl. K. Pattimura, Tanjung Kasuari, Sorong - West Papua, Indonesia, e-mail: sepri.papua31@gmail.com

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Comment [A35]: Please display full name.