

# Estimation of Biomass and Carbon at Mangroves Tapak in Semarang City

by Dadan Z, Hendra I, Ratna S, Basuki R Meuthia A.j, Shantika Ms, Fitri Damayanti

---

**Submission date:** 03-Nov-2022 04:25AM (UTC-0700)

**Submission ID:** 1943327556

**File name:** IJSR\_2021\_Mangrove\_Tapak\_in\_Semarang\_Indonesia.pdf (420.15K)

**Word count:** 5534

**Character count:** 28282

# Estimation of Biomass and Carbon at Mangroves Tapak in Semarang City, Province of Central Java, Indonesia

Dadan Zulkifli<sup>1</sup>, Hendra Irawan<sup>1</sup>, Ratna Suharti<sup>1</sup>, Basuki Rachmad<sup>1</sup>, Meuthia A. Jabbar<sup>2</sup>, Shantika Maylana Sastraprawira<sup>2</sup>, Fitri Dumayanti<sup>2</sup>

<sup>1</sup>Lecturer of Aquatic Resources Management Technology, Jakarta Technical University of Fisheries, Indonesia

17

<sup>2</sup>IF Sciences de l'14<sup>e</sup> siècle Environment, Université de Bordeaux, France; Plentzia Marine Station, University of the Basque Country, Spain; Institute of Tropical Biodiversity and Sustainable Development, Universiti Malaysia Terengganu, Malaysia

10

<sup>3</sup>Student of Aquatic Resources Management Technology, Jakarta Technical University of Fisheries, Indonesia <sup>4</sup>AUP Ponorogo-Jakarta Selatan 12520  
E-mail: beffyleukq@tritaf@gmail.com

4

**Abstract:** The increasing of greenhouse gases as a result of various human activities that can cause global warming become one of the issues in the world today. The increasing of greenhouse gases causing global warming has a direct impact on climate change and sea level rise. Mangrove ecosystems that are widely available <sup>19</sup> coastal areas have a very effective ability to reduce the concentration of carbon dioxide ( $CO_2$ ) in nature. The objective of study is to determine the potential of carbon stored in mangrove stands in the Tapak Area, Semarang City at Central Java Province, Indonesia. The method used is non-destructive test with allometric equations in mangrove stands and survey. Data collection was done by purposive sampling. The results showed that there were 3 types of mangroves in Dukuh Tapak, namely Rhizophora mucronata, Avicennia marina, and Rhizophora apiculata. Mangrove in Dukuh Tapak has a biomass content of 563.87 tons / ha equivalent to 231.78 tons C / ha with a total of 469 trees.

**Keywords:** biomass, carbon stock, climate change, greenhouse, global warming, mangrove ecosystem, mangrove stands

## 1. Introduction

The coastal area has a strategic meaning because it is an interface area between terrestrial and marine ecosystems (Ward et al., 2020), and has very rich potential for natural resources and environmental services (Susantoro et al., 2019). In this shallow sea area, there are several productive marine ecosystems such as mangroves, estuaries, coral reefs and seagrass beds (Kathiresan, 2011).

The character of the Northern region of Java is a major area in the development of industrial areas and is the main area of land transportation routes on the island of Java (Handayani et al., 2020). High population density and industrial activities can have an impact on increasing the use of natural resources and the volume of waste (Pimentel, 2015; Ray, 2011). All of these activities have an impact on coastal areas which can affect the environmental quality of coastal areas, especially in decreasing the quality of coastal ecosystems (Visbeck, 2013).

Mangrove forests are found in almost all islands in Indonesia in 30 provinces (Daalat et al., 2018). Mangrove forests are a form of ecosystem that plays an important role in the coastal areas of Indonesia (Malik et al., 2019) which are scattered in several provinces in various archipelagic groups (Miron, 2018). The uniqueness of the mangrove ecosystem in Indonesia is that it has the highest species diversity in the world.

Ecological damage to the utilization of mangrove forests in coastal areas that is not managed properly will reduce the function of the mangrove forests which will have a negative impact on the potential and function of other forest ecosystems as habitats (Rudianto et al., 2020). Whereas mangroves can absorb and store large amounts of organic carbon in sediments and biomass of plants, trees and leaves which makes them an important natural <sup>67</sup> carbon sink (Putra, 2019).

Mangrove ecosystems play an important role in global mitigation efforts by reducing  $CO_2$  concentrations (Kibria, 2013). According to (Iksan, 2019) the value of carbon contained in mangrove vegetation is the potential for mangroves to store carbon in the form of biomass. Calculation of carbon stock in a mangrove ecosystem can be used to see the ability of the mangrove ecosystem to absorb gases that can be operated globally (Putra, 2019).

15

Some efforts being made to control the <sup>6</sup> concentration of carbon in the atmosphere can be used to reduce the amount of  $CO_2$  in the atmospheric (Jaikishun, 2017), especially in the Northern region of Java.

## 2. Material and Method

### 2.1 Research Location

The research was conducted in March 2 - May 15, 2020 at Mangrove Tapak, Tegurejo Village, Semarang City Province

Volume 10 Issue 7, July 2021

[www.ijsr.net](http://www.ijsr.net)

Licensed Under Creative Commons Attribution CC BY

of Central Java. The boat lift net size used in this research is 6 GT. Location of the research presented in Figure 1.



Figure 1: Research location

## 2.2 Tools and Material

The tools used in this research are GPS used to determine the coordinate point, camera used to record activities during research, stationery used to record the results of the interview, transect rope used to transect boundary, roll meter used to measure distances between transects, plots rope used to define mangrove boundaries that will be identified in the form of seedlings, saplings and trees, cloth meter used to measure the diameter of mangrove tree trunks, wooden stakes used to be a pole for transect lines and plots, mangrove identification table used to assist in identifying mangrove species in the field, plastic sample used to store leaf, stem and fruit samples in mangroves, mangrove worksheets used to be data entry sheet, compass used to make the rope pulled perpendicular to the shoreline, pH meter used to measure water pH, thermometer used to measure temperature and refractometer used to measure the salinity of waters.

## 2.3 Research Methods

The method used is a non-destructive test with allometric equations method in mangrove stands and also survey method by conducting direct operations in the field. The non-destructive test with allometric equations method is the method done by measuring the height or diameter of the tree and using allometric equations to extrapolate the biomass. The survey method is an investigation carried out to obtain facts from the symptoms that exist and look for factual information, both about social, economic, or political institutions of a group or region in the form of primary and secondary data. Primary data obtained by direct observation to obtain data, conduct interviews to dig up information, in this case the interviewees are mangrove stakeholders. Documenting by taking pictures directly using the camera to support and visualize the results associated with this research. While secondary data is obtained through supporting data obtained through Internet, books and others references. The research station was determined by using a purposive sampling technique. The procedure for measuring biomass and mangrove carbon stock was carried out by means of a non-destructive test (NDT) where the allometric formula was known to be measured. The total biomass in mangrove trees is calculated using allometric equations by calculating the

aboveground biomass. The carbon concentration in organic matter is usually around 50% (Ikhsan et al. 2019), therefore the estimated amount of carbon stored per component can be calculated by multiplying the total weight of the biomass by the carbon concentration. The analysis includes vegetation index, biological index, associated biota, mangrove biomass, carbon and total carbon stock as well as water quality parameters.

## 2.4 Data Analysis

The analysis used in this research is an analysis of species and relative density, species and relative frequencies, importance value index, diversity index, index of evenness, dominance index, environment parameter of waters, calculation of carbon biomass and measurement of total carbon.

- 1) **Species Density (Di)** →  $Di = ni/A \rightarrow (Di = i\text{-species density}; ni = number of individual boxes of species-i; A = total area of sampling)$  (Akhranti, 2020)
- 2) **Relative Species Density (RDi)** →  $RDi = ni/\Sigma n \times 100\% \rightarrow RD_i = Relative\ Density; ni = total\ number\ of\ individuals\ of\ species-i\ (ind); \Sigma n = total\ number\ of\ individuals\ of\ all\ species. (ind)$  (Akhranti, 2020)
- 3) **Species Frequencies (Fi)** →  $Fi = pi/\Sigma P \rightarrow Fi = Frequency\ of\ the\ ith\ type; pi = Number\ of\ sample\ plots\ where\ species\ of\ i,\ is\ found; \Sigma P = The\ total\ number\ of\ sample\ plots\ created$  (Akhranti, 2020)
- 4) **Species Relative Frequency (RFi)** →  $RFi = Fi/\Sigma F \times 100\% \rightarrow RF_i = Relative\ frequency\ (%); Fi = Frequency\ of\ the\ i^{\text{th}}\ species\ (ind); \Sigma F = total\ frequency\ of\ all\ species\ (ind)$  (Akhranti, 2020)
- 5) **Closure of Species (Ci)** →  $Ci = \Sigma BA/A \rightarrow BA = \pi d^2/4, \pi = 3.14; A = The\ total\ area\ of\ the\ sampling\ area\ (m^2)$  (Akhranti, 2020)
- 6) **Relative Closure of Species (RCi)** →  $RCi = Ci/\Sigma C \times 100\% \rightarrow RC_i = closing\ relative\ type; Ci = area\ of\ species\ cover; \Sigma C = total\ area\ of\ closure\ for\ all\ species.$  (Akhranti, 2020)
- 7) **Species Important Value Index (IVI)** →  $IVI = Important\ values\ of\ a\ species\ range\ from\ 0-300\%. This\ important\ value\ provides\ an\ overview\ of\ the\ influence\ or\ role\ of\ a\ type\ of\ mangrove\ in\ the\ mangrove\ community.$  (Akhranti, 2020)
- 8) **Diversity Index (H')** →  $H' = (\Sigma Pi \ln Pi) \rightarrow H' = Shannon-Wiener\ diversity\ index; \Sigma\ is\ summation; Pi\ is\ the\ proportion\ of\ individuals\ found\ in\ the\ ith\ species; \ln = Natural\ logarithm; (Shannon, 1949)$
- 9) **Index of Evenness (E)** →  $E = H'/H'^{\max} \rightarrow E = Evenness\ index; H' = Shannon-Wiener\ diversity\ index; \ln = Natural\ logarithm; S = Number\ of\ species; H'^{\max} = Maximum\ value/\ln S;$  (Odum, 1959)
- 10) **Dominance Index (D)** →  $D = \sum_i ni(ni-1)/N(N-1) \rightarrow D = dominance\ index - Simpson; ni = number\ of\ individual\ in\ the\ i^{\text{th}}\ species; N = total\ number\ of\ entities\ in\ the\ dataset;$  (Simpson, 1949)
- 11) **Water Environment Parameters**

**Table 1:** Standard criteria for aquatic environment parameters (minister of Environment Decree of Indonesia, Number 51/2004)

No.	Water Environment Parameters	Quality Standards
1	Potential Hydrogen (pH)	7-8.5
2	Temperature (°C)	28-32
3	Salinity (‰)	30-34

## 12) Calculation of Carbon Biomass

**Table 2:** Allometric models of above-ground biomass of several mangrove species (Rustam et al., 2015)

No.	Species type	Model of allometric	Source
1	<i>Avicennia alba</i>	$B = 0.07921 * D^{2.7885}$	(Tue et al., 2014)
2	<i>Avicennia marina</i>	$B = 0.1848 * D^{2.534}$	(Dhamawan & Siregar, 2008)
3	<i>Rhizophora apiculata</i>	$B = 0.043 * D^{2.63}$	(Amira, 2008)
4	<i>Rhizophora mucronata</i>	$B = 0.1466 * D^{2.3126}$	(Dhamawan, 2013)
5	<i>Sonneratia alba</i>	$B = 0.3841 * p * D^{2.011}$	(Kauffman & Cole, 2010)
6	<i>Scyphiphora hydrophyllacea</i>	$B = 0.251 * p * D^{2.46}$	(Komiyama et al., 2005)
7	<i>Bruguiera gymnorhiza</i>	$B = p * 0.0754 * D^{2.95}$	(Kauffman & Donato, 2012)
8	<i>Bruguiera cylindrica</i>	$B = 0.351 * p * D^{2.49}$	(Komiyama et al., 2005)
9	<i>Ceriops tagal</i>	$B = 0.251 * p * D^{2.48}$	(Komiyama et al., 2005)
10	<i>Xylocarpus granatum</i>	$B = 0.1832 * D^{2.21}$	(Tadian, 2008)
11	<i>Lumnitzera littorea</i>	$B = 0.251 * p * D^{2.09}$	(Komiyama et al., 2005)
12	<i>Lumnitzera racemosa</i>	$B = 0.251 * p * D^{2.48}$	(Komiyama et al., 2005)

Information:

B = Biomass (kg); D = Diameter at breast height (cm); p = wood density (g/cm³)

## 13) Measurement of Total Carbon Stock

The carbon concentration in organic matter is usually around 50% (Macius, 2017), therefore the estimated amount of carbon stored per component can be calculated by multiplying the total weight of the biomass by the carbon concentration with the following formula (Hetland, 2016): Carbon stock = Biomass per unit area × 0.50

Based on the calculation of carbon stock, the amount of CO<sub>2</sub>

absorption by mangroves will be obtained using the formula: CO<sub>2</sub> absorption = (Mr CO<sub>2</sub>) / (Ar C) × Content C → Mr CO<sub>2</sub> = Molecular weight of the compound (44) and Ar C = Relative molecular weight of atom C (12)

## 3. Result and Discussion

### 3.1 Identification of Mangrove Species

The research which was conducted at the Mangroves of Dukuh Tapak, Tugurejo Village, Semarang City, found 3 species of mangrove trees. The mangrove species consist of Rhizophora m<sup>16</sup> unata, Avicennia marina, and Rhizophora apiculata. It is line with previous research conducted by Martuti et al., 2017). The distribution of mangrove species can be seen in Table 3

**Table 3:** Distribution of mangroves site

Station	Mangrove Types		
	Rhizophora mucronata	Avicennia marina	Rhizophora apiculata
1	121 stands	43 stands	21 stands
2	140 stands	47 stands	-
3	7 stands	90 stands	-
Total	268 stands	180 stands	21 stands

### 3.2 Identification of Mangrove Areas

The area of mangroves in the coastal area of Semarang City is 84.47 ha and the largest area is in the Tugu District which has an area of 52.4 ha of mangrove. This situation tends to show a decrease in the area of mangroves and this is due to the change in land use for industry and the presence of coastal abrasion due to sea level rise (Nugroho et al., 2020). Mangrove in Tapak village itself has an area of mangrove vegetation in 1999 was 20.37 ha, then in 2003 it was 19.12 ha, and in 2015 it was 19.27 ha (Perdina et al., 2016). This mangrove is located on the northern coast of the sea and close to the industrial area of the city of Semarang

### 3.3 Identification of Mangrove Vegetation

The results of observations of mangrove vegetation at station 1 in the mangroves of Dukuh Tapak, Tugurejo Village, Semarang City used an identification book (Noor, 2008). The mangrove vegetation analyzed by calculating the density (D<sub>i</sub>), relative density (DR<sub>i</sub>), relative frequency (FR<sub>i</sub>), closure relative (C<sub>i</sub>), and important value index (IVI) (Ismail et al., 2017) can be shown at figure 2 and 3.




Figure 2: Species relative and density

Based on the figure above, station 1 and 2 has criteria for medium mangrove density with a value of 1001 and 1012 trees / ha respectively, but station 3 has criteria for low mangrove density with a value of 525 trees/ha. There are three

species of mangrove plants at station 1 consisting of Rhizophora mucronata, Rhizophora apiculata and Avicennia marina, two species at station 2 and 3 consisting of Rhizophora mucronata dan Avicennia marina.




Figure 3: Species Relative Frequency and Closure

The highest relative frequency value and relative closure at station 1 was 55.56% and 50% for the species Rhizophora mucronata, indicating that Rhizophora mucronata had the most distribution in each plot. Station 2 has a smaller relative frequency value that is 42.86%, which means that this species of mangroves has less distribution in each area. However, Rhizophora mucronata had a relatively higher cover than Avicennia marina with a ratio of 60%: 40%. So it can be concluded that Rhizophora mucronata has a more important role than Avicennia marina in the area because it has characteristics and morphology that supports it to compete with other species and it can be stated that the condition of the waters in the area is good for mangrove growth. At station 3, Avicennia marina has a relative frequency value of 85.71% and a relative cover of 97.07%. This shows that the Avicennia marina has the most distribution and is evenly distributed in each compartment. Avicennia marina has a relative frequency value of 85.71% and a relative cover of 97.07%. This indicates that Avicennia marina has the most distribution and this species is evenly distributed in each plot.

#### 3.4 Diversity, Evenness and Dominance Index Vegetation

The calculation of the diversity, evenness and dominance index is carried out to determine the level of mangrove biodiversity, the evenness level of a mangrove species and the magnitude of the mangrove species that dominates an area. The results of the calculation of the diversity, evenness and dominance index can be seen in (Table 4).

Table 4: Diversity, evenness and dominance index at research stations at Duku Tapak Mangrove

Index	Station		
	1	2	3
Diversity Index	0.9	0.6	0.3
Evenness Index	0.8	0.8	0.4
Dominance Index	0.5	0.6	0.9

Based on table 4 above, the diversity index value ( $H'$ ) is obtained ranging from 0.3 to 0.9. According to Agustini's

criteria (2016), this value is included in the low category ( $H' \leq 2.0$ ). This shows that the mangrove ecosystem has less productivity, less stable water conditions and high ecological pressure. There are several possible factors causing damage to the mangrove ecosystem, namely pollution, conversion of mangrove forests that do not pay attention to environmental factors and excessive <sup>22</sup>forestation. This is reinforced by the statement of (Ilman et al., 2020) which states that the damage to the mangrove ecosystem is caused by the conversion of functions for fishponds and housing areas, overexploitation and water pollution which triggers degradation of habitat and biodiversity in this area.

Furthermore, the evenness index value at each research station ranged from 0.4 to <sup>21</sup>0.8. In accordance with Agustini's criteria (2016), this value is included in the low category (at station 3) and classified as high (at stations 1 and 2). This shows that the species found at stations 1 and 2 tend to have evenness, it means that there is not particular species dominate a station. If the evenness index value is small, then the <sup>21</sup>evenness in the community is less, it means that the number of individuals of each species is not the same, so there is a tendency to be dominated by certain species. Conversely, the greater the evenness index value indicates that in the community there is no particular dominion species.

The dominance index value at each research station ranged from 0.5 to 0.9. The highest dominance index value was found at station 3 (0.9) and the lowest at station 1 (0.5). Based on Agustini's criteria (2016), station 3 has a high dominance value which is indicated by the presence of a dominant species, namely Avicennia marina which has more than half of the total individuals at station 3. Meanwhile, at station 1 the dominance value is low, there are not mangrove species that dominate the community. This means that the species in the mangrove community tend to be evenness and the ecological conditions are still stable. And for station 2 the dominance value is moderate as indicated by the presence of a dominant species, namely Avicennia marina which has

more than half of the total individuals at the station.

### 3.5 Identification of Biota Types

The biota found at the sampling station were aves species such as large egrets (*Egretta alba*), Prejak jawa (*Prinia familiaris*), and turtles doves (*Geopelia striata*). This biota was seen passing through the air several times when mangrove samples were taken. Then also found types of fish such as gelodok fish (*Periophthalmodon sp.*), Milkfish (*Chanos chanos*) and shrimp (*Litopenaeus vannamei*). Gelodok fish (*Periophthalmodon sp.*) are often found on riverbanks in mangroves. These fish have fast and agile movements in moving places, making it difficult for researchers to catch these fish. Milkfish (*Chanos chanos*) and shrimp (*Litopenaeus vannamei*) are found in mangrove areas located in traditional ponds. At the mangrove location Tapak milkfish and shrimp are commonly cultivated by the local community. At the sampling station, reptiles, molluscs and bivalves were also found, such as snakes (*Hypsiglossus matanensis*), mangrove crabs (*Scylla Spp.*), Snails (*Achatina fulica*), sea snails (*Gibbula divaricata*), tree snails (*Amphidromus*).

### 3.6 Water Environment Parameters

The following is the water quality of the Tapak mangrove sampling station, Tugurejo Village, Semarang City, obtained during the final practice (Table 5).

**Table 5:** Water environment parameters of Tapak's Mangrove

No	Water Environment Parameters	Station 1		Station 2		Station 3		Quality Standards (Kepmen LH/51/2005)
		T1	T2	T1	T2	T1	T2	
1	pH	8.4	8	8.1	8	7.9	7.7	7-8.5
2	Temperature (°C)	32	33	33.3	35.2	32.8	32.5	28-32
3	Salinitas (%)	32	32	30	30.5	27	27	30-34

### 3.7 Water Environment Parameters

Based on the research results, the total value of biomass and carbon stock of mangrove vegetation in Dukuh Tapak, Tugurejo Village, Semarang City was shown in (Table 6).

Station	Mangrove Species	Species Number	Stand Biomass (tonnes/ha)	Carbon Stock (tonnes/ha)	CO2 absorption
1	<i>Rhizophora mucronata</i>	121	60.92	28.02	102.74
	<i>Rhizophora apiculata</i>	21	5.07	2.34	8.56
	<i>Avicennia macrocarpa</i>	43	86.8	39.92	146.4
	$\Sigma$	185	152.79	70.28	257.7
2	<i>Rhizophora mucronata</i>	140	83.14	38.25	140.24
	<i>Avicennia macrocarpa</i>	47	88.14	40.54	148.66
	$\Sigma$	187	171.28	78.29	288.9
3	<i>Rhizophora mucronata</i>	7	3.23	1.49	132.47
	<i>Avicennia macrocarpa</i>	90	176.57	81.22	170.79

Station	Mangrove Species	Species Number	Stand Biomass (tonnes/ha)	Carbon Stock (tonnes/ha)	CO2 absorption
	$\Sigma$	97	179.8	82.71	303.26
	Total Number	469	503.87	231.78	849.86

The research results showed that the biomass content was 503.87 tonnes/ha. This is equivalent to 231.78 tonnes C/ha and this condition is estimated to be able to absorb 849.86 tonnes of CO<sub>2</sub>/ha of carbon dioxide (CO<sub>2</sub>). The Tapak Mangrove has an area of 19.27 ha, so with the existing area it is estimated that the Tapak Mangrove has a biomass content of 9709.57 tonnes/ha, equivalent to 4466.40 tonnes C/ha and it is able to absorb CO<sub>2</sub> about 16376.80 tonnes/ha. Based on carbon storage data obtained from each research station, the highest carbon storage is at station 3 with a total of 97 trees, this allows the area to be able to store carbon of 82.71 tonnes/C, while the lowest carbon content is at station 1 with the number of 185 trees, the ability to store carbon in this area is estimated at 70.28 tonnes/C. Station 3 has the highest biomass potential (179.8 tonnes/ha) while station 1 has the lowest biomass potential (152.78 tonnes/ha). This is because station 3, which is located near the estuary, has a high density and the age of the existing stands is older than others so that the mangroves growing in this location have a relatively large diameter.

### 4. Conclusion

The types of mangroves found in the study locations were different for each station. At station 1 there were 3 types of mangroves, while at stations 2 and 3 only 2 types were found. This condition is possible because each type of mangrove has a different survival rate and substrate.

The quality of mangrove waters in Tapak, Tugurejo Village, Semarang City for the degree of acidity or pH has a value ranging from 7 to 8.4, and the temperature at the observation location has a value ranging from 32°C to 35.2°C. If it is compared with the sea water quality standard, the Ministerial Decree No. 51 of 2004, the pH value indicates the state of the waters in the location has passed the specified threshold. Biota in tropical waters generally lives naturally in the upper limit of the highest temperature, if there is a change from the upper limit it will disrupt physiological processes that can cause the death of biota. The salinity of the waters in the Tapak Semarang mangrove area was found to range from 27‰ to 32‰. If it is compared with the sea water quality standard, the Ministerial Decree No. 51 of 2004, the pH value indicates that the waters in the location are fairly normal.

The mangroves in Tapak, Tugurejo Urban Village, Semarang City are able to absorb carbon from the air. The mangroves in Dukuh Tapak have a biomass content of 503.87 tonnes/ha, a carbon stock content of 231.78 tonnes C/ha with a total of 469 trees.

### 5. Acknowledgment

Authors are thankful to our colleagues at Jakarta Technical University of Fisheries for their assistance in the field works. We are indebted to Ilham, Ph.D as Director of Fisheries Ja-

kura Technical University of Fisheries, Indonesia, for his support to finish this article and we would like also to express our gratitude to our colleagues at Université de Bordeaux, University of the Basque Country and Universiti Malaysia Terengganu, as well as all lecturers of Jakarta Technical University of Fisheries.

## References

- [1] Nicolas D. Ward, J. Patrick Megoigal, Ben Bond-Lamberty, Vanessa L. Bailey, David Butman, Elizabeth A. Canuel, Heida Diefenderfer, Neil K. Ganju, Miguel A. Goñi, Emily B. Grimaldi, Charles S. Hopkinson, Tatung Khunghonkar, J. Adam Langley, Nate G. McDowell, Allison N. Myers-Pigg, Rebecca B. Neumann, Christopher L. Osburn, René M. Price, Joel Rowland, Aditi Sengupta, Marc Simard, Peter E. Thornton, Maria Tzortziou, Rodrigo Vargas, Pamela B. Weisenborn and Lisamarie Windham-Myers, "Representing the function and sensitivity of coastal interfaces in Earth system models," *Nature Communications* Vol. 11 no. 2458, pp. 1-14, 18 Mei 2020, <https://doi.org/10.1038/s41467-020-16236-2>
- [2] Tri Muji Susantoro, Ketut Wikantika, Lissa Fuji Yusman, Alex Tan, M. Firman Ghazali, "Monitoring of Mangrove Growth and Coastal Changes on The North Coast of Brebes, Central Java, Using Landsat Data," *International Journal of Remote Sensing and Earth Sciences* Vol. 16 No.2, pp. 197-214, 26 December 2019, <http://dx.doi.org/10.30536/j.ijreses.2019.v16.a3221>
- [3] Kathiresan Kanasamy and Alikunhi M Nabeel, "Tropical Coastal Ecosystem: Rerally Explored for Their Interaction" *Ecologia* Vol. 1 no. 1, pp. 1-22, 2011, <https://doi.org/10.3923/ecologia.2011.1.22>
- [4] August Daslat, Widodo Setyo Pranowo, Syahrial Nur Amri, "Mangrove Forest Change in Nusa Penida Marine Protected Area, Bali - Indonesia Using Landsat Satellite Imagery", *International Journal of Remote Sensing and Earth Sciences* Vol. 15 no. 2, pp.141-156, December 2018, <https://dx.doi.org/10.30536/j.ijreses.2018.v15.a2955>
- [5] Abdul Malik, Abdul Rahim, Uca Sideng, Abdul Rasyid, and Jamaluddin Jumaddin, "Biodiversity assessment of mangrove vegetation for the sustainability of eco-tourism in West Sulawesi, Indonesia," *AACL Bioflux*, Volume 12, no. 4, pp. 1458-1466, 2019, <http://www.bioflux.com.ro/aacl>
- [6] Alino Miron, "The Archipelagic Status Reconsidered in Light of The South China Sea and Ditzgit Integrity Awards," *Indonesian Journal of International Law*, Vol. 15 no. 3, pp. 306-340, 10 Maret 2018, <http://dx.doi.org/10.17304/ijil.vol15.3.729>
- [7] Golam Kibria, "Mangrove Forests-Its role in Livelihoods, Carbon sinks and Disaster mitigation," Vol. 8 No. 7, pp.1-16, January 2013, <https://doi.org/10.13140/RG.2.1.1848.8800>
- [8] Mahamad Iksan, WOD Al Zarliani , La Nare S. Hafidhawati A. & Fitriani Baena, "Biomass and Carbon Uptake of Mangrove Forests Pohorua Village, Muna Regency," *International Journal of Applied Biology*, Vol. 3 No 2, pp. 57-64, 29 December 2019
- [9] Armya Atlanta Putra, Rudianto and Cira Satrya Utama Dewi, "Analysis of the Ability of Mangrove Sequestration and Carbon Stock in Pejukan Village, Buleleng Regency, Bali," *Jurnal Ilmu dan Teknologi Kelautan Tropis*, Vol. 11 No. 3, Hlm. 511-526, December 2019, <https://doi.org/10.29244/jikkt.v11i3.24049>
- [10] Sirpiol Jaikishun, A.A. Ansari, Philip Dasilva, and Alan Hosen, "Carbon storage potential of mangrove forest in Guyana," *Bonorowo Wetlands*, Vol. 7, no. 1, pp. 43-54 June 2017, <https://doi.org/10.13037/bonorowo/w070109>
- [11] Wiwandari Handayani, Uchendu Eugene Chigbu, Iwan Rudianto and Intan Hapsari Surya Putri, "Urbanization and Increasing Flood Risk in the Northern Coast of Central Java—Indonesia: An Assessment towards Better Land Use Policy and Flood Management," *Land*, Vol. 9, No. 343, pp. 01-22, 23 September 2020, <https://doi.org/10.3390/land9100343>
- [12] David Pimentel, X. Huang, A. Cordova and M. Pimentel, "Impact of a Growing Population on Natural Resources: The Challenge for Environmental Management," *Frontiers*, Vol. 13, No. 12, pp. 01-22, [http://www.frontiersjournal.com/issues/vol3/vol3-12\\_Pimentel.htm](http://www.frontiersjournal.com/issues/vol3/vol3-12_Pimentel.htm)
- [13] Sarbapriya Ray and Ishita Aditya Ray, "Impact of Population Growth on Environmental Degradation: Case of India," *Journal of Economics and Sustainable Development*, Vol.2, No.8, 04 November 2011, <http://www.iiste.org/>
- [14] Martin Visbeck, Ulrike Kronfeld-Goharani, Barbara Neumann, Wilfried Rickels, Jörn Schmidt, and Erik van Doorn, "Establishing a Sustainable Development Goal for Oceans and Coasts to Face the Challenges of our Future Ocean," *Kiel Working Paper*, No. 1847, pp. 01-23, June 2013, <https://www.researchgate.net/publication/248391393>
- [15] Radianto Rudianto, Dietrich G. Bengen and Ferry Karinaawan, "Causes and Effects of Mangrove Ecosystem Damage on Carbon Stocks and Absorption in East Java, Indonesia," *Sustainability Journal*, Vol. 12, No. 24, pp. 01-17, 10 December 2020, <https://doi.org/10.3390/su122410319>
- [16] Mahamad Iksan, W O D Al Zarliani, La Nare, S. Hafidhawati, A, and Fitriani Baena,"Biomass and Carbon Uptake of Mangrove Forests Pohorua Village, Muna Regency, *International Journal of Applied Biology*, Vol. 3, No. 2, pp. 57-64, 2019
- [17] I Akhrianti and A Gustomi, "Important value aspect of mangrove community at coastal area of Pangkalpinang City, Bangka Island", *IOP Conf. Series: Earth and Environmental Science*, Vol. 599, pp. 1-10, 24 November 2020, <https://doi.org/10.1088/1755-1315/599/1/012056>
- [18] C.E. Shannon and W. Weaver, "The mathematical theory of communication", *Science*, Vol. 185, pp. 27-39, 1949
- [19] Odum, E.P., *Introduction to ecology* (translation). Gadjah Mada University Press, Yogyakarta, p. 574, 1993.
- [20] E.H. Simpson, "Measurement of diversity", *Nature*; Vol. 163, p.688
- [21] Minister of Environment Decree of Indonesia, Number 51/2004, Standard criteria for aquatic environment parameters of Minister of Environment, 2004.
- [22] Rustam, Agustin, Novi Sasetyo Adi, Restu Nur Afifati, Terry L. Kepel, Dan August Daslat, T.T. "Blue Car-

- bon : Program Inisiatif Blue Carbon Indonesia Kep. Derawan-Berau, Kalimantan Timur." Diakses 29 April 2020.
- [23] Carlos A. Salas Macias, Julio C. Alegre Orihuela and Sergio Iglesias Abad," Estimation of above-ground live biomass and carbon stocks in different plant formations and in the soil of dry forests of the Ecuadorian coast", Food and Energy Security, pp. 1-7, 28 July 2017.
- [24] Hetland, J., P. Yowargana, S. Ledac, and F. Kraxner. 2016". Carbon-negative emissions: systemic impacts of biomass conversion: a case study on CO<sub>2</sub> capture and storage options", Int. J. Greenhouse Gas Control, Vol. 49, pp. 330–342, June 2016, <https://doi.org/10.1016/j.ijgpc.2016.03.017>
- [25] Nana K. Tri Martuti, Dewi L. Setyowati, Satya B. Nugraha, and Ditha P. Mutintar," Carbon stock potency of mangrove ecosystem at Tapak Sub-village, Semarang, Indonesia", AACL Bioflux, Volume 10, No. 6, pp. 1524-1533, 2017, <http://www.bioflux.com.vn/aac>
- [26] Patric Eric Rakandika Nugroho, Suryanti, and Pudjiono Wahyu Purnomo," Analysis of Changes in Mangrove Area in the North Coast of Central Java Province Indonesia", Indonesian Journal of Fisheries Science and Technology, Vol. 16, no. 3, pp. 208-218, 29 October 2020, <http://ejournal.undip.ac.id/index.php/saintek>
- [27] Perdana, A.F., Petrus Subardjo, and Agus A.D.S., "The study of changes in the area of mangrove vegetation using the NDVI method. Landsat 7 ETM + and Landsat 8 ETM + imagery in 1999, 2003, and 2015 in Pesisir Desu Tapak, Kec. Tuga, Semarang City (translation)," Journal Oceanografi, vol. 5, no. 3, pp. 301-308, 2016, <http://ejournal-s1.undip.ac.id/index.php/josc>
- [28] Noor, Y.S., M. Khazali, & I.N.N. Saryadipuera. Guide Book for Introduction to Mangroves in Indonesia (translation), Ditjen PKA Departemen Kehutanan dan Wetlands International Indonesia Programme. Bogor, 2008.
- [29] Mohd Hasmadi Ismail, Mohd Fodli Ahmad Fuad, Pakhrizad Hassan Zaki, and Noor Janatun Na'im Jemali," Analysis of importance value index of unlogged and logged peat swamp forest in Nenasi Forest Reserve, Peninsular Malaysia", Bosco Rawa Wetlands, Volume 7, No. 2, pp. 74-78, December 2017, <https://doi.org/10.13057/boscorawo/w070203>
- [30] Agustini, Nella Tri, Zamzal Ta'alidin, Dan Dewi Purnama," Mangrove Community Structure in Kahyagu Village, Enggano Island" (translation), Jurnal Enggano, Vol. 1, no. 1, pp. 19–31, 2016.

# Estimation of Biomass and Carbon at Mangroves Tapak in Semarang City

---

ORIGINALITY REPORT



PRIMARY SOURCES

---

- |   |   |        |
|---|---|--------|
| 1 | <a href="http://hdl.handle.net">hdl.handle.net</a><br>Internet Source   | 1 %    |
| 2 | Kauffman J.B., Arifanti V.B., Basuki I.,<br>Kurnianto S., Novita N., Murdiyarno D.,<br>Donato D.C., Warren M.W.. "Protocols for the<br>measurement, monitoring, and reporting of<br>structure, biomass, carbon stocks and<br>greenhouse gas emissions in tropical peat<br>swamp forests", Center for International<br>Forestry Research (CIFOR) and World<br>Agroforestry Centre (ICRAF), 2016<br>Publication | 1 %    |
| 3 | <a href="http://jonuns.com">jonuns.com</a><br>Internet Source   | $<1$ % |
| 4 | <a href="http://lib.unnes.ac.id">lib.unnes.ac.id</a><br>Internet Source   | $<1$ % |
| 5 | <a href="http://www.discoveryjournals.org">www.discoveryjournals.org</a><br>Internet Source   | $<1$ % |
-

- 6 Jeffrey A. Bennett, Mohammad Abotalib, Fu Zhao, Andres F. Clarens. "Life cycle meta-analysis of carbon capture pathways in power plants: Implications for bioenergy with carbon capture and storage", International Journal of Greenhouse Gas Control, 2021 <1 %  
Publication
- 
- 7 Sri Wahyuningsih. "Assessment of Water Quality Status in Bondet Estuary, Cirebon", Tomini Journal of Aquatic Science, 2022 <1 %  
Publication
- 
- 8 Submitted to Universitas Dian Nuswantoro <1 %  
Student Paper
- 
- 9 gss.geo.ugm.ac.id <1 %  
Internet Source
- 
- 10 Ilham, Ravi Fotedar, Sukham Munilkumar. "Effects of organic selenium supplementation on growth, glutathione peroxidase activity and histopathology in juvenile barramundi (*Lates calcarifer* Bloch 1970) fed high lupin meal-based diets", Aquaculture, 2016 <1 %  
Publication
- 
- 11 download.garuda.kemdikbud.go.id <1 %  
Internet Source
- 
- 12 www.hindawi.com <1 %  
Internet Source
-

- 13 Terry Louise Kepel, La Ode Nurman Mbay, R. Bambang Aditya Nugraha, M. Hikmat Jayawiguna, Nasir Sudirman, Peter Mangindaan. "Tekanan Ekologi dan Nilai Moneter Karbon Biru Ekosistem Mangrove Muara Gembong, Teluk Jakarta", Jurnal Kelautan Nasional, 2021  
Publication <1 %
- 14 threatenedtaxa.org Internet Source <1 %
- 15 www.slideshare.net Internet Source <1 %
- 16 Anang Wahyu Sejati, Imam Buchori, Nattaya Mlatti Lakshita, I.G. Andika Wiratmaja, Desyta Ulfiana. "A 50-years flood frequency in Coastal area of Java, Indonesia: Integrating historical hotspot and Spatial Autocorrelation Analysis", Research Square Platform LLC, 2022  
Publication <1 %
- 17 www.mdpi.com Internet Source <1 %
- 18 Yunita Yunita, Fitria Lestari, Yuli Febrianti. "Antibacterial activity lemongrass leaves of Staphylococcus aureus inhibition one", JPBIO (Jurnal Pendidikan Biologi), 2020  
Publication <1 %

- 19 Internet Source <1 %
- 
- 20 [www.ijsh-journals.org](http://www.ijsh-journals.org) <1 %  
Internet Source
- 
- 21 [www.worldscientificnews.com](http://www.worldscientificnews.com) <1 %  
Internet Source
- 
- 22 "Blue Carbon Dynamics of the Indian Ocean", Springer Science and Business Media LLC, 2022 <1 %  
Publication
- 
- 23 I A Syari, J D N Manik, I Akhrianti, A Pamungkas. "Marine debris: Sources, characteristics, and environmental impact on Baturusa River, Bangka Belitung", IOP Conference Series: Earth and Environmental Science, 2021 <1 %  
Publication
- 
- 24 Submitted to University of Birmingham <1 %  
Student Paper
- 
- 25 Nadzirum Mubin, Idham Sakti Harahap, Giyanto ., "DIVERSITY OF SUBTERRANEAN TERMITES (BLATTODEA: TERMITOIDEA) ON VARIOUS TYPES HABITAT AROUND IPB UNIVERSITY CAMPUS, BOGOR, INDONESIA", JURNAL HAMA DAN PENYAKIT TUMBUHAN TROPIKA, 2020 <1 %  
Publication
-

---

Exclude quotes      On

Exclude bibliography    On

Exclude matches      Off