

# The Effect of Organic and Inorganic Phenolic Compounds on Hemoglobin, Hematocrit, and Cortisol Hormone of *Clarias Gariepinus*

<sup>1</sup>Afandi Saputra, <sup>2</sup>Sri Andayani, <sup>2</sup>Happy Nusyam, <sup>1</sup>Djumbuh Rukmono, <sup>1</sup>Maria Goreti Eny, <sup>1</sup>Mugi Mulyono, <sup>1</sup>Sri Budiani, <sup>1</sup>Suharyadi, <sup>1</sup>Erni Marlina, <sup>1</sup>Moch. Nurhudah, <sup>1</sup>Effi A. Thaib, <sup>1</sup>Amyda S. Panjaitan, <sup>1</sup>Sinung Rahardjo, <sup>1</sup>Guntur Prabowo, <sup>1</sup>Ilham, <sup>1</sup>Fitriska Hapsyari, Laishevtcev A.<sup>3,4</sup>

<sup>1</sup> Aquaculture Departement, Jakarta Fisheries University, Indonesia

<sup>2</sup> Faculty of Fisheries and Marine Sciences, Universitas Brawijaya, Indonesia

<sup>3</sup> Federal Scientific Centre VIEV, Moscow, Russia

<sup>4</sup> Laboratory of Biological Control and Antimicrobial Resistance, Orel State University named after I.S. Turgenev, Orel City, Russia

Corresponding Author Email: afandi.saputra@yahoo.co.id

## Abstract.

Phenolic are compounds found in the environment which can cause fish stressed or xenobiotic. In this research, the African catfish (*Clarias gariepinus*) were injected a single dose of phenol (inorganic) and *P.crocatum* Polyphenols (organic). The animals were exposed (75% of 96 hours LC50) with phenol concentration (3 mg/kg) and *P.crocatum* Polyphenols (1.8 mg/kg). The limited information about toxicology and pathological consequences in the fish exposed to phenolic compounds needs to be studied. Furthermore, the hematology was measured on 3 days after injection. The metabolic effects of hemoglobin and hematocrit decreased significantly ( $P < 0.05$ ) while the cortisol hormone increased ( $P < 0.05$ ) in a short-term after 3 days of exposure.

**Key Words:** phenolic, *Clarias gariepinus*, *P. crocatum*, cortisol, hemoglobin

## Introduction.

The phenol residue comes from industrial and pharmaceutical waste as well as and natural decomposition (Buikema *et al.*, 1979). Red Betel (*P. crocatum*) is one of the most aromatic plants in which the leaves contain phenolic compounds (flavonoids and tannin) and have a high phenol content (Saputra *et al.*, 2016). Phenolic compounds become a concern specifically because of its toxic effect in the water environment. However, this compound is difficult to detect due to the lack of taste and smell (Tilak *et al.*, 2007). A cytotoxic activity of red betel leaves on *Artemia* and African catfish was tested in-vivo based on the LC50 and showed that these plants have a potential of cytotoxic compounds. (Nursyam *et al.*, 2017 dan Emrizal *et al.*, 2014)

The habitat of *Clarias gariepinus* is in freshwaters which have a risk to be exposed to chemicals such as phenols (James *et al.*, 2008). Nursyam *et al.*, (2017) and Ibrahim (2012) reported that in-vivo test with an injection of phenolic compounds in fish *C. gariepinus* shows abnormal signs of glucose, BUN, and ALP in the blood and necrosis of liver and kidneys. Even though there is some research on the in-vivo effect of phenolic compounds towards fish blood parameters, there is no information about the residues and effects of organic phenolic compounds from plants to the toxicity on catfish. Therefore, further research is needed to determine the in-vivo influence of phenolic compounds of Red Betel Leaf (*Piper crocatum*) on Hemoglobin, Hematocrit, and Cortisol Hormones of African Catfish (*Clarias gariepinus*).

## Material and Method

**Tested Animals.** The African catfish (*Clarias gariepinus*) used were in the size of 12 to 15 grams. There was 20 fish/tub obtained from the Aquaculture Laboratory of Jakarta Fisheries University. The commercial feed was given twice a day at a dose of 3%. Before

**Comment [U1]:** added AFRICAN CATFISH before the latin name

**Comment [U2]:** second name is placed in front for all

**Comment [U3]:** Rewrite the entire paragraph, because first you should write the purpose of the study, then write the methods you use, and then write the results in a systematic manner. You have written molecular identification result first and then write morphological results. Write properly.

**Comment [U4]:** more precise keywords added

**Comment [U5]:** Introduction is about 300-500 words, covering the aims of the research and provide an adequate background, avoiding a detailed literature survey or a summary of the results. Please add some references to support your research background.

**Comment [U6]:** Not bold

the experiment was carried out, the fish was adapted in a concrete tub for 1 month. The quality of DO water conservancy was 3.99-6.53 mg/l with a pH of 7.0-7.6 at 28-30°C.

#### **Isolation and characterization of Organic and Inorganic phenolic compounds.**

The organic phenolic compounds used were obtained from the isolation of red betel leaf (*P. crocatum*) using maceration, thin layer chromatography, column chromatography, and characterized by using UV-Vis and Infrared spectrophotometers (Saputra *et al.* 2016; Bigoniya and Singh, 2014; Sundang *et al.* 2012; Vasconcelos *et al.* 2010). The inorganic phenolic compounds were obtained from the commercial chemical store.

Comment [U7]: Not bold

**Sub-lethal toxicity study.** The concentration of LC50 from inorganic phenol (3 mg/kg) and organic phenolic (1.80 mg/kg) is equal to 75% of LC50-96 hours. The exposure to phenolic compounds was done for 3 days after a single injection (Saputra *et al.* 2016).

Comment [U8]: Not bold

**Blood sample.** The blood was collected 3 days after exposure to phenolic compounds. The blood was taken by using a 1 ml syringe on the cardinal vein and then centrifuged at 3000 rpm at 4°C for 15 minutes. The serum was separated and stored at a temperature of 15- 20°C (Saputra *et al.* 2016; Varadarajan *et al.*, 2014; Roche and Boge, 2000). The measurement of hemoglobin level was carried out by the Sahli method (Wedemeyer and Yasutake, 1977) in Hb/100ml while the hematocrit level was done by the capillary system (Anderson and Siwicki, 1995). On the other hand, the cortisol concentration plasma (Cort) was assessed by radioimmunoassay (RIA) (Foster and Dunn, 1974).

Comment [U9]: Not bold

**Data analysis.** The data were analyzed by using the Analysis of Variance (ANOVA) with Minitab 16 for Windows.

Comment [U10]: Not bold

## **Results**

Comment [U11]: should be results a discussion

**Hemoglobin.**—The data analysis showed that the hemoglobin after being exposed with phenolic compounds have a significantly different situation in between the 5% level of treatment, Fhit: 35.58 ( $P > 0.05$ ). Based on the Figure 1, the hemoglobin rate of African catfish without treatment (control) is 13.67 Hb/100ml in which it is in line with the research from Alamanda *et al.*, (2007) that a generally healthy African catfish have a hemoglobin level ranged from 12-14 Hb/100 ml of blood.

Different conditions indicate that after the hemoglobin is injected, the phenolic compound was significantly decreased in the treatment of organic phenolic that is 7.67 Hb/100ml. In the inorganic phenolic treatment, it is shown that the hemoglobin was lower compared to the treatment of organic phenolic that is 6.67 Hb/100ml. According to Nusyam *et al.*, (2017) and Wahjuningrum *et al.*, (2008), low and high hemoglobin levels indicate that the fish might be sick or stressed. Roche and Boge (2000) stated that the increase or decrease in the plasma concentration of hemoglobin is a sign that there is an interference from the cells. Harikrishnan *et al.*, (2003) also asserted that the damage in red blood cells causes a decrease in hemoglobin content.




Figure 1. The Hemoglobin of African Catfish after Organic and Inorganic Phenolic Injection

Comment [U12]: for tables made 1 page in the discussion

**Hematocrit:** The results of data analysis showed that the hematocrit condition after being exposed to phenolic compounds was significantly different in between treatments in the level of 5%, Fhit: 17.35 ( $P>0.05$ ). Based on the Figure 2 below, the hematocrit in African catfish without treatment (control) amounts to 37 %. This is in accordance with the opinion of Bastiawan *et al.*, (1995) that the normal hematocrit of African catfish is ranged from 30.8 to 45.5%.

The inorganic phenolic treatment showed that the hematocrit value was much lower than the organic treatment which was 23.33%. This value indicates that the treatment of inorganic phenolic greatly affects the decrease in the African catfish hematocrit. Tsuzuki *et al.*, (2001) stated that when the fish hematocrit is less than 22%, anemia will occur whereas if the hematocrit is greater than 60%, the fish will be in a state of stress. Anderson and Siwicki (1994) confirmed that low hematocrit can indicate the occurrence of fish contamination.




Figure 2. The Hematocrit of African Catfish after Organic and Inorganic Phenolic Injection

Comment [U13]: for tables made 1 page in the discussion

**Cortisol-hormone:** This hormone is the result of endocrine gland secretion. The endocrine glands build the endocrine system which has the function of regulating the work of the body's physiological processes. The value of the cortisol hormone in catfish after being injected with phenolic compounds can be seen in Figure 3.




Figure 3. Cortisol Hormone of African Catfish after Organic and Inorganic Phenolic Injection

Comment [U14]: for tables made 1 page in the discussion

The results showed that the cortisol hormone after being exposed to phenolic compounds was significantly different in between the treatments (5%), Fhit: 3.86 ( $P > 0.05$ ). Based on the results of the study, the cortisol in African catfish without treatment (control) is 412.76 nmol/L. The increase of cortisol hormone in phenolic treatment was compared with the control, in which the increase of the hormone was 198.86% in the organic phenolic treatment and 186.09% in the inorganic phenol treatment. Roche and Boge (2000) reported that the cortisol hormone in fish treatment was increased by 185.7-218%. According to Zaki *et al.*, (2012), there was an increase in *C. gariepinus* fish cortisol after being exposed to phenol for 3 months at 32.7%.

John *et al.*, (1994) explained that the increased cortisol in phenol treatment can be associated with fish poisoning and handling. On the one hand, Roche and Boge (2000) believed that the increase in fish stressors after a single or fraction phenol injection is due to the disturbances in cells. The damage to endocrine cells in both liver and kidneys causes an abnormal biochemical blood value that one of which plays a major role in hemoglobin. Low Hb in tannin injection treatment showed that the spread of oxygen in the body of African catfish was disrupted. Barton (1987) described that a physiological stress will cause a new homeostatic condition in fish through a changing metabolism. This response to stress is controlled by the endocrine system through the release of cortisol hormones and catecholamine.

**Conclusions.** The results from blood tests show that there is an increase in the cortisol hormone while the other parameters such as hemoglobin and hematocrit experience a decrease. This condition explains that the treatment of organic phenolic exposure is comparable to the inorganic phenol exposure. It can be seen from the changes in the blood hemoglobin and hematocrit as well as the cortisol hormones of African catfish.

Comment [U15]: center position and capital letters

## References

Comment [U16]: center position and capital letters

1. Alamanda IE, Handajani NS & Budiharjo, A, Penggunaan metode hematologi dan pengamatan endoparasit darah untuk penetapan kesehatan Ikan Lele Dumbo (*Clarias gariepinus*) di kolam budidaya Desa Mangkubumen Boyolali. *Biodiversitas*, 1 (2017) 34.
2. Anderson DP & Siwicki AK, Basic hematology and serology for fish health programs: Fish Health Section. *Asian Fisheries Society*, 21 (1995) 185.
3. Anderson DP & Siwicki AK, Simplified assays for measuring nonspecific defense mechanisms in fish: Fish Health Section. *Fisheries Soc. Meeting*, 2 (1994) 26.
4. Barton BA, Schreck CB & Barton LD, Effects of chronic cortisol administration and daily acute stress on growth, physiological conditions, and stress responses in juvenile rainbow trout. *Diseases of Aquatic Organisms*, 2 (1987) 173.
5. Bastiawan D, Tauhid M, Alifuddin & Dermawati TS, Perubahan hematologi dan jaringan kan Lele Dumbo (*Clarias gariepinus*) Yang diinfeksi Cendawan *Apharioomyces* sp. *Jurnal Penelitian Perikanan Indonesia*, 22 (1995) 106.
6. Bigoniya P & Singh K, Ulcer protective potential of standardized hesperidin, a citrus flavonoid isolated from *Citrus sinensis*. *Rev Bras Farma*, 24 (2014) 330.
7. Buikema AL, Ginniss MJ & Cairns J, Phenolics in aquatic ecosystems: a selected review of recent literature. *Mar Environ Res*, 2 (1979) 87.
8. Emrizal A, Fernando R, Yuliandri K, Rullah, Nola IRN, Reni A, Susanty R, Yerti F, Ahmad HM, Sirat D, & Arbain, Cytotoxic activities of fractions and two isolated compounds from Sirih Merah (Indonesian red betel), *Piper crocatum* Ruiz and Pav. *Procedia Chemistry*, 13 (2014) 79.
9. Foster LB & Dunn RT, Single antibody technique for radioimmunoassay of cortisol in unextracted serum or plasma. *Clin Chem*, 20 (1974).

10. Harikrishnan R, Rani MN & Balasundaram C, Haematological and biochemical parameters in common carp, *Cyprinus carpio*, following herbal treatment for *Aeromonas hydrophilla* infection. *Aquaculture*, 221 (2003) 41.
11. Ibrahim MD, Experimental exposure of African catfish *Clarias gariepinus* (Burchell, 1822) to phenol: Clinical evaluation, tissue alterations and residue assessment. *Journal of Advanced Research*, 3 (2012) 177.
12. James MO, Stuchal LD & Nyagode BA, Glucuronidation and sulfonation in vitro of the major endocrine-active metabolites of methoxychlor in the channel catfish, *Ictalurus punctatus* and induction following treatment with 3-methylcholanthrene. *Aquatic Toxicology*, 2 (2008) 227.
13. John F, Carragler & Christine MR, Primary and secondary stress responses in golden perch *Macquoria ambigua*. *J. comp. Biochem. Physiol*, 107 (194) 40.
14. Mukherjee D, Bhattacharya S, Kumar V & Moitra J, Biological significance of [<sup>14</sup>C] phenol accumulation in different organs of a murrel, *Channa punctatus*, and the common carp, *Cyprinus carpio*. *Biomed. Environ. Sci*, 3 (1990) 337.
15. Nursyam H, Andayani S & Saputra A, The effect of tannin from red betel (*Piper crocatum*) leaves towards blood biochemistry and histology of North African catfish (*Clarias gariepinus*). *AAFL Bioflux*, 10 (2017) 1386.
16. Roche H & Boge G, Fish blood parameters as a potential tool for identification of stress caused by environmental factors or chemical intoxication. *Mar. Environ. Res*, 125 (2000) 27.
17. Saputra A, Andayani S & Nusyam H, Total quantity of phenol and isolation methanol tannin extract of red betel leaf (*Piper crocatum*). *International Journal of PharmTech Research*, 9 (2016) 146.
18. Sundang M, Nasir SNS, Sipaut CS & Othman H. (2012), Antioxidant activity, phenolic, flavonoid and tannin content of *Piper betle* and *Leucosyke capitella*. *Malaysian Journal of Fundamental and Applied Sciences*, 1 (2012) 1.
19. Taysse L, Troutaud D, Khan NA & Deschaux P, Structure activity relationship of phenolic compounds (phenol, pyrocatechol and hydroquinone) on natural lymphocytotoxicity of carp (*Cyprinus carpio*). *Toxicology*, 1 (1995) 207.
20. Tilak KS, Veeraiah K, Butchiram MS, Effect of phenol on haematological components of Indian major carps *Catla catla*, *Labeo rohita* and *Cirrhinus mrigala*. *J Environ Biol*, 2 (2007) 177.
21. Tsuzuki MY, Ogawa K & Strussmann CA, Physiological responses after stress and subsequent recovery at different salinities in adults pejerrey *Odontesthes bonariensis*. *Aquaculture*, 200 (2001) 349.
22. Vasconcelos PCP, Andreo MA, Vilegas W, Hiruma-Lima CA & Pellizzon CH, Effect of *Mouriri pusa* tannins and flavonoids on prevention and treatment against experimental gastric ulcer. *Journal of Ethnopharmacology*, 131 (2010) 146.
23. Varadarajan R, Sankar H, Jose J & Philip B, Sublethal effects of phenolic compounds on biochemical, histological and ionoregulatory parameters in a tropical teleost fish *Oreochromis mossambicus* (Peters). *International Journal of Scientific and Research Publications*, 3 (2014) 2250.
24. Wahjuningrum D, Ashry N & Suryati S, Pemanfaatan ekstrak daun Ketapang *Terminalia Cattapa* untuk pencegahan dan pengobatan Ikan Patin (*Pangasionodon hypophthalmus*) yang Terinfeksi (*Aeromonas hydrophila*). *Jurnal Akuakultur Indonesia*, 7 (2008) 82.
25. Wedemeyer GA & Yasutake WT, *Clinical methods for the assessment of the effects of environmental stress on fish health* (Department of the Interior Fish and Wildlife Service, USA), 1997, 80.
26. Zaki MS, El-Batrawy N & Taha NM, Dose phenol toxicity affected endocrine status in African Catfish (*Clarias gariepinus*). *Life Science Journal*, 9 (2012) 636.

Comment [U17]: give numbering

Authors:

Afandi Saputra, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [afandi.saputra@yahoo.co.id](mailto:afandi.saputra@yahoo.co.id).

Sri Andayani, Faculty of Fisheries and Marine Sciences, Universitas Brawijaya, Jl. Veteran, Malang, Indonesia, e-mail: [srianday\\_08@yahoo.com](mailto:srianday_08@yahoo.com)  
Happy Nussyam, Faculty of Fisheries and Marine Sciences, Universitas Brawijaya, Jl. Veteran, Malang, Indonesia, e-mail: [happy\\_nursyam@yahoo.com](mailto:happy_nursyam@yahoo.com)  
Djumbuh Rukmono, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [d\\_rukmono@yahoo.co.id](mailto:d_rukmono@yahoo.co.id)  
Maria Goreti Eny, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [eny.kristiany@gmail.com](mailto:eny.kristiany@gmail.com)  
Mugi Mulyono, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [mugimulyono@kkp.go.id](mailto:mugimulyono@kkp.go.id)  
Sri Budiani, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [s.budiani@yahoo.com](mailto:s.budiani@yahoo.com)  
Suharyadi, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [suharyadi.fahri@gmail.com](mailto:suharyadi.fahri@gmail.com)  
Erni Marlina, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [ernie.marlina@gmail.com](mailto:ernie.marlina@gmail.com)  
Moch. Nurhudah, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [mochnurhudah@kkp.go.id](mailto:mochnurhudah@kkp.go.id)  
Effi A. Thaib, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [effithaib@gmail.com](mailto:effithaib@gmail.com)  
Amyda S. Panjaitan, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [amyda\\_3r@yahoo.com](mailto:amyda_3r@yahoo.com)  
Sinung Rahardjo, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [sinungrahardjo@yahoo.com](mailto:sinungrahardjo@yahoo.com)  
Guntur Prabowo, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [dhgunturprabowo@yahoo.com](mailto:dhgunturprabowo@yahoo.com)  
Ilham, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [ilham.jfu@kkp.go.id](mailto:ilham.jfu@kkp.go.id)  
Fitriska Hapsyari, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [hapsari\\_fitriska@yahoo.com](mailto:hapsari_fitriska@yahoo.com)



# RUSSIAN JOURNAL OF AGRICULTURAL AND SOCIO- ECONOMIC SCIENCES

ISSN 2226-1184



## MANUSCRIPT ACCEPTANCE CERTIFICATE

### **Title of Articles/Manuscripts:**

**“ EFFECT OF ORGANIC AND INORGANIC PHENOLIC COMPOUNDS ON HEMOGLOBIN, HEMATOCRIT, AND CORTISOL HORMONE OF AFRICAN CATFISH (*Clarias gariepinus*)”**

### **By**

**Afandi Saputra<sup>1\*</sup>, Sri Andayani<sup>2</sup>, Happy Nusyam<sup>2</sup>, Djumbuh Rukmono<sup>1</sup>, Maria Goreti Eny<sup>1</sup>, Mugi Mulyono<sup>1</sup>, Sri Budiani<sup>1</sup>, Suharyadi<sup>1</sup>, Erni Marlina<sup>1</sup>, Moch. Nurhudah<sup>1</sup>, Effi A. Thaib<sup>1</sup>, Amyda S. Panjaitan<sup>1</sup>, Sinung Rahardjo<sup>1</sup>, Guntur Prabowo<sup>1</sup>, Ilham<sup>1</sup>, Fitriska Hapsyari<sup>1</sup>, Laishevtcev A.<sup>3,4</sup>**

### **Institution:**

<sup>1</sup> Aquaculture Departement, Jakarta Fisheries University

<sup>2</sup> Fac. Of Fisheries and Marine Science. Universitas Brawijaya, Malang, Indonesia

<sup>3</sup>Federal Scientific Centre VIEV, Moscow, Russia

<sup>4</sup> Laboratory of Biological Control and Antimicrobial Resistance, Orel State University named after I.S. Turgenev, Orel City, Russia

**Category:** Research article

I, hereby acknowledge that the above article/manuscript has been received and accepted for publication in **ISSUE 2(96), Desember 2019 RJOAS**

**Date:** 20/11/2019

**Sergey Plygun**

(Editor in-chief)

#### International bibliographic systems and databases:

- [AGRIS: International Information System for the Agricultural Sciences and Technology](#)
- [DOAJ: Directory of Open Access Journals](#)
- [CAB Abstracts](#)
- [Open Access Infrastructure for Research in Europe \(OpenAIRE\)](#)
- [Research Papers in Economics \(RePEc\)](#)
- [Bielefeld Academic Search Engine \(BASE\)](#)
- [EBSCOhost](#)
- [Microsoft Academic](#)
- [Dimensions](#)
- [Mendeley](#)

#### Russian bibliographic systems and databases:

- [Cyberleninka](#)
- [ELS of Publishing House "Lan"](#)
- [ELS of Publishing House "Infra-M"](#)
- [ELS "IPRbooks"](#)
- [GlobalF5](#)
- [SocioNet](#)





---

## Manuscript Submission – Afandi Saputra

6 messages

---

**Afandi Saputra** <[afandi.saputra@yahoo.co.id](mailto:afandi.saputra@yahoo.co.id)>

Oct 14, 2019 at 7:49 PM

To: [rjoas\[a\]yandex.ru](mailto:rjoas[a]yandex.ru)

Dear Editor,

I hereby send you a manuscript for possible publication in your journal. We would like to publish the article " THE EFFECT OF ORGANIC AND INORGANIC PHENOLIC COMPOUNDS ON HEMOGLOBIN, HEMATOCRIT, AND CORTISOL HORMONE OF AFRICAN CATFISH (CLARIAS GARIEPINUS) "Thank you :)

Regards,  
Afandi Saputra

---

 **Manuscript - Saputra et al., SUBMIT.docx**  
133 Kb

---

**Plygun** <[rjoas\[a\]yandex.ru](mailto:rjoas[a]yandex.ru)>

Oct 23, 2019 at 9:51 AM

To: [afandi.saputra@yahoo.co.id](mailto:afandi.saputra@yahoo.co.id)

Reviewer A

Dear **Mr. Afandi Saputra**

Evaluation process of the above mentioned manuscript has been reviewed. The comments of the reviewer(s) are included at the bottom of this letter.

The reviewer(s) have recommended publication, but also suggest some revisions to your manuscript. Therefore, I invite you to respond to the reviewer(s) comments and revise your manuscript within the period of defined time.

Because we are trying to facilitate timely publication of manuscripts submitted to journal, your revised manuscript should be uploaded as soon as possible. If it is not possible for you to submit your revision in a reasonable amount of time, we may have to consider your paper as a new submission.

Once again, thank you for submitting your manuscript to this journal and I look forward to receiving your revision.

Truly yours,

Recommendation: Revisions Required

Thanks and I look forward to your update.

Sincerely,

Editorial Team

[Quoted text hidden]

---

**1 attachments**

 **11411 Review editor dan riviewers.pdf**  
657 Kb



Afandi S <[afandi.saputra@yahoo.co.id](mailto:afandi.saputra@yahoo.co.id)>

---

## Manuscript Submission – Afandi Saputra

6 messages

---

**Afandi Saputra** <[afandi.saputra@yahoo.co.id](mailto:afandi.saputra@yahoo.co.id)>

Oct 29, 2019 at 11:55 PM

To: [rjoas\[a\]yandex.ru](mailto:rjoas[a]yandex.ru)

Dear RJOAS editor, I have corrected the reviewers, herewith I am attaching the repair document

Regards,  
Afandi Saputra

---

 **Manuscript - Saputra et al., Revision.docx**  
3.840 Kb

---

**Plygun** <[rjoas\[a\]yandex.ru](mailto:rjoas[a]yandex.ru)>

Nov 5, 2019 at 6:34 AM

To: [afandi.saputra@yahoo.co.id](mailto:afandi.saputra@yahoo.co.id)

Dear Afandi Saputra,

We would like to publish the article " THE EFFECT OF ORGANIC AND INORGANIC PHENOLIC COMPOUNDS ON HEMOGLOBIN, HEMATOCRIT, AND CORTISOL HORMONE OF AFRICAN CATFISH (CLARIAS GARIEPINUS)". Please find the attached file for further information regarding how to publish your manuscript, including the acceptance letter.

Thanks and I look forward to your update.

Sincerely,

Editorial Team

[Quoted text hidden]

---

### 1 attachments

 **Publication\_PackageInstruction.pdf**  
195 Kb

---



---

## Manuscript Submission – Afandi Saputra

6 messages

---

**Afandi Saputra** <[afandi.saputra@yahoo.co.id](mailto:afandi.saputra@yahoo.co.id)>

Nov 16, 2019 at 11:19 PM

To: [rjoas\[a\]yandex.ru](mailto:rjoas[a]yandex.ru)

Dear Editor RJOAS,

Here is the payment receipt. Thank you :)

---

**You paid 225 \$ USD**

to POD SERVICE LIMITED

[Details](#)

### Purchase details

Receipt number: 122Q34567LT8827768

We'll send confirmation to: [afandi.saputra@yahoo.co.id](mailto:afandi.saputra@yahoo.co.id)

### Merchant details

POD SERVICE LIMITED

Regards,  
Afandi SaputraA

[Quoted text hidden]

---

**Plygun** <[rjoas\[a\]yandex.ru](mailto:rjoas[a]yandex.ru)>

Nov 20, 2019 at 10:13 AM

To: [afandi.saputra@yahoo.co.id](mailto:afandi.saputra@yahoo.co.id)

Dear authors,

Thanks for the documents.

To be published on December 12 (96). Please keep an eye on the journal's homepage. Thank you for publishing with us.

Editor

[Quoted text hidden]

---

### 1 attachments

 **Acceptance letter – RJOAS Dec 12(96).pdf**  
479Kb

**THE EFFECT OF ORGANIC AND INORGANIC PHENOLIC COMPOUNDS  
ON HEMOGLOBIN, HEMATOCRIT, AND CORTISOL HORMONE  
OF AFRICAN CATFISH (*CLARIAS GARIEPINUS*)**

**Saputra Afandi<sup>1\*</sup>, Andayani Sri<sup>2</sup>, Nusyam Happy<sup>2</sup>, Rukmono Djumbuh<sup>1</sup>,  
Eny Maria Goreti<sup>1</sup>, Mulyono Mugi<sup>1</sup>, Budiani Sri<sup>1</sup>, Suharyadi<sup>1</sup>, Marlina Erni<sup>1</sup>,  
Nurhudah M.<sup>1</sup>, Thaib Effi A.<sup>1</sup>, Panjaitan Amyda S.<sup>1</sup>, Rahardjo Sinung<sup>1</sup>,  
Prabowo Guntur<sup>1</sup>, Ilham<sup>1</sup>, Hapsyari Fitriska<sup>1</sup>, Laishevtcev A.<sup>3,4</sup>**

<sup>1</sup>Department of Aquaculture, Jakarta Fisheries University, Indonesia

<sup>2</sup>Faculty of Fisheries and Marine Science, University of Brawijaya, Malang, Indonesia

<sup>3</sup>Federal Scientific Centre VIEV, Moscow, Russia

<sup>4</sup>Laboratory of Biological Control and Antimicrobial Resistance,  
Orel State University named after I.S. Turgenev, Orel City, Russia

\*E-mail: [afandi.saputra@yahoo.co.id](mailto:afandi.saputra@yahoo.co.id)

**ABSTRACT**

Phenolic are compounds found in the environment which can cause fish stressed or xenobiotic. The limited information about toxicology and pathological consequences in the fish exposed to phenolic compounds needs to be studied. In this research, the African catfish (*Clarias gariepinus*) were injected a single dose of phenol (inorganic) and *P.crocatum* Polyphenols (organic). The animals were exposed (75% of 96 hours LC50) with phenol concentration (3 mg/kg) and *P.crocatum* Polyphenols (1.8 mg/kg). Furthermore, the hematology was measured on 3 days after injection. The metabolic effects of hemoglobin and hematocrit decreased significantly ( $P < 0.05$ ) while the cortisol hormone increased ( $P < 0.05$ ) in a short-term after 3 days of exposure.

**KEY WORDS**

Phenolic, *Clarias gariepinus*, *P. crocatum polyphenols*, cortisol, hemoglobin, hematocrit.

Phenolic compounds are xenobiotic compounds that affect anemia and disrupt the endocrine system and hormones (Writer *et al.*, 2010; Hori *et al.*, 2006; Awachie and Ezenwaji, 1998). This compound is classified as toxic chronic and immunotoxic which usually found in fish tissue (Taysse *et al.*, 1995; Mukherjee *et al.*, 1990). Phenol waste is accumulated from human activities such as the waste from industry, pharmaceuticals, and agriculture (Buikema *et al.*, 1979).

The phenol residue comes from industrial and pharmaceutical waste as well as and natural decomposition (Buikema *et al.*, 1979). Red Betel (*P. crocatum*) is one of the most aromatic plants in which the leaves contain phenolic compounds (flavonoids and tannin) and have a high phenol content (Saputra *et al.*, 2016). Phenolic compounds become a concern specifically because of its toxic effect in the water environment. However, this compound is difficult to detect due to the lack of taste and smell (Tilak *et al.*, 2007). A cytotoxic activity of red betel leaves on *Artemia* and African catfish was tested in-vivo based on the LC50 and showed that these plants have a potential of cytotoxic compounds. (Nursyam *et al.*, 2017 and Emrizal *et al.*, 2014)

The habitat of *Clarias gariepinus* is in freshwaters which have a risk to be exposed to chemicals such as phenols (James *et al.*, 2008). Nursyam *et al.*, (2017) and Ibrahim (2012) reported that in-vivo test with an injection of phenolic compounds in fish *C. gariepinus* shows abnormal signs of glucose, BUN, and ALP in the blood and necrosis of liver and kidneys. Even though there is some research on the in-vivo effect of phenolic compounds towards fish blood parameters, there is no information about the residues and effects of organic phenolic compounds from plants to the toxicity on catfish. Therefore, further research is needed to

determine the in-vivo influence of phenolic compounds of Red Betel Leaf (*Piper crocatum*) on Hemoglobin, Hematocrit, and Cortisol Hormones of African Catfish (*Clarias gariepinus*).

## MATERIALS AND METHODS OF RESEARCH

*Tested Animal.* The African catfish (*Clarias gariepinus*) used were in the size of 12 to 15 grams. There was 20 fish/tub obtained from the Aquaculture Laboratory of Jakarta Fisheries University. The commercial feed was given twice a day at a dose of 3%. Before the experiment was carried out, the fish was adapted in a concrete tub for 1 month. The quality of DO water conservancy was 3.99-6.53 mg/l with a pH of 7.0-7.6 at 28-30°C.

*Isolation and characterization of Organic and Inorganic phenolic compounds.* The organic phenolic compounds used were obtained from the isolation of red betel leaf (*P. crocatum*) using maceration, thin layer chromatography, column chromatography, and characterized by using UV-Vis and Infrared spectrophotometers (Saputra *et al.* 2016; Bigoniya and Singh, 2014; Sundang *et al.* 2012; Vasconcelos *et al.* 2010). The inorganic phenolic compounds were obtained from the commercial chemical store.

*Sub-lethal toxicity study.* The concentration of LC50 from inorganic phenol (3 mg/kg) and organic phenolic (1.80 mg/kg) is equal to 75% of LC50-96 hours. The exposure to phenolic compounds was done for 3 days after a single injection (Saputra *et al.* 2016)

*Blood sample.* The blood was collected 3 days after exposure to phenolic compounds. The blood was taken by using a 1 ml syringe on the cardinal vein and then centrifuged at 3000 rpm at 4°C for 15 minutes. The serum was separated and stored at a temperature of 15- 20°C (Saputra *et al.* 2016; Varadarajan *et al.*, 2014; Roche and Boge, 2000). The measurement of hemoglobin level was carried out by the Sahli method (Wedemeyer and Yasutake, 1977) in Hb/100ml while the hematocrit level was done by the capillary system (Anderson and Siwicki, 1995). On the other hand, the cortisol concentration plasma (Cort) was assessed by radioimmunoassay (RIA) (Foster and Dunn, 1974).

*Data analysis.* The data were analyzed by using the Analysis of Variance (ANOVA) with Minitab 16 for windows.

## RESULTS AND DISCUSSION

The data analysis showed that the hemoglobin after being exposed with phenolic compounds have a significantly different situation in between the 5% level of treatment, Fhit: 35.58 (P> 0.05). Based on the Figure 1, the hemoglobin rate of African catfish without treatment (control) is 13.67 Hb/100ml in which it is in line with the research from Alamanda *et al.*, (2007) that a generally healthy African catfish have a hemoglobin level ranged from 12-14 Hb/100 ml of blood.

Different conditions indicate that after the hemoglobin is injected, the phenolic compound was significantly decreased in the treatment of organic phenolic that is 7.67 Hb/100ml. In the inorganic phenolic treatment, it is shown that the hemoglobin was lower compared to the treatment of organic phenolic that is 6.67 Hb/100ml. According to Nusyam *et al.*, (2017) and Wahjuningrum *et al.*, (2008), low and high hemoglobin levels indicate that the fish might be sick or stressed. Roche and Boge (2000) stated that the increase or decrease in the plasma concentration of hemoglobin is a sign that there is an interference from the cells. Harikrishnan *et al.*, (2003) also asserted that the damage in red blood cells causes a decrease in hemoglobin content.

The results of data analysis showed that the hematocrit condition after being exposed to phenolic compounds was significantly different in between treatments in the level of 5%, Fhit: 17.35 (P>0.05). Based on the Figure 2 below, the hematocrit in African catfish without treatment (control) amounts to 37 %. This is in accordance with the opinion of Bastiawan *et al.*, (1995) that the normal hematocrit of African catfish is ranged from 30.8 to 45.5%.

The inorganic phenolic treatment showed that the hematocrit value was much lower than the organic treatment which was 23.33%. This value indicates that the treatment of inorganic phenolic greatly affects the decrease in the African catfish hematocrit. Tsuzuki *et*

*al.*, (2001) stated that when the fish hematocrit is less than 22%, anemia will occur whereas if the hematocrit is greater than 60%, the fish will be in a state of stress. Anderson and Siwicki (1994) confirmed that low hematocrit can indicate the occurrence of fish contamination.




Figure 1 – The Hemoglobin of African Catfish after Organic and Inorganic Phenolic Injection




Figure 2 – The Hematocrit of African Catfish after Organic and Inorganic Phenolic Injection




Figure 3 – Cortisol Hormone of African Catfish after Organic and Inorganic Phenolic Injection

This hormone is the result of endocrine gland secretion. The endocrine glands build the endocrine system which has the function of regulating the work of the body's physiological

processes. The value of the cortisol hormone in catfish after being injected with phenolic compounds can be seen in Figure 3.

The results showed that the cortisol hormone after being exposed to phenolic compounds was significantly different in between the treatments (5%), Fhit: 3.86 ( $P > 0.05$ ). Based on the results of the study, the cortisol in African catfish without treatment (control) is 412.76 nmol/L. The increase of cortisol hormone in phenolic treatment was compared with the control, in which the increase of the hormone was 198.86% in the organic phenolic treatment and 186.09% in the inorganic phenol treatment. Roche and Boge (2000) reported that the cortisol hormone in fish treatment was increased by 185.7-218%. According to Zaki *et al.*, (2012), there was an increase in *C.gariepinus* fish cortisol after being exposed to phenol for 3 months at 32.7%.

John *et al.*, (1994) explained that the increased cortisol in phenol treatment can be associated with fish poisoning and handling. On the one hand, Roche and Boge (2000) believed that the increase in fish stressors after a single or fraction phenol injection is due to the disturbances in cells. The damage to endocrine cells in both liver and kidneys causes an abnormal biochemical blood value that one of which plays a major role in hemoglobin. Low Hb in tannin injection treatment showed that the spread of oxygen in the body of African catfish was disrupted. Barton (1987) described that a physiological stress will cause a new homeostatic condition in fish through a changing metabolism. This response to stress is controlled by the endocrine system through the release of cortisol hormones and catecholamine.

## CONCLUSION

The results from blood tests show that there is an increase in the cortisol hormone while the other parameters such as hemoglobin and hematocrit experience a decrease. This condition explains that the treatment of organic phenolic exposure is comparable to the inorganic phenol exposure. It can be seen from the changes in the blood hemoglobin and hematocrit as well as the cortisol hormones of African catfish.

## REFERENCES

1. Alamanda, I.E., N.S. Handajani, and A. Budiharjo. 2007. Penggunaan metode hematologi and pengamatan endoparasit darah untuk penetapan kesehatan Ikan Lele Dumbo (*Clarias gariepinus*) di kolam budidaya Desa Mangkubumen Boyolali. Biodiversitas, (1):34-38.
2. Anderson, D.P., and A.K. Siwicki. 1995. Basic hematology and serology for fish health programs: Fish Health Section, Asian Fisheries Society. 185-202pp.
3. Anderson, D.P., and A.K. Siwicki. 1994. Simplified assays for measuring nonspecific defense mechanisms in fish: Fish Health Section, Fisheries Soc. Meeting. 26 pp.
4. Barton, B.A., C.B. Schreck, and L.D. Barton. 1987. Effects of chronic cortisol administration and daily acute stress on growth, physiological conditions, and stress responses in juvenile rainbow trout. Diseases of Aquatic Organisms, (2):173-185.
5. Bastiawan, D., Taukhid, M. Alifuddin, and T.S. Dermawati. 1995. Perubahan hematologi and jaringan kan Lele Dumbo (*Clarias gariepinus*) Yang diinfeksi Cendawan *Aphariomyces* sp. Jurnal Penelitian Perikanan Indonesia: 106-115.
6. Bigoniya, P., and K. Singh. 2014. Ulcer protective potential of standardized hesperidin, a citrus flavonoid isolated from *Citrus sinensis*. Rev Bras Farma, (24):330-340.
7. Buikema, Jr.A.L., M.J. Mc. Ginniss, and Jr.J. Cairns. 1979. Phenolics in aquatic ecosystems: a selected review of recent literature. Mar Environ Res, (2):87-181.
8. Emrizal., A. Fernando, R. Yuliandri, K. Rullah, Nola, Indrayani, N.R, Reni, A. Susanty, R. Yerti, F. Ahmad, H.M. Sirat, D. Arbain. 2014. Cytotoxic activities of fractions and two isolated compounds from Sirih Merah (Indonesian red betel), *Piper crocatum* Ruiz and Pav. Procedia Chemistry, (13):79-84.

9. Foster, L.B., and R.T. Dunn. 1974. Single antibody technique for radioimmunoassay of cortisol in unextracted serum or plasma. *Clin Chem*, (20):365–368.
10. Harikrishnan, R., M.N. Rani, and C. Balasundaram. 2003. Haematological and biochemical parameters in common carp, *Cyprinus carpio*, following herbal treatment for *Aeromonas hydrophilla* infection. *Aquaculture*, 221:41-50.
11. Ibrahim. M.D. 2012. Experimental exposure of African catfish *Clarias gariepinus* (Burchell, 1822) to phenol: Clinical evaluation, tissue alterations and residue assessment. *Journal of Advanced Research* (3):177–183.
12. James, M.O., L.D. Stuchal, B.A. Nyagode. 2008. Glucuronidation and sulfonation in vitro of the major endocrine-active metabolites of methoxychlor in the channel catfish, *Ictalurus punctatus* and induction following treatment with 3-methylcholanthrene. *Aquatic Toxicology*, (2):227–238.
13. John, F., Carragler, and M.R. Christine. 1994. Primary and secondary stress responses in golden perch *Macquoria ambigua*. *J. comp. Biochem. Physiol*, 107A (1):40-56.
14. Mukherjee, D., S. Bhattacharya, V. Kumar, and J. Moitra. 1990. Biological significance of [<sup>14</sup>C] phenol accumulation in different organs of a murrel, *Channa punctatus*, and the common carp, *Cyprinus carpio*. *Biomed. Environ. Sci*, (3):337–342.
15. Nursyam, H., S. Andayani and A. Saputra. 2017. The effect of tannin from red betel (*Piper crocatum*) leaves towards blood biochemistry and histology of North African catfish (*Clarias gariepinus*). *AACL Bioflux.*, Volume 10, Issue 5, pp 1386-1393.
16. Roche, H., and G. Boge. 2000. Fish blood parameters as a potential tool for identification of stress caused by environmental factors or chemical intoxication. *Mar. Environ. Res. Part C* 125:27–44.
17. Saputra, A., S. Andayani and H. Nusyam. 2016. Total quantity of phenol and isolation methanol tannin extract of red betel leaf (*Piper crocatum*). *International Journal of PharmTech Research*. Vol.9, No.7, pp 146-153.
18. Sundang, M, S.N.S. Nasir, C.S. Sipaut, and H. Othman. 2012. Antioxidant activity, phenolic, flavonoid and tannin content of *Piper betle* and *Leucosyke capitella*. *Malaysian Journal of Fundamental and Applied Sciences*. (1):1-6.
19. Taysse, L., D. Troutaud, N.A. Khan, P. Deschaux. 1995. Structure activity relationship of phenolic compounds (phenol, pyrocatechol and hydroquinone) on natural lymphocytotoxicity of carp (*Cyprinus carpio*). *Toxicology*, (1-3):207–214.
20. Tilak, K.S., K. Veeriah, M.S. Butchiram. 2007. Effect of phenol on haematological components of Indian major carps *Catla catla*, *Labeo rohita* and *Cirrhinus mrigala*. *J Environ Biol*, (2):177–179.
21. Tsuzuki, M.Y., K. Ogawa, C.A. Strussmann. 2001. Physiological responses after stress and subsequent recovery at different salinities in adults pejerrey *Odontesthes bonariensis*. *Aquaculture*, 200:349-362.
22. Vasconcelos, P.C.P, M.A. Andreo, W. Vilegas, C.A. Hiruma-Lima, C.H. Pellizzon. 2010. Effect of Mouriri pusa tannins and flavonoids on prevention and treatment against experimental gastric ulcer. *Journal of Ethnopharmacology*, 131:146–153.
23. Varadarajan, R, H. Sankar H.S, J. Jose and B. Philip. 2014. Sublethal effects of phenolic compounds on biochemical, histological and ionoregulatory parameters in a tropical teleost fish *Oreochromis mossambicus* (Peters). *International Journal of Scientific and Research Publications*, (3):2250-3153.
24. Wahjuningrum, D., N. Ashry and S. Suryati. 2008. Pemanfaatan ekstrak daun Ketapang *Terminalia Cattapa* untuk pencegahan and pengobatan Ikan Patin (*Pangasionodon hypophthalmus*) yang Terinfeksi (*Aeromonas hydrophila*). *Jurnal Akuakultur Indonesia* 7(1):82-88.
25. Wedemeyer G.A., and W.T. Yasutake. 1977. Clinical methods for the assessment of the effects of environmental stress on fish health. Technical Paper of the U.S. Fish and Wildlife Service. Vol. 89. U.S. Department of the Interior Fish and Wildlife Service, Washington, D.C., USA. 18p.
26. Zaki, M.S., N. El-Batrawy, and N. M. Taha. 2012. Dose phenol toxicity affected endocrine status in African Catfish (*Clarias gariepinus*). *Life Science Journal*, 9(1):636-639.



# The Effect of Organic and Inorganic Phenolic Compounds on Hemoglobin, Hematocrit, and Cortisol Hormone of African Catfish (*Clarias Gariepinus*)

<sup>1</sup>Afandi Saputra, <sup>2</sup>Sri Andayani, <sup>2</sup>Happy Nusyam, <sup>1</sup>Djumbuh Rukmono, <sup>1</sup>Maria Goreti Eny, <sup>1</sup>Mugi Mulyono, <sup>1</sup>Sri Budiani, <sup>1</sup>Suharyadi, <sup>1</sup>Erni Marlina, <sup>1</sup>Moch. Nurhudah, <sup>1</sup>Effi A. Thaib, <sup>1</sup>Amyda S. Panjaitan, <sup>1</sup>Sinung Rahardjo, <sup>1</sup>Guntur Prabowo, <sup>1</sup>Ilham, <sup>1</sup>Fitriska Hapsyari

<sup>1</sup> Aquaculture Departement, Jakarta Fisheries University, Indonesia

<sup>2</sup> Faculty of Fisheries and Marine Sciences, Universitas Brawijaya, Indonesia

Corresponding Author Email: afandi.saputra@yahoo.co.id

**Abstract.** Phenolic are compounds found in the environment which can cause fish stressed or xenobiotic. The limited information about toxicology and pathological consequences in the fish exposed to phenolic compounds needs to be studied. In this research, the African catfish (*Clarias gariepinus*) were injected a single dose of phenol (inorganic) and *P.crocatum* Polyphenols (organic). The animals were exposed (75% of 96 hours LC50) with phenol concentration (3 mg/kg) and *P.crocatum* Polyphenols (1.8 mg/kg). Furthermore, the hematology was measured on 3 days after injection. The metabolic effects of hemoglobin and hematocrit decreased significantly ( $P < 0.05$ ) while the cortisol hormone increased ( $P < 0.05$ ) in a short-term after 3 days of exposure.

**Key Words:** phenolic, *Clarias gariepinus*, *P. crocatum* polyphenols, cortisol, hemoglobin, hematocrit.

**Introduction.** Phenolic compounds are xenobiotic compounds that affect anemia and disrupt the endocrine system and hormones (Writer *et al.*, 2010; Hori *et al.*, 2006; Awachie and Ezenwaji, 1998). This compound is classified as toxic chronic and immunotoxic which usually found in fish tissue (Taysse *et al.*, 1995; Mukherjee *et al.*, 1990). Phenol waste is accumulated from human activities such as the waste from industry, pharmaceuticals, and agriculture (Buikema *et al.*, 1979).

The phenol residue comes from industrial and pharmaceutical waste as well as and natural decomposition (Buikema *et al.*, 1979). Red Betel (*P. crocatum*) is one of the most aromatic plants in which the leaves contain phenolic compounds (flavonoids and tannin) and have a high phenol content (Saputra *et al.*, 2016). Phenolic compounds become a concern specifically because of its toxic effect in the water environment. However, this compound is difficult to detect due to the lack of taste and smell (Tilak *et al.*, 2007). A cytotoxic activity of red betel leaves on *Artemia* and African catfish was tested in-vivo based on the LC50 and showed that these plants have a potential of cytotoxic compounds. (Nursyam *et al.*, 2017 dan Emrizal *et al.*, 2014)

The habitat of *Clarias gariepinus* is in freshwaters which have a risk to be exposed to chemicals such as phenols (James *et al.*, 2008). Nursyam *et al.*, (2017) and Ibrahim (2012) reported that in-vivo test with an injection of phenolic compounds in fish *C. gariepinus* shows abnormal signs of glucose, BUN, and ALP in the blood and necrosis of liver and kidneys. Even though there is some research on the in-vivo effect of phenolic compounds towards fish blood parameters, there is no information about the residues and effects of organic phenolic compounds from plants to the toxicity on catfish. Therefore, further research is needed to determine the in-vivo influence of phenolic compounds of Red Betel Leaf (*Piper crocatum*) on Hemoglobin, Hematocrit, and Cortisol Hormones of African Catfish (*Clarias gariepinus*).

## Material and Method

**Tested Animals.** The African catfish (*Clarias gariepinus*) used were in the size of 12 to 15 grams. There was 20 fish/tub obtained from the Aquaculture Laboratory of Jakarta Fisheries University. The commercial feed was given twice a day at a dose of 3%. Before

the experiment was carried out, the fish was adapted in a concrete tub for 1 month. The quality of DO water conservancy was 3.99-6.53 mg/l with a pH of 7.0-7.6 at 28-30°C.

**Isolation and characterization of Organic and Inorganic phenolic compounds.** The organic phenolic compounds used were obtained from the isolation of red betel leaf (*P. crocatum*) using maceration, thin layer chromatography, column chromatography, and characterized by using UV-Vis and Infrared spectrophotometers (Saputra *et al.* 2016; Bigoniya and Singh, 2014; Sundang *et al.* 2012; Vasconcelos *et al.* 2010). The inorganic phenolic compounds were obtained from the commercial chemical store.

**Sub-lethal toxicity study.** The concentration of LC50 from inorganic phenol (3 mg/kg) and organic phenolic (1.80 mg/kg) is equal to 75% of LC50-96 hours. The exposure to phenolic compounds was done for 3 days after a single injection (Saputra *et al.* 2016).

**Blood sample.** The blood was collected 3 days after exposure to phenolic compounds. The blood was taken by using a 1 ml syringe on the cardinal vein and then centrifuged at 3000 rpm at 4°C for 15 minutes. The serum was separated and stored at a temperature of 15- 20°C (Saputra *et al.* 2016; Varadarajan *et al.*, 2014; Roche and Boge, 2000). The measurement of hemoglobin level was carried out by the Sahli method (Wedemeyer and Yasutake, 1977) in Hb/100ml while the hematocrit level was done by the capillary system (Anderson and Siwicki, 1995). On the other hand, the cortisol concentration plasma (Cort) was assessed by radioimmunoassay (RIA) (Foster and Dunn, 1974).

**Data analysis.** The data were analyzed by using the Analysis of Variance (ANOVA) with Minitab 16 for Windows.

## Results

**Hemoglobin.** The data analysis showed that the hemoglobin after being exposed with phenolic compounds have a significantly different situation in between the 5% level of treatment, Fhit: 35.58 ( $P > 0.05$ ). Based on the Figure 1, the hemoglobin rate of African catfish without treatment (control) is 13.67 Hb/100ml in which it is in line with the research from Alamanda *et al.*, (2007) that a generally healthy African catfish have a hemoglobin level ranged from 12-14 Hb/100 ml of blood.

Different conditions indicate that after the hemoglobin is injected, the phenolic compound was significantly decreased in the treatment of organic phenolic that is 7.67 Hb/100ml. In the inorganic phenolic treatment, it is shown that the hemoglobin was lower compared to the treatment of organic phenolic that is 6.67 Hb/100ml. According to Nusyam *et al.*, (2017) and Wahjuningrum *et al.*, (2008), low and high hemoglobin levels indicate that the fish might be sick or stressed. Roche and Boge (2000) stated that the increase or decrease in the plasma concentration of hemoglobin is a sign that there is an interference from the cells. Harikrishnan *et al.*, (2003) also asserted that the damage in red blood cells causes a decrease in hemoglobin content.




Figure 1. The Hemoglobin of African Catfish after Organic and Inorganic Phenolic Injection

**Hematocrit.** The results of data analysis showed that the hematocrit condition after being exposed to phenolic compounds was significantly different in between treatments in the level of 5%, Fhit: 17.35 ( $P > 0.05$ ). Based on the Figure 2 below, the hematocrit in African catfish without treatment (control) amounts to 37 %. This is in accordance with the opinion of Bastiawan *et al.*, (1995) that the normal hematocrit of African catfish is ranged from 30.8 to 45.5%.

The inorganic phenolic treatment showed that the hematocrit value was much lower than the organic treatment which was 23.33%. This value indicates that the treatment of inorganic phenolic greatly affects the decrease in the African catfish hematocrit. Tsuzuki *et al.*, (2001) stated that when the fish hematocrit is less than 22%, anemia will occur whereas if the hematocrit is greater than 60%, the fish will be in a state of stress. Anderson and Siwicki (1994) confirmed that low hematocrit can indicate the occurrence of fish contamination.




Figure 2. The Hematocrit of African Catfish after Organic and Inorganic Phenolic Injection

**Cortisol hormone.** This hormone is the result of endocrine gland secretion. The endocrine glands build the endocrine system which has the function of regulating the work of the body's physiological processes. The value of the cortisol hormone in catfish after being injected with phenolic compounds can be seen in Figure 3.




Figure 3. Cortisol Hormone of African Catfish after Organic and Inorganic Phenolic Injection

The results showed that the cortisol hormone after being exposed to phenolic compounds was significantly different in between the treatments (5%),  $F_{hit}: 3.86$  ( $P > 0.05$ ). Based on the results of the study, the cortisol in African catfish without treatment (control) is 412.76 nmol/L. The increase of cortisol hormone in phenolic treatment was compared with the control, in which the increase of the hormone was 198.86% in the organic phenolic treatment and 186.09% in the inorganic phenol treatment. Roche and Boge (2000) reported that the cortisol hormone in fish treatment was increased by 185.7-218%. According to Zaki *et al.*, (2012), there was an increase in *C.garipepinus* fish cortisol after being exposed to phenol for 3 months at 32.7%.

John *et al.*, (1994) explained that the increased cortisol in phenol treatment can be associated with fish poisoning and handling. On the one hand, Roche and Boge (2000) believed that the increase in fish stressors after a single or fraction phenol injection is due to the disturbances in cells. The damage to endocrine cells in both liver and kidneys causes an abnormal biochemical blood value that one of which plays a major role in hemoglobin. Low Hb in tannin injection treatment showed that the spread of oxygen in the body of African catfish was disrupted. Barton (1987) described that a physiological stress will cause a new homeostatic condition in fish through a changing metabolism. This response to stress is controlled by the endocrine system through the release of cortisol hormones and catecholamine.

**Conclusions.** The results from blood tests show that there is an increase in the cortisol hormone while the other parameters such as hemoglobin and hematocrit experience a decrease. This condition explains that the treatment of organic phenolic exposure is comparable to the inorganic phenol exposure. It can be seen from the changes in the blood hemoglobin and hematocrit as well as the cortisol hormones of African catfish.

## References

Alamanda IE, Handajani NS & Budiharjo, A, Penggunaan metode hematologi dan pengamatan endoparasit darah untuk penetapan kesehatan Ikan Lele Dumbo

- (*Clarias gariepinus*) di kolam budidaya Desa Mangkubumen Boyolali. *Biodiversitas*, 1 (2017) 34.
- Anderson DP & Siwicki AK, Basic hematology and serology for fish health programs: Fish Health Section. *Asian Fisheries Society*, 21 (1995) 185.
- Anderson DP & Siwicki AK, Simplified assays for measuring nonspecific defense mechanisms in fish: Fish Health Section. *Fisheries Soc. Meeting*, 2 (1994) 26.
- Barton BA, Schreck CB & Barton LD, Effects of chronic cortisol administration and daily acute stress on growth, physiological conditions, and stress responses in juvenile rainbow trout. *Diseases of Aquatic Organisms*, 2 (1987) 173.
- Bastiawan D, Tauhid M, Alifuddin & Dermawati TS, Perubahan hematologi dan jaringan kan Lele Dumbo (*Clarias gariepinus*) Yang diinfeksi Cendawan *Apharioomyces* sp. *Jurnal Penelitian Perikanan Indonesia*, 22 (1995) 106.
- Bigoniya P & Singh K, Ulcer protective potential of standardized hesperidin, a citrus flavonoid isolated from *Citrus sinensis*. *Rev Bras Farma*, 24 (2014) 330.
- Buikema AL, Ginniss MJ & Cairns J, Phenolics in aquatic ecosystems: a selected review of recent literature. *Mar Environ Res*, 2 (1979) 87.
- Emrizal A, Fernando R, Yuliandri K, Rullah, Nola IRN, Reni A, Susanty R, Yerti F, Ahmad HM, Sirat D, & Arbain, Cytotoxic activities of fractions and two isolated compounds from Sirih Merah (Indonesian red betel), *Piper crocatum* Ruiz and Pav. *Procedia Chemistry*, 13 (2014) 79.
- Foster LB & Dunn RT, Single antibody technique for radioimmunoassay of cortisol in unextracted serum or plasma. *Clin Chem*, 20 (1974).
- Harikrishnan R, Rani MN & Balasundaram C, Haematological and biochemical parameters in common carp, *Cyprinus carpio*, following herbal treatment for *Aeromonas hydrophilla* infection. *Aquaculture*, 221 (2003) 41.
- Ibrahim MD, Experimental exposure of African catfish *Clarias gariepinus* (Burchell, 1822) to phenol: Clinical evaluation, tissue alterations and residue assessment. *Journal of Advanced Research*, 3 (2012) 177.
- James MO, Stuchal LD & Nyagode BA, Glucuronidation and sulfonation in vitro of the major endocrine-active metabolites of methoxychlor in the channel catfish, *Ictalurus punctatus* and induction following treatment with 3-methylcholanthrene. [Aquatic Toxicology](#), 2 (2008) 227.
- John F, Carragler & Christine MR, Primary and secondary stress responses in golden perch *Macquoria ambigua*. *J. comp. Biochem. Physiol*, 107 (194) 40.
- Mukherjee D, Bhattacharya S, Kumar V & Moitra J, Biological significance of [<sup>14</sup>C] phenol accumulation in different organs of a murrel, *Channa punctatus*, and the common carp, *Cyprinus carpio*. *Biomed. Environ. Sci*, 3 (1990) 337.
- Nursyam H, Andayani S & Saputra A, The effect of tannin from red betel (*Piper crocatum*) leaves towards blood biochemistry and histology of North African catfish (*Clarias gariepinus*). *AAFL Bioflux*, 10 (2017) 1386.
- Roche H & Boge G, Fish blood parameters as a potential tool for identification of stress caused by environmental factors or chemical intoxication. *Mar. Environ. Res*, 125 (2000) 27.
- Saputra A, Andayani S & Nusyam H, Total quantity of phenol and isolation methanol tannin extract of red betel leaf (*Piper crocatum*). *International Journal of PharmTech Research*, 9 (2016) 146.
- Sundang M, Nasir SNS, Sipaut CS & Othman H. (2012), Antioxidant activity, phenolic, flavonoid and tannin content of *Piper betle* and *Leucosyke capitella*. *Malaysian Journal of Fundamental and Applied Sciences*, 1 (2012) 1.
- Taysse L, Troutaud D, Khan NA & Deschaux P, Structure activity relationship of phenolic compounds (phenol, pyrocatechol and hydroquinone) on natural lymphocytotoxicity of carp (*Cyprinus carpio*). *Toxicology*, 1 (1995) 207.
- Tilak KS, Veeraiah K, Butchiram MS, Effect of phenol on haematological components of Indian major carps *Catla catla*, *Labeo rohita* and *Cirrhinus mrigala*. *J Environ Biol*, 2 (2007) 177.

- Tsuzuki MY, Ogawa K & Strussmann CA, Physiological responses after stress and subsequent recovery at different salinities in adults pejerrey *Odontesthes bonariensis*. *Aquaculture*, 200 (2001) 349.
- Vasconcelos PCP, Andreo MA, Vilegas W, Hiruma-Lima CA & Pellizzon CH, Effect of *Mouriri pusa* tannins and flavonoids on prevention and treatment against experimental gastric ulcer. *Journal of Ethnopharmacology*, 131 (2010) 146.
- Varadarajan R, Sankar H, Jose J & Philip B, Sublethal effects of phenolic compounds on biochemical, histological and ionoregulatory parameters in a tropical teleost fish *Oreochromis mossambicus* (Peters). *International Journal of Scientific and Research Publications*, 3 (2014) 2250.
- Wahjuningrum D, Ashry N & Suryati S, Pemanfaatan ekstrak daun Ketapang *Terminalia Cattapa* untuk pencegahan dan pengobatan Ikan Patin (*Pangasionodon hypophthalmus*) yang Terinfeksi (*Aeromonas hydrophila*). *Jurnal Akuakultur Indonesia*, 7 (2008) 82.
- Wedemeyer GA & Yasutake WT, *Clinical methods for the assessment of the effects of environmental stress on fish health* (Department of the Interior Fish and Wildlife Service, USA), 1997, 80.
- Zaki MS, El-Batrawy N & Taha NM, Dose phenol toxicity affected endocrine status in African Catfish (*Clarias gariepinus*). *Life Science Journal*, 9 (2012) 636.

Received: 23 August 2016. Accepted: 25 September 2016. Published online: 18 October 2016.

Authors:

Afandi Saputra, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail: [afandi.saputra@yahoo.co.id](mailto:afandi.saputra@yahoo.co.id).

Sri Andayani, Faculty of Fisheries and Marine Sciences, Universitas Brawijaya, Jl. Veteran, Malang, Indonesia, e-mail:

Happy Nusyam, Faculty of Fisheries and Marine Sciences, Universitas Brawijaya, Jl. Veteran, Malang, Indonesia, e-mail:

Djumbuh Rukmono, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

Maria Goreti Eny, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

Mugi Mulyono, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

Sri Budiani, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

Suharyadi, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

Erni Marlina, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

Moch. Nurhudah, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

Effi A. Thaib, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

Amyda S. Panjaitan, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

Sinung Rahardjo, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

Guntur Prabowo, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

Ilham, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

Fitriska Hapsyari, Aquaculture Department, Jakarta Fisheries University, Jl. AUP Pasar Minggu, Jakarta, Indonesia, e-mail:

# RUSSIAN JOURNAL OF AGRICULTURAL AND SOCIO- ECONOMIC SCIENCES



ISSN 2226-1184

Dear Dr Saputra, 5 November 2019

Greetings and Seasons Wishes.

We would like to inform you that,

your paper titled, “ **EFFECT OF ORGANIC AND INORGANIC PHENOLIC COMPOUNDS ON HEMOGLOBIN, HEMATOCRIT, AND CORTISOL HORMONE OF AFRICAN CATFISH (*Clarias gariepinus*)**”

is selected for publication in: Vol. 96., Issue 12, Upcoming Regular issue, 2019 of Russian Journal Of Agricultural And Socio-Economic Sciences (RJOAS)

Every author will have to pay a nominal cost of **US \$ 225** for his full paper. Journals do not have subscription base/any sponsorship and are available for free search .Thus for existence/financial independencies of the journal, authors have to pay as nominal charges. (Cost towards processing, maintenance of paper in secured data storage system, databases; to meet various overhead expenses like paper setting costs, employees charges, WEB cost, electr. Cost, Office cost, communication cost, maintenance cost, time cost etc.).

Thus your paper is charged with, For Authors 3.174.750 IDR.

**\*\*Pay the charges ONLY AND ONLY to the following Representative in his Bank account at INDONESIA. ---**

Thanks,

Expecting a reply at your earliest for further processing of your paper.

Sincerely

***Dr.Sergey Plygun***

*Editor, Journal Of Agricultural And Socio-Economic Sciences (RJOAS)*

**THE EFFECT OF ORGANIC AND INORGANIC PHENOLIC COMPOUNDS  
ON HEMOGLOBIN, HEMATOCRIT, AND CORTISOL HORMONE  
OF AFRICAN CATFISH (*CLARIAS GARIEPINUS*)**

**Saputra Afandi<sup>1\*</sup>, Andayani Sri<sup>2</sup>, Nusyam Happy<sup>2</sup>, Rukmono Djumbuh<sup>1</sup>,  
Eny Maria Goreti<sup>1</sup>, Mulyono Mugi<sup>1</sup>, Budiani Sri<sup>1</sup>, Suharyadi<sup>1</sup>, Marlina Erni<sup>1</sup>,  
Nurhudah M.<sup>1</sup>, Thaib Effi A.<sup>1</sup>, Panjaitan Amyda S.<sup>1</sup>, Rahardjo Sinung<sup>1</sup>,  
Prabowo Guntur<sup>1</sup>, Ilham<sup>1</sup>, Hapsyari Fitriska<sup>1</sup>, Laishevtcev A.<sup>3,4</sup>**

<sup>1</sup>Department of Aquaculture, Jakarta Fisheries University, Indonesia

<sup>2</sup>Faculty of Fisheries and Marine Science, University of Brawijaya, Malang, Indonesia

<sup>3</sup>Federal Scientific Centre VIEV, Moscow, Russia

<sup>4</sup>Laboratory of Biological Control and Antimicrobial Resistance,  
Orel State University named after I.S. Turgenev, Orel City, Russia

\*E-mail: [afandi.saputra@yahoo.co.id](mailto:afandi.saputra@yahoo.co.id)

**ABSTRACT**

Phenolic are compounds found in the environment which can cause fish stressed or xenobiotic. The limited information about toxicology and pathological consequences in the fish exposed to phenolic compounds needs to be studied. In this research, the African catfish (*Clarias gariepinus*) were injected a single dose of phenol (inorganic) and *P.crocatum* Polyphenols (organic). The animals were exposed (75% of 96 hours LC50) with phenol concentration (3 mg/kg) and *P.crocatum* Polyphenols (1.8 mg/kg). Furthermore, the hematology was measured on 3 days after injection. The metabolic effects of hemoglobin and hematocrit decreased significantly ( $P < 0.05$ ) while the cortisol hormone increased ( $P < 0.05$ ) in a short-term after 3 days of exposure.

**KEY WORDS**

Phenolic, *Clarias gariepinus*, *P. crocatum* polyphenols, cortisol, hemoglobin, hematocrit.

Phenolic compounds are xenobiotic compounds that affect anemia and disrupt the endocrine system and hormones (Writer *et al.*, 2010; Hori *et al.*, 2006; Awachie and Ezenwaji, 1998). This compound is classified as toxic chronic and immunotoxic which usually found in fish tissue (Taysse *et al.*, 1995; Mukherjee *et al.*, 1990). Phenol waste is accumulated from human activities such as the waste from industry, pharmaceuticals, and agriculture (Buikema *et al.*, 1979).

The phenol residue comes from industrial and pharmaceutical waste as well as and natural decomposition (Buikema *et al.*, 1979). Red Betel (*P. crocatum*) is one of the most aromatic plants in which the leaves contain phenolic compounds (flavonoids and tannin) and have a high phenol content (Saputra *et al.*, 2016). Phenolic compounds become a concern specifically because of its toxic effect in the water environment. However, this compound is difficult to detect due to the lack of taste and smell (Tilak *et al.*, 2007). A cytotoxic activity of red betel leaves on *Artemia* and African catfish was tested in-vivo based on the LC50 and showed that these plants have a potential of cytotoxic compounds. (Nursyam *et al.*, 2017 and Emrizal *et al.*, 2014)

The habitat of *Clarias gariepinus* is in freshwaters which have a risk to be exposed to chemicals such as phenols (James *et al.*, 2008). Nursyam *et al.*, (2017) and Ibrahim (2012) reported that in-vivo test with an injection of phenolic compounds in fish *C. gariepinus* shows abnormal signs of glucose, BUN, and ALP in the blood and necrosis of liver and kidneys. Even though there is some research on the in-vivo effect of phenolic compounds towards fish blood parameters, there is no information about the residues and effects of organic phenolic compounds from plants to the toxicity on catfish. Therefore, further research is needed to



determine the in-vivo influence of phenolic compounds of Red Betel Leaf (*Piper crocatum*) on Hemoglobin, Hematocrit, and Cortisol Hormones of African Catfish (*Clarias gariepinus*).

## MATERIALS AND METHODS OF RESEARCH

**Tested Animal.** The African catfish (*Clarias gariepinus*) used were in the size of 12 to 15 grams. There was 20 fish/tub obtained from the Aquaculture Laboratory of Jakarta Fisheries University. The commercial feed was given twice a day at a dose of 3%. Before the experiment was carried out, the fish was adapted in a concrete tub for 1 month. The quality of DO water conservancy was 3.99-6.53 mg/l with a pH of 7.0-7.6 at 28-30°C.

**Isolation and characterization of Organic and Inorganic phenolic compounds.** The organic phenolic compounds used were obtained from the isolation of red betel leaf (*P. crocatum*) using maceration, thin layer chromatography, column chromatography, and characterized by using UV-Vis and Infrared spectrophotometers (Saputra *et al.* 2016; Bigoniya and Singh, 2014; Sundang *et al.* 2012; Vasconcelos *et al.* 2010). The inorganic phenolic compounds were obtained from the commercial chemical store.

**Sub-lethal toxicity study.** The concentration of LC50 from inorganic phenol (3 mg/kg) and organic phenolic (1.80 mg/kg) is equal to 75% of LC50-96 hours. The exposure to phenolic compounds was done for 3 days after a single injection (Saputra *et al.* 2016)

**Blood sample.** The blood was collected 3 days after exposure to phenolic compounds. The blood was taken by using a 1 ml syringe on the cardinal vein and then centrifuged at 3000 rpm at 4°C for 15 minutes. The serum was separated and stored at a temperature of 15- 20°C (Saputra *et al.* 2016; Varadarajan *et al.*, 2014; Roche and Boge, 2000). The measurement of hemoglobin level was carried out by the Sahli method (Wedemeyer and Yasutake, 1977) in Hb/100ml while the hematocrit level was done by the capillary system (Anderson and Siwicki, 1995). On the other hand, the cortisol concentration plasma (Cort) was assessed by radioimmunoassay (RIA) (Foster and Dunn, 1974).

**Data analysis.** The data were analyzed by using the Analysis of Variance (ANOVA) with Minitab 16 for windows.

## RESULTS AND DISCUSSION

The data analysis showed that the hemoglobin after being exposed with phenolic compounds have a significantly different situation in between the 5% level of treatment, Fhit: 35.58 (P> 0.05). Based on the Figure 1, the hemoglobin rate of African catfish without treatment (control) is 13.67 Hb/100ml in which it is in line with the research from Alamanda *et al.*, (2007) that a generally healthy African catfish have a hemoglobin level ranged from 12-14 Hb/100 ml of blood.

Different conditions indicate that after the hemoglobin is injected, the phenolic compound was significantly decreased in the treatment of organic phenolic that is 7.67 Hb/100ml. In the inorganic phenolic treatment, it is shown that the hemoglobin was lower compared to the treatment of organic phenolic that is 6.67 Hb/100ml. According to Nusyam *et al.*, (2017) and Wahjuningrum *et al.*, (2008), low and high hemoglobin levels indicate that the fish might be sick or stressed. Roche and Boge (2000) stated that the increase or decrease in the plasma concentration of hemoglobin is a sign that there is an interference from the cells. Harikrishnan *et al.*, (2003) also asserted that the damage in red blood cells causes a decrease in hemoglobin content.

The results of data analysis showed that the hematocrit condition after being exposed to phenolic compounds was significantly different in between treatments in the level of 5%, Fhit: 17.35 (P>0.05). Based on the Figure 2 below, the hematocrit in African catfish without treatment (control) amounts to 37 %. This is in accordance with the opinion of Bastiawan *et al.*, (1995) that the normal hematocrit of African catfish is ranged from 30.8 to 45.5%.

The inorganic phenolic treatment showed that the hematocrit value was much lower than the organic treatment which was 23.33%. This value indicates that the treatment of inorganic phenolic greatly affects the decrease in the African catfish hematocrit. Tsuzuki *et*

*al.*, (2001) stated that when the fish hematocrit is less than 22%, anemia will occur whereas if the hematocrit is greater than 60%, the fish will be in a state of stress. Anderson and Siwicki (1994) confirmed that low hematocrit can indicate the occurrence of fish contamination.




Figure 1 – The Hemoglobin of African Catfish after Organic and Inorganic Phenolic Injection




Figure 2 – The Hematocrit of African Catfish after Organic and Inorganic Phenolic Injection




Figure 3 – Cortisol Hormone of African Catfish after Organic and Inorganic Phenolic Injection

This hormone is the result of endocrine gland secretion. The endocrine glands build the endocrine system which has the function of regulating the work of the body's physiological

processes. The value of the cortisol hormone in catfish after being injected with phenolic compounds can be seen in Figure 3.

The results showed that the cortisol hormone after being exposed to phenolic compounds was significantly different in between the treatments (5%), Fhit: 3.86 ( $P > 0.05$ ). Based on the results of the study, the cortisol in African catfish without treatment (control) is 412.76 nmol/L. The increase of cortisol hormone in phenolic treatment was compared with the control, in which the increase of the hormone was 198.86% in the organic phenolic treatment and 186.09% in the inorganic phenol treatment. Roche and Boge (2000) reported that the cortisol hormone in fish treatment was increased by 185.7-218%. According to Zaki *et al.*, (2012), there was an increase in *C.gariepinus* fish cortisol after being exposed to phenol for 3 months at 32.7%.

John *et al.*, (1994) explained that the increased cortisol in phenol treatment can be associated with fish poisoning and handling. On the one hand, Roche and Boge (2000) believed that the increase in fish stressors after a single or fraction phenol injection is due to the disturbances in cells. The damage to endocrine cells in both liver and kidneys causes an abnormal biochemical blood value that one of which plays a major role in hemoglobin. Low Hb in tannin injection treatment showed that the spread of oxygen in the body of African catfish was disrupted. Barton (1987) described that a physiological stress will cause a new homeostatic condition in fish through a changing metabolism. This response to stress is controlled by the endocrine system through the release of cortisol hormones and catecholamine.

## CONCLUSION

The results from blood tests show that there is an increase in the cortisol hormone while the other parameters such as hemoglobin and hematocrit experience a decrease. This condition explains that the treatment of organic phenolic exposure is comparable to the inorganic phenol exposure. It can be seen from the changes in the blood hemoglobin and hematocrit as well as the cortisol hormones of African catfish.

## REFERENCES

1. Alamanda, I.E., N.S. Handajani, and A. Budiharjo. 2007. Penggunaan metode hematologi and pengamatan endoparasit darah untuk penetapan kesehatan Ikan Lele Dumbo (*Clarias gariepinus*) di kolam budidaya Desa Mangkubumen Boyolali. Biodiversitas, (1):34-38.
2. Anderson, D.P., and A.K. Siwicki. 1995. Basic hematology and serology for fish health programs: Fish Health Section, Asian Fisheries Society. 185-202pp.
3. Anderson, D.P., and A.K. Siwicki. 1994. Simplified assays for measuring nonspecific defense mechanisms in fish: Fish Health Section, Fisheries Soc. Meeting. 26 pp.
4. Barton, B.A., C.B. Schreck, and L.D. Barton. 1987. Effects of chronic cortisol administration and daily acute stress on growth, physiological conditions, and stress responses in juvenile rainbow trout. Diseases of Aquatic Organisms, (2):173-185.
5. Bastiawan, D., Tauhid, M. Alifuddin, and T.S. Dermawati. 1995. Perubahan hematologi and jaringan kan Lele Dumbo (*Clarias gariepinus*) Yang diinfeksi Cendawan *Aphariomyces* sp. Jurnal Penelitian Perikanan Indonesia: 106-115.
6. Bigoniya, P., and K. Singh. 2014. Ulcer protective potential of standardized hesperidin, a citrus flavonoid isolated from *Citrus sinensis*. Rev Bras Farma, (24):330-340.
7. Buikema, Jr.A.L., M.J. Mc. Ginniss, and Jr.J. Cairns. 1979. Phenolics in aquatic ecosystems: a selected review of recent literature. Mar Environ Res, (2):87-181.
8. Emrizal., A. Fernando, R. Yuliandri, K. Rullah, Nola, Indrayani, N.R, Reni, A. Susanty, R. Yerti, F. Ahmad, H.M. Sirat, D. Arbain. 2014. Cytotoxic activities of fractions and two isolated compounds from Sirih Merah (Indonesian red betel), *Piper crocatum* Ruiz and Pav. Procedia Chemistry, (13):79-84.

9. Foster, L.B., and R.T. Dunn. 1974. Single antibody technique for radioimmunoassay of cortisol in unextracted serum or plasma. *Clin Chem*, (20):365–368.
10. Harikrishnan, R., M.N. Rani, and C. Balasundaram. 2003. Haematological and biochemical parameters in common carp, *Cyprinus carpio*, following herbal treatment for *Aeromonas hydrophilla* infection. *Aquaculture*, 221:41-50.
11. Ibrahim. M.D. 2012. Experimental exposure of African catfish *Clarias gariepinus* (Burchell, 1822) to phenol: Clinical evaluation, tissue alterations and residue assessment. *Journal of Advanced Research* (3):177–183.
12. James, M.O., L.D. Stuchal, B.A. Nyagode. 2008. Glucuronidation and sulfonation in vitro of the major endocrine-active metabolites of methoxychlor in the channel catfish, *Ictalurus punctatus* and induction following treatment with 3-methylcholanthrene. *Aquatic Toxicology*, (2):227–238.
13. John, F., Carragler, and M.R. Christine. 1994. Primary and secondary stress responses in golden perch *Macquoria ambigua*. *J. comp. Biochem. Physiol*, 107A (1):40-56.
14. Mukherjee, D., S. Bhattacharya, V. Kumar, and J. Moitra. 1990. Biological significance of [<sup>14</sup>C] phenol accumulation in different organs of a murrel, *Channa punctatus*, and the common carp, *Cyprinus carpio*. *Biomed. Environ. Sci*, (3):337–342.
15. Nursyam, H., S. Andayani and A. Saputra. 2017. The effect of tannin from red betel (*Piper crocatum*) leaves towards blood biochemistry and histology of North African catfish (*Clarias gariepinus*). *AAFL Bioflux.*, Volume 10, Issue 5, pp 1386-1393.
16. Roche, H., and G. Boge. 2000. Fish blood parameters as a potential tool for identification of stress caused by environmental factors or chemical intoxication. *Mar. Environ. Res. Part C* 125:27–44.
17. Saputra, A., S. Andayani and H. Nusyam. 2016. Total quantity of phenol and isolation methanol tannin extract of red betel leaf (*Piper crocatum*). *International Journal of PharmTech Research*. Vol.9, No.7, pp 146-153.
18. Sundang, M, S.N.S. Nasir, C.S. Sipaut, and H. Othman. 2012. Antioxidant activity, phenolic, flavonoid and tannin content of *Piper betle* and *Leucosyke capitella*. *Malaysian Journal of Fundamental and Applied Sciences*. (1):1-6.
19. Taysse, L., D. Troutaud, N.A. Khan, P. Deschaux. 1995. Structure activity relationship of phenolic compounds (phenol, pyrocatechol and hydroquinone) on natural lymphocytotoxicity of carp (*Cyprinus carpio*). *Toxicology*, (1-3):207–214.
20. Tilak, K.S., K. Veeraiyah, M.S. Butchiram. 2007. Effect of phenol on haematological components of Indian major carps *Catla catla*, *Labeo rohita* and *Cirrhinus mrigala*. *J Environ Biol*, (2):177–179.
21. Tsuzuki, M.Y., K. Ogawa, C.A. Strussmann. 2001. Physiological responses after stress and subsequent recovery at different salinities in adults pejerrey *Odontesthes bonariensis*. *Aquaculture*, 200:349-362.
22. Vasconcelos, P.C.P, M.A. Andreo, W. Vilegas, C.A. Hiruma-Lima, C.H. Pellizzon. 2010. Effect of Mouriri pusa tannins and flavonoids on prevention and treatment against experimental gastric ulcer. *Journal of Ethnopharmacology*, 131:146–153.
23. Varadarajan, R, H. Sankar H.S, J. Jose and B. Philip. 2014. Sublethal effects of phenolic compounds on biochemical, histological and ionoregulatory parameters in a tropical teleost fish *Oreochromis mossambicus* (Peters). *International Journal of Scientific and Research Publications*, (3):2250-3153.
24. Wahjuningrum, D., N. Ashry and S. Suryati. 2008. Pemanfaatan ekstrak daun Ketapang *Terminalia Cattapa* untuk pencegahan and pengobatan Ikan Patin (*Pangasionodon hypophthalmus*) yang Terinfeksi (*Aeromonas hydrophila*). *Jurnal Akuakultur Indonesia* 7(1):82-88.
25. Wedemeyer G.A., and W.T. Yasutake. 1977. Clinical methods for the assessment of the effects of environmental stress on fish health. Technical Paper of the U.S. Fish and Wildlife Service. Vol. 89. U.S. Department of the Interior Fish and Wildlife Service, Washington, D.C., USA. 18p.
26. Zaki, M.S., N. El-Batrawy, and N. M. Taha. 2012. Dose phenol toxicity affected endocrine status in African Catfish (*Clarias gariepinus*). *Life Science Journal*, 9(1):636-639.