



## INCREASED COLOR BRIGHTNESS WITH THE ADDITION OF ASTAXANTHIN IN KOI FISH (*CYPRINUS RUBROFUSCUS*) FEED

Received: 00-00-0000

Accepted: 00-00-0000

Published: 00-00-0000

### ABSTRACT

**Introductions:** This research was conducted experimentally from May to July 2021 at the Teaching Factory (TEFA) Cultivation Laboratory of the Jakarta Fisheries Business Expert Polytechnic. This study aimed to determine the effect of adding astaxanthin to feed on increasing the colour, survival and growth of koi fish. **Method:** The research method used is quantitative. The design was completely randomized (CRD) with four treatment levels and three replications. The analysis of variance used was an analysis of variance (ANOVA). **Result:** The results showed that astaxanthin added to feed significantly affected changes in the colour quality of koi fish. The best result was the addition of astaxanthin 125 mg/kg feed ( $20.29 \pm 0.10$ ). Results The average weight measurement (ABW) at the end of the study obtained the highest value in treatment A (control) of 10.37 g. The results of the average weight measurement (ABW) at the end of the study obtained the highest value in the treatment A (control) of 10.37 g, the highest specific growth rate (SGR) in treatment A (control) of 12.00%, the best Feed Conversion Ratio (FCR) in the treatment with the addition of astaxanthin 100 mg 1.12. **Conclusion:** Astaxanthin is a safe and effective way to enhance the color of koi fish without any negative impact on their growth or survival. By adding astaxanthin to koi fish feed in the correct dosage, owners can achieve optimal results in color enhancement, improving the quality and value of the fish in the market.

**Keywords:** astaxanthin, *Cyprinus rubrofuscus*, fish colour.

**Comment [A1]:** should include the implications of the research

Corresponding Author:

E-mail:



### INTRODUCTION

Koi fish (*Cyprinus rubrofuscus*) is well-known and popular with ornamental fish lovers and business people (Iskandar et al., 2021). The selling price of koi fish continues to increase, and the price classification is very varied. The main factors determining the price of koi fish are beautiful colour patterns and exotic body shapes, beautiful body shapes and movements, as well as very lucrative economic value, causing koi fish to continue to be studied both from the aspect of feed nutrition and a genetic standpoint (Simbolon, et al., 2002).

beautiful body shape and colour of koi fish (*Cyprinus rubrofuscus*) make this fish have high economic value (Irawati et al., 2022). Indicators of beauty in ornamental fish can be seen in their brilliant colours, physical shape and completeness, behaviour, and health or stamina conditions (Andriani et al., 2019). According to (Malini & Agustin, 2018), the beauty of koi fish lies in its back, which has an attractive colour. The colour of koi fish is related to the number and location of pigment cells (chromatophores) in the dermis layer. It is said by (Simbolon et al., 2002) that the body colour of the fish is caused by the availability of carotene from natural food, while the fish that are reared get their carotene source from artificial feed.

The primary function of pigments in fish food will naturally form colour changes in fish so that fish scales are more contrasting. The addition of pigmentation ingredients in the feed will encourage an increase in colour pigments in the fish's body and be able to maintain the dyes in the

DOIs:

1

body (Nur et al., 2020). The increase in colour intensity in fish is influenced by two factors, namely internal factors and external factors (Collins et al., 2022). Internal factors are factors originating from within the body of the fish that are fixed, such as age, size, genetics, sex, and the fish's ability to absorb the nutrients in the feed (Indarti et al., 2012)

The pigment manipulation technique by enriching the content of pigment cells in the fish's body through feeding containing astaxanthin (chlorophyll pink) is a way to obtain bright and even colours in fish (Sitorus, 2015). Provision of feed-containing supplements needs to be done in order to improve and improve colour quality (Maolana et al., 2018).

Carotenoids are dye-forming components that give red and yellow colours. Astaxanthin is the most effective and widely found type of carotenoid pigment for colouring (Yulianti et al., 2014). Nasution (1997) in (Andriani et al., 2019) stated that the greater the carotenoid content in the fish's body, the brighter the colour of the fish. According to (Sukarman & Hirnawati, 2014), astaxanthin is a synthetic carotenoid commonly used in aquaculture. Thus the need for carotenoids must be provided through feed, especially in intensive rearing systems (Sukarman & Hirnawati, 2014).

Based on the background above, this study aimed to determine and evaluate the effect of adding astaxanthin to feed on increasing the colour of koi fish (*Cyprinus rubrofusca*).

**Comment [A2]:** restate the purpose of the research in a special paragraph

## METHODS

This research activity was conducted from May to July 2021 at the Cultivation Laboratory of the Jakarta Fisheries Business Expert Polytechnic. The materials used in this study were 120 koi fish of the Kohaku type with a size of  $6 \pm 7$  cm. Other ingredients used are artificial feed, prologue and astaxanthin. The design used was a completely randomized design (CRD) with four treatments and three replications with doses A (Control/without astaxanthin), B (75 mg/kg feed), C (100 mg/kg feed), D (125 mg/kg feed). Kg of feed). The feed is poured into the astaxanthin solution, mixed with a prologue of as much as 2gr/kg of feed and 150 ml/kg of water, and then dried.

The research method used is quantitative. The analysis of variance used was an analysis of variance (ANOVA) with a 5% degree of freedom. If the statistical test results show a significant difference between the treatments ( $P < 0.05$ ), proceed with the Tukey HSD test. Then the data were processed using SPSS software version 26 to find out whether there was an effect on each treatment.


The increase in colour in koi fish was observed using the Modified-Toca Color Finder (M-TFC), carried out six times during 60 days of rearing. The fish colour was observed visually by three panellists with eyes that were not myopic and not colour-blind (Indarti et al., 2012). (Sitorus, 2015) said that determining the colour score using M-TCF paper on each fish. The score on the paper starts with the numbers 1,2,3, with colour gradations starting from light orange to dark red, where the higher the value, the darker the colour.

## RESULTS AND DISCUSSION

The results showed that adding astaxanthin to the feed had a significant effect ( $P < 0.05$ ) on the brightness of the koi fish's colour. Based on the Tukey HSD test output, sig  $< 0.05$  was obtained

between the treatments of adding astaxanthin 125mg/kg of feed to other treatments. The increase in the colour of koi fish at the end of maintenance can be seen in Figure 1

Comment [A3]: don't disconnect




**Table 1**

**Average Color Value of Koi Fish**

Astaxanthin dosage	Absolute Color Change
Control	18.58 ± 0.25 <sup>a</sup>
75 mg/kg feed	18.71 ± 0.21 <sup>b</sup>
100 mg/kg feed	19.64 ± 0.67 <sup>b</sup>
125 mg/kg feed	20.29 ± 0.10 <sup>c</sup>

The results of the average weight measurement (*Average Body Weight / ABW*) at the end of the study obtained the highest value in treatment A (control) of 10.37 g, the highest specific growth rate (*Specific Growth Rate / SGR*) in treatment A (control) of 12.00 %, the best Feed Conversion Ratio (*FCR*) in the treatment with the addition of 100 mg astaxanthin 1.12.

Survival or survival is the ratio between the number of individuals who live at the end of the experiment with the number of individuals who live at the end of the experiment at e start of the experiment. Koi fish survival at the end of rearing can be seen in Figure 2.



The best survival was obtained in treatment B with 75 mg of astaxanthin w, a value of 93.3% and treatment C of 90.0%.

Good water quality plays a vital role in the life of fish, where fish need a suitable and qualified environment to grow properly. The results of measuring water quality parameters during maintenance can be seen in Table 2.

**Table 2**  
**Koi Fish Rearing Water Quality**

Parameter	Unit	Average
Temperature	°C	26.5-28.7
pH	-	7.6-7.9
Dissolved Oxygen/DO	mg l <sup>-1</sup>	4.8-5.5

The results of water quality parameter measurements in Table 2 show that the temperature ranges from 26.8-29.5°C, the pH ranges from 7.6-7.9 a, and the DO ranges from 4.8-5.5 mg l<sup>-1</sup>. This water quality range can be said to be optimal for fish life. According to (Purba et al., 2021), a suitable temperature range for koi fish is 27-29°C. According to (Kelabora, 2010) that temperature can affect the growth and appetite of fish. The ideal pH for freshwater biota is 7-8.5 (Muslim & Atjo, 2021). The minimum dissolved oxygen concentration for keeping koi fish is mg l<sup>-1</sup> (Simamora et al., 2021).

#### CONCLUSION

The addition of astaxanthin to fish feed had a very significant effect ( $P < 0.05$ ) on increasing the brightness of the colour of koi fish (*Cyprinus subfuscous*). However, it did not significantly affect the average growth (Average Body Weight / ABW), specific growth rate (SGR), Feed Conversion Ratio (FCR) and survival rate (SR). The best results were obtained at the additional dose of 125 mg/kg of feed with an average colour increase of  $20.29 \pm 0.10$ .

#### REFERENCES

- Andriani, Y., Wulandari, AP, Pratama, RI, & Zidni, I. (2019). Improving the Quality of Koi Fish (*Cyprinus carpio*) in the PBC Fish Farm Group in Cisaat District, Sukabumi. *Agrocreative Scientific Journal of Community Service*, 5 (1), 33-38. <https://doi.org/10.29244/agrokreatif.5.1.33-38>
- Collins, MA, Browne, CM, Edwards, TL, Ling, N., Tempero, GW, Gleeson, DM, Crockett, K., & Quaife, J. (2022). How low can they go: A comparison between dog (*Canis familiaris*) and environmental DNA detection of invasive koi carp (*Cyprinus rubrofusca*). *Applied Animal Behavior Science*, p. 255, 105729. <https://doi.org/https://doi.org/10.1016/j.applanim.2022.105729>
- Indarti, S., Muhaemin, M., & Hudaidah, S. (2012). Modified Toca Color Finder (M-TCF) and Chromatophore As Estimators of Comet Fish (*Carasius Auratus Auratus*) Color Brightness Levels That Are Fed With Different Proportions Of Shrimp Head Meal (TKU). *E-Journal of Aquaculture Engineering and Technology*, 1 (1).
- Irawati, T., Helilusatiningih, N., & Soenyoto, E. (2022). Innovation in Development of Integrated Field Laboratory Faculty of Agriculture Kadiri Islamic University East Java. *Formosa Journal of Applied Sciences*, 1 (6), 1051-1060.
- Iskandar, A., Amalia, D., Aji, HS, Hendriana, A., & Darmawangsa, GM (2021). Optimization of

- Cyprinus rubrofusca* Koi Fish Hatchery at Mina Karya Koi, Sleman, Yogyakarta. *SIGANUS: Journal of Fisheries and Marine Science*, 3 (1), 154–159.
- Kelabora, DM (2010). Effect of temperature on the survival and growth of carp (*Cyprinus carpio*) larvae. *Worst Fisheries Periodical*, 38 (1).
- Malini, DM, & Agustin, R. (2018). The Effect of Adding *Spirulina fusiformis* Flour to Feed on the Brightness of the Color of Koi Fish (*Cyprinus carpio* L). *Journal of Pro-Life*, 5 (2), 579–588.
- Maolana, V., Madyowati, SO, & Hayati, N. (2018). The effect of adding carrot juice (*Daucus carota* l) to feed on increasing colour at the enlargement of fish. Maolana, V., Madyowati, SO, & Hayati, N. (2018). Effect of adding carrot juice (*Daucus carota* l) in feed to increase. *Techno-Fish*, 1 (2), 78–85.
- Muslim, I., & Atjo, AA (2021). Hatching Response of Goldfish (*Cyprinus carpio*) Eggs at Different Temperature Levels. *SIGANUS: Journal of Fisheries and Marine Science*, 2 (2), 147–153.
- Nur, LA, Liliyanti, MA, & Kalih, LS (2020). The Influence of Natural Pigment Addition in Feed to Color Brightness and Growth of Juvenile Koi (*Cyprinus Carpio*). *Indonesian Journal of Aquaculture and Fisheries*, 2 (1), 40–43.
- Purba, M., Putriningtias, A., & Komariyah, S. (2021). Addition of Natural  $\beta$ -Carotene Flour in Feed on Increasing Color Brightness and Growth of Koi Fish (*Cyprinus carpio*). *Journal of Aquaculture, Teuku Umar University*, 4 (2), 10–20.
- Simamora, EK, Mulyani, C., & Isma, MF (2021). The Effect of Different Feeding on the Growth and Survival of Koi Goldfish (*Cyprinus Carpio*) Seeds. *Scientific Journal of the Aquatic Ocean*, 5 (1), 9–16.
- Simbolon, DFH, Sitanggang, LP, & Harefa, YM. (2002). *Addition of Marigold Flour (Agetes erecta) to Artificial Feed to Increase the Brightness of the Color of Koi Fish (Cyprinus carpio)*. *Budiyanto*, 65–71.
- Sitorus, AMG (2015). *The Effect of Concentration of Astaxanthin Flour in Feed on Increasing the Color of Goldfish (Carrasius auratus)*. University of Northern Sumatra.
- Sukarman, & Hirnawati, R. (2014). An alternative synthetic carotenoid (astaxanthin) to improve the colour quality of goldfish (*Carassius auratus*). *Widyariset*, 17 (3), 333–342.
- Yulianti, E., Maharani, H., & Diantari, R. (2014). The Effectiveness of Astaxanthin In Increasing the Brightness of Clown Fish (*Amphiprion ocellaris*) Color. *E-Journal of Aquaculture Engineering and Technology*, 3 (1), 313–318. <https://doi.org/10.23960/jrtbp.v3i1.468p313-318>



© 2023 by the authors. It was submitted for possible open-access publication under the terms and conditions of the Creative Commons Attribution (CC BY SA ) license (<https://creativecommons.org/licenses/by-sa/4.0/>).

21:12

4G



[JWS] Editor Decision

Inbox



**admin admin** 27 Apr

Amyda Suryati Panjaitan: We have reached a decision regarding your submission to



**Asep Mulyana** 2 days ago

to me



Amyda Suryati Panjaitan:

The editing of your submission, "Increased Color Brightness with The Addition of Astaxanthin in Koi Fish (Cyprinus Rubrofusca) Feed," is complete. We are now sending it to production.

Submission URL: <https://jws.rivierapublishing.id/index.php/jws/authorDashboard/submission/281>

Show quoted text

Reply

Reply all

Forward





jwsofficial@gmail.com

rivierapublishing.id

Perumahan Palm Asri Pasalakan Blok C No 5  
Kel. Pasalakan Kec. Sumber Kab. Cirebon

**ACADEMIC PAPER ACCEPTANCE LETTER**

Date: 10<sup>th</sup> April 2023  
Number: 070/RP-JWS/IV/2023

Dear (s), Amyda Suryati Panjaitan

**Title: INCREASED COLOR BRIGHTNESS WITH THE ADDITION OF ASTAXANTHIN IN KOI FISH (CYPRINUS RUBROFUSCUS) FEED**

After peer review process, your article has been provisionally accepted for publication in the **Journal of World Science**, in the forthcoming issue, Volume 2 Number 4, April 2023. All papers are published in the English language. All submitted manuscripts are subject to peer-review by the leading specialists for the respective topic.

Thus, this information is conveyed, it is hoped that cooperation during the publication process is expected.  
Thank you

Regards,  
Chief Editor



Vip Paramarta

Editorial Manager  
Journal of World Science  
Online ISSN 2828-8726 | Print ISSN 2828-9307  
<https://jws.rivierapublishing.id/index.php/jws/index>

International Indexed :



National Indexed:

