

The growth rate of Fenneropenaeus indicus in Seneboy Waters, South Sorong, Papua, Indonesia

by Cek Turnitin


Submission date: 24-May-2023 11:40AM (UTC+0800)

Submission ID: 2100552891

File name: 2021.1150-1162.pdf (979.96K)

Word count: 6454

Character count: 32642



The growth rate of *Fenneropenaeus indicus* in Seneboy Waters, South Sorong, Papua, Indonesia

¹Helman N. Yusuf, ¹Ali Suman, ²Achmad Suhermanto, ³Adnal Yeka, ³Indra Syamson, ³Ratih M. Rahmani, ³Yulia Fitri, ⁴Bongbongan Kusmedy, ⁴Erick Nugraha

¹Fishing Capture, Research Institute for Marine Fisheries Cibinong, Bogor, West Java, Indonesia; ²Faculty of Capture Fishing, Marine and Fisheries Polytechnic of Sorong, West Papua, Indonesia; ³Faculty of Capture Fishing, Marine and Fisheries Polytechnic of Pariaman, Padang, West Sumatera, Indonesia; ⁴Faculty of Fishing Technology, Jakarta Technical University of Fisheries, South Jakarta, Indonesia. Corresponding author: E. Nugraha, nugraha_eriq1@yahoo.co.id

Abstract. *Fenneropenaeus indicus* is one of the economically important species of shrimp cultivated in South Sorong Seneboy waters and currently its production tends to decrease. Studying its biological aspects provides data and information needed for assessing its exploitation and managing its conservation. This study aimed to examine the biological aspects of *F. indicus* as a material for shrimp fisheries management in the waters of Seneboy, South Kalimantan. The study was conducted between January-November 2016 in the waters of Seneboy. The results showed that the average size of the caught female *F. indicus* was 13.2-52.8 mm carapace length (CL), with a mode size of 11-33 mm CL, and the average size of male was 16-49.6 mm CL with a mode size of 29-31 mm CL. The sex ratio of male and female shrimp was 1:1.8. *F. indicus* spawning season is suspected to last all year and peak in February and October. The average size of the first time caught ($L_c=31.1$ CL) shrimp was smaller than the average size of the first time cooked gonads ($L_m=35$ mm), so that most of the shrimp caught were small and had not spawned. In female *F. indicus*, the growth rate (K), asymptotic carapace length (CL) and age at length 0 (t0) were of: 1.1 cm year⁻¹, 54.6 mm CL and -0.14883 years, respectively, while in male they were of: 1.0 cm year⁻¹, 52.6 mm and -0.136202 years, respectively. The natural mortality rate (M), death rate due to capture (F), total death rate (Z) and exploitation rate (E) of female rock lobsters were 2.64 cm year⁻¹, 1.53 cm year⁻¹, 2.64 cm year⁻¹ and 0.48 cm year⁻¹, respectively, while in male *F. indicus* amounted to 2.64 cm year⁻¹, 1.42 cm year⁻¹, 1.22 cm year⁻¹ and 0.56 cm year⁻¹, corresponding to an optimal utilization of the *F. indicus* resources. The status of *F. indicus* in Seneboy waters is "not fully exploited".

Key Words: biology aspect, shrimp fisheries management, sex ratio, spawning season.

Introduction. *Fenneropenaeus indicus* is one of the important commodities in the waters of Seneboy, South Sorong. *F. indicus* has economic value and contributes to the improvement of the fishermen's life in South Sorong. Seneboy fishermen catch *F. indicus* using a 4 m nylon trammel net as fishing gear, operated by 3 people from vessels of about 6 m in length, made of wood and fiber, of an average tonnage of 3 GT and a main engine power of 15 HP. The mesh size is of 1.5-7.25 inch, with a polyethylene head rope of Ø 8 mm. The length of the ground rope is 660 m. This effective and environmentally friendly fishing gear is used for catching *F. indicus* and other types of demersal fish which are bycatch (BRPL 2016). The catching time per trip ranges from 6 to 9 hours (from 07.00 to 16.00 hours, which is one day fishing) and the fishing ground ranges from 1 to 20 miles, with a distance of about 1-4 hours from the landing base. The shrimp fishing grounds are located around Wawonket Cape, Dua Island, Daram Island, Sibabu Cape, Komokara Cape, Sabuga Island, Sele Cape, Misool Islands, Sinabu Cape, Banana Island and Segeat Island. The shrimp

fishing season is 12 months with the highest season in July-November, with a catch of around 10-100 kg trip⁻¹. The production of shrimp and other crustaceans in South Sorong waters, in the period 2009 to 2016, reached 4,051 tons, with an annual average of 506.42 tons or 38.75% total production. The annual average production of *F. indicus* was 475.75 tons, with the highest production in 2010 of 557 tons and the lowest in 2013 of 350 tons (DKP Sorong 2017).

Shrimp catching activity in Seneboy, South Sorong, is the main support to meet the economic needs of trammel net fishermen in Temibuanan Regency. Catching shrimp at this location is carried out very intensively, so that fishing activities need to be supervised. The biology and population parameters provide the control and management criteria to the marine authorities. Research on several aspects of biology and population parameters was extensively conducted in several waters in Indonesia: Arafura (Naamin 1984), Cilacap (Suman 1992; Suman & Boer 2005; Hargiyatno et al 2013; Saputra & Subiyanto 2007), Strait Madura (Setyohadi et al 1999), Kotabaru (Suman & Umar 2010), Bone (Kembaren & Ernawati 2015), Tarakan (Kembaren & Suman 2013) and Meulaboh (Yusuf et al 2017).

The study provides information on the exploitation status of *F. indicus*, facing the market demand and an increasing commercial value. Rational management actions need to be taken in order to maintain a sustainable balance of the shrimp resources and the population parameters estimation is required as supporting data and information. The results of this study are expected to complement and update previous studies, underlying the management of *F. indicus* resources in the waters of Seneboy, South Sorong, and surroundings.

Material and Method

Research location and time. *F. indicus* sampling was carried out with a trammel net, obtained from fishermen and shrimp collectors in Seneboy Temibuana, South Sorong, from January to October 2016. The enumerators were taking samples using a random sampling method. The data collected included carapace length (CL), weight, sex, gonad maturity level and fishing area (Figure 1).



Figure 1. Fishing ground of trammel net in Seneboy, South Sorong (<https://www.mapsofworld.com/where-is/sorong.html>).

Data analysis. Biological data analysis included the size distribution, sex ratio, gonad maturity level, average size of first caught specimens (L_c). The analyzed population parameters were the growth pattern (CL_∞ , K), mortality parameters (Z , M , F) and utilization rate (E). The distribution of carapace length measurements is presented in the form of a bar chart with a length class of 2 mm. The level of maturity of female *F. indicus* gonads was observed visually based on Motosh (1981), namely by looking at the size of the shrimp ovary, located on its back. Shrimp sex ratios were analyzed using the Chi-square test (Walpole 1993).

The length at first matured (L_m) is obtained by entering the carapace length and P_{L_m} values into a graphical logistic function (King 1995), using the following equation:

$$P_{L_m} = \frac{1}{1 + \exp(aCL + b)}$$

Where:

P_{L_m} - logistic curve of proportion mature by length, L_m was determined by a/b ;

a - the intercept of regression;

b - the slope of regression;

CL - the length of shrimp carapace.

The value of carapace length caught for the first time (L_c) is obtained through a logistic function approach, with the Sparre & Venema (1992) equation:

$$S_{L_c} = \frac{1}{1 + \exp(a - b \cdot CL)}$$

Where:

S_{L_c} - logistic curve and L_c was determined by a/b ;

a - the intercept of regression;

b - the slope of regression;

CL - the length of shrimp carapace.

The method used in the study of population parameters is an analytical model based on the composition of the age structure of the shrimp (Sparre & Venema 1992). The growth parameters of Von Bertalanffy, namely the asymptotic carapace length (CL_∞) and the growth coefficient (K) were estimated by the ELEFAN 1 program in the FISAT II program (Gayanilo et al 2005). An estimation of the theoretical age (t_0) was carried out with the Pauly's (1983) empirical equation, namely:

$$\log(-t_0) = (-0.3922) - 0.2752 \log CL_\infty - 1.038 \log K$$

The value of the natural mortality (M) is estimated through the Pauly's equation, by including the effect of the mean water temperature (T) on the mortality rate, based on empirical observations (Pauly et al 1984):

$$\log M = (-0.0066) - 0.279 \log CL_\infty + 0.6542 \log K + 0.4634 \log T$$

The total mortality (Z) value was estimated using the length-converted catch curve method in the FISAT II program package (Pauly 1983; Gayanilo et al 2005). The capture mortality and exploitation rate are estimated using the Sparre & Venema (1992) equations:

$$E = \frac{F}{F + M}$$

$$E = \frac{F}{Z}$$

3

Where:

E - the exploitation rate;

F - the Fishing mortality;

M - natural mortality;

Z - the total mortality.

Results and Discussion

Biological aspects. A number of 2,085 *F. indicus* specimens were collected during the study were: 1,174 females (56.31%) and 911 males (46.69%). The carapace length of female *F. indicus* ranged from 13.2 to 52.8 mm, for a weight ranging from 9.6 to 88.4 g, with two modes in the ranges of 29-31 and 31.1-33 mm CL. The carapace length of male *F. indicus* ranged between 16-49.6 mm, for a weight ranging from 15.4 to 78.6 g, with a mode in the range of 29-31 mm CL (Figure 2).




Figure 2. Length frequency of *Fenneropenaeus indicus* in Seneboy waters.

The sex ratio determination of *F. indicus* in the waters of Seneboy South Sorong is based on the chi squared test, which indicates an imbalance between the male and female shrimp populations: in most cases, female shrimp dominate the catch. Although the average sex ratio of *F. indicus* males and females was 1:1.78, it showed a balanced value of 1:1 in February, March, September and October (Table 1).

Table 1
Sex ratio of *Fenneropenaeus indicus* in Seneboy waters, January-October 2016

Months	Sex ratio of <i>F. indicus</i>				X ²	Comparison
	Male	Female	M:F ratio			
2016)						P=5%
January	66	162	1	2.45	40.42	Not balanced
February	100	98	1	1.48	0.02	Balanced
March	94	106	1	1.61	0.72	Balanced
April	78	99	1	1.50	2.49	Not balanced
May	149	97	1	1.47	10.99	Not balanced
June	56	144	1	2.18	38.72	Not balanced
July	100	161	1	2.44	14.26	Not balanced
August	62	138	1	2.09	28.88	Not balanced
September	119	81	1	1.23	7.22	Balanced
October	87	88	1	1.33	0.01	Balanced
Total	911	1174	1	1.78	143.73	Not balanced

The highest proportion of mature female *F. indicus* (the maturity level of gonads III and the maturity level of gonads IV) was in January, February, September and October, with the peak maturing of gonads in October. The proportion of females with mature gonads in January, February, September and October was 60, 44, 46 and 40% (Figure 3).




Figure 3. Proportion of *Fenneropenaeus indicus* females with mature gonads in Seneboy waters, January–October 2016.

The size of the first time capture (L_c) corresponds to the probability level of 50% of the trammel net selectivity, meaning that the catch distribution of *F. indicus* can influence the availability of resources. The logistic function graph with a class interval of 2 mm CL indicates that the average value at the first capture of *F. indicus* is 31.1 mm CL, with an average value (L_m) of 35 mm CL at the first gonad maturation (Figure 4).




Figure 4. Average length at first *Fenneropenaeus indicus* capture with a trammel net.

Population parameters. The growth parameters of *F. indicus* in the Seneboy waters is processed from long frequency distribution data carapace by tracking any mode shift the distribut¹² of carapace length frequencies in a curved sequence of times Von Bertalanffy's growth. Based on the results of the ana⁴⁰is, it was found that infinite carapace length (L_∞) of ²⁷male *F. indicus* was 54.6 mm, the growth rate (K) was 1.1 year^{-1} and the male was 52.6 mm, the growth rate was 1.0 year^{-1} (Figure 5). From the value of these two parameters a growth equation and a key relationship can be created between carapace lengths with aged shrimp by using multiple variations in age value (t).




Figure 5. Growth model of male (A) and female (B) *Fenneropenaeus indicus* by ELEFAN I in Seneboy waters.

The asymptotic carapace length (L_{∞}) of female *F. indicus* was 54.6 mm CL and 52.6 mm CL for males, while the age of female *F. indicus* at 0 (t_0) was -0.14883 years and -0.136202 for males. So that the equation for the Von Bertalanffy equation for female *F. indicus* is $L_t = 54.6 (1 - e^{-1.100(t - 0.14883)})$ and for male it is $L_t = 52.6 (1 - e^{-1.000(t - 0.136202)})$ (Figure 6). The maximum size range of *F. indicus* is estimated at the age of 1.8 years or around 20 months. The average size size of 31.1 mm CL in caught *F. indicus* (L_c) was estimated at the age of 7-8 months. The average mature gonad size in females (L_m) was of 35 mm CL, at the age of 9-10 months.




Figure 6. Von Bertalanffy curve of *Fenneropenaeus indicus* in Seneboy waters.

The total mortality rate (Z) of female and male *F. indicus* based on the length-converted catch curve is 3.16 year^{-1} , with a natural mortality rate (M) of 1.64 year^{-1} and the mortality rate due to fishing (F) of 1.52 year^{-1} . The total mortality rate of male *F. indicus* was 2.64 year^{-1} , the natural mortality rate was 1.42 year^{-1} and the mortality rate due to fishing was 1.22 year^{-1} . Based on these mortality parameters, the utilization rate of *F. indicus* was 0.48 for females and 0.46 for males (Figure 7).




Figure 7. Growth curve of male (A) and female (B) *Fenneropenaeus indicus* in waters of Seneboy.

Discussion. The size distribution of *F. indicus* caught in the study area was later than in Kaimana waters, ranging from 25–49 mm, and the average size was 28.98 mm for males and 34.96 for females. The size of *F. indicus* specimens caught in Meulaboh waters ranged between 17.8 and 79.54 mm for females and between 18.3 and 64.36 mm for males. The size of *Penaeus merguensis* in Central Java waters ranged between 20 and 62 mm for males and between 14 and 68 mm for females, and averaged 17 to 47 mm in West Java waters (Tirtadanu & Panggabean 2018; Yusuf 2017; Suman et al 1988). Differences in shrimp size at several locations can be caused by different gears, environmental conditions and fishing pressure (Olin et al 2017; Wilson et al 2010; Matthews 1982; Ogbonna 2001). Given the larger size in Seneboy waters, it is supposed that the species is not over-utilized.

The sex ratio of male and female shrimp in the study area was not balanced: the females dominated the catch. The unbalanced sex ratio is similar to the *F. indicus* caught in Meulaboh waters (Yusuf 2017; Saputra 2008; Saputra et al 2013; Wedjadmiko & Yulianti 2003; Wedjadmiko 2009; Suman et al 1991), suggesting that the fishing pressure of shrimp resources in Seneboy waters has not disturbed the population renewal so that the sustainability of shrimp stocks is still well maintained.

The peak indicated that a higher frequency of mature gonads in *F. indicus* occurred in February and October, suggesting the spawning seasons for the shrimp (Martosubroto 1978). For comparison, the spawning peaks occurred in November for several types of Penaeid shrimp from the waters of West Kalimantan (including *P. merguensis*), in January and August for the *Metapenaeus monoceros* from the Southern Java waters and from September to October for the *M. monoceros* from the Gulf of Carpentaria, Australia (Kembaren & Suman 2013; Suman et al 2005; Crocos et al 2000; Pillai & Thirumilu 2013; Gerami et al 2013). The spawning season for *F. indicus* in the study area lasted throughout the year, with peaks in February and October. Therefore, from the perspective of the

resources sustainability, it is preferable to suspend the *F. indicus* fishing activities in Seneboy waters during February and October.

The first catch average size (Lc) of *F. indicus* in the study area was of 31.1 mm CL and the mean value at the first gonad maturity was 35 mm CL. The results of research conducted in the waters of the Cilacap Segara Anakan Lagoon, using floating nets, showed an Lc value of 20.2 mm, for a total length of about 90 mm (Saputra 2008). Chan (1998) stated that *F. indicus* females can reach a total length of 230 mm, although they generally measure less than 170 mm. These results are different from the research conducted in the waters of North Central Java, where the carapace length of *P. merguensis* was of 29.4 mm (Tirtadanu & Ernawati 2016), the carapace length in the Mayangan waters was 28.9 mm (Wedjatmiko & Yulianti 2003) and the Lc in the Dolak waters, Arafura Sea was 28.78 mm (Hargiyatno & Sumiono 2012). The differences of Lc values are not only caused by the mesh size of the net used, but they are also due to the operations duration and to the location of the fishing ground. The differences of Lc values are not only caused by the mesh size of the net used, but they are also due to the operations duration and to the location of the fishing ground. According to Susetiono & Setyono (1990), the *P. merguensis* from shallow waters tends to be smaller than in deeper waters (e.g. Cilacap), with an average size (Lc) of 51 mm CL (Saputra et al 2013). The shrimp size differences at several locations can be caused by different gears, environmental conditions and fishing pressure (Olin et al 2017; Wilson et al 2010).

According to FAO (2008), the Lm value is 50% of *F. indicus*, at a total length of 130-149 mm. This shows that the value of $L_{50\%} < Lm_{50\%}$ means that *F. indicus* recruitment is threatened by overfishing. According to Teikwa & Mgaya's (2003) research, the *F. indicus* carapace length size of the first caught males is of 34 mm and 39 mm carapace in female shrimps. The results of a study conducted by Melmammblessy (2011) indicated that in the Arafura waters, the average size of the first catch (Lc) had a carapace length of 60.26 mm. The average size of the *F. indicus* caught for the first time (Lc) in the study area was smaller than the average size at the first maturity of gonads (Lm), indicating that most of the shrimp caught had not spawned. In the long term, this situation is detrimental to the population's sustainability, hampering the shrimp's recruitment process in the Seneboy waters. In this regard, it is necessary to adjust the size of the mesh in such a manner that the smallest carapace length of the caught *F. indicus* cannot be under 13.2 mm CL.

The growth rate (K) of *F. indicus* in the research area shows that they have a fast growth rate: 1.1 year^{-1} in females and 1.0 year^{-1} in males, with an asymptotic carapace length (L_{∞}) of 54.6 mm CL for females and 52.6 mm CL for males. The growth parameters in different locations were: $L_{\infty}=35.7 \text{ mm}$ and $K=1.26 \text{ year}^{-1}$ in the waters of the Segara Anakan Lagoon Cilacap, 1.4 year^{-1} and $L_{\infty} 44.3 \text{ mm}$ in the Kota Baru waters and 44.7 mm CL for males and 51.25 mm CL for females in the Kaimana waters (Table 2).

The mortality rate (Z) of *F. indicus* in the Seneboy waters is relatively low, namely 3.16 in males and 2.6 in females. Some *Penaeid* shrimp in other waters, namely the waters of the Cilacap Segara Anakan Lagoon, obtained an M value of 2.6 cm year^{-1} , an F value of $1.35 \text{ cm year}^{-1}$ (*P. merguensis* in Segara Anakan Lagoon Cilacap is 1.43 year^{-1} , the natural mortality rate is $1.85 \text{ cm year}^{-1}$ and catching $2.32 \text{ cm year}^{-1}$, Madura waters natural mortality of $2.17-2.22 \text{ cm year}^{-1}$ and catching $1.11-1.56 \text{ cm year}^{-1}$ (Saputra 2008; Kembaren et al 2012; Setyohadi et al 1999; Hediando et al 2016).

The highest mortality rates (ind year^{-1}) were recorded: 9 for males and 9.47 for females in the waters of Kota Baru, 10.58 in the Kakinada waters, 7.01 in the Khoozestan waters and 5.78 in the Vietnamese waters (Tirtadanu et al 2017; Devi 1987; Ansari et al 2014; Dinh et al 2010). The natural mortality is influenced by several factors such as food availability, disease, environment, competition and the presence of predatory species (Sparre & Venema 1992; Niamaimandi et al 2007). *Penaeid* prawn is the prey of several demersal fish, such as *Lutjanus gibbus*, *Eleutheronema tetradactylum* and *Arius* sp. (Bachok et al 2004; Titrawani et al 2013). Death due to fishing (F) of both male and female *F.*

indicus is lower than the natural mortality rate (M), indicating that the *F. indicus* population has not been exploited intensively by fishermen in the waters of Seneboy, West Papua. Garcia (1988) suggests that the average natural mortality rate (M) of penaeid shrimp is $2.4 \pm 0.3 \text{ year}^{-1}$ for adult shrimp. Naamin (1984) found that the F value of *P. merguensis* in Arafura waters varies according to the development of the fishing effort, ranging from 0.55 year^{-1} (at the start of the exploitation) to 8.99 year^{-1} .

37

Table 2

Asymptotic length (L_{∞}), growth rate (K) and exploitation rate (E) of *Fenneropenaeus indicus* in some areas

Location	Sexes	L_{∞} (mm CL)	K	t0	Authors
Cilacap, Central Jawa	Combined	35.7	1.26	-	Saputra (2008)
Kota Baru, South Kalimantan	Combined	44.3	1.4	-	Suman et al (2010)
Kaimana, West Papua	Combined	44.7	1.38	-	Tirtadanu & Pangabebean (2018)
Cilacap, Central Jawa	Male	40.7	1	0.56	Suman & Prisantosa (2017)
	Female	54.2	1.1	0.36	
Tarakan, North Kalimantan	Male	45.2	1.55	0.76	Chodriyah & Suman (2017)
	Female	57.6	1.33	0.76	
Sampit, Central Kalimantan	Combined	57.8	1.45	0.66	Nurdin & Kembaren (2015)
Tanah Laut, South Kalimantan	Combined	55	1.05	0.74	Suman et al (2017)
	Male	44.5	1.05	0.66	
Cenderawasih Bay, Papua	Female	48.7	1.15	0.55	Kembaren & Ernawati (2015)
	Male	44.7	1.4	0.49	
Kaimana, West Papua	Female	51.25	1.37	0.44	Tirtadanu & Pangabebean (2018)
	Male	54.6	1.1	0.148	
Seneboy, West Papua	Male	54.6	1.1	0.148	Present research
	Female	52.6	1.0	0.136	

In this study, the exploitation rates (E) of female and male *F. indicus* were 0.48 and 0.46. This condition explains that the fishing rate is greater than the natural mortality rate. The optimum exploitation rate stated by Gulland (1983) is 0.5, so that the rate of exploitation of *F. indicus* in Seneboy waters is close to the optimum value and is still sustainable. The level of exploitation of *F. indicus* in several areas has exceeded the optimum limit, namely in the waters of the Segara Anakan Cilacap Lagoon, Meulaboh waters, Tarakan Bay, Tanah Laut and Cenderawasih, where the level of exploitation indicates overfishing (Saputra 2008; Yusuf et al 2017; Chodriyah & Suman 2017; Suman et al 2017; Kembaren & Ernawati 2015). The exploitation of *F. indicus* has approached the maximum, suggesting that a reduction of the fishing pressure is required, with at least 5% of the current total effort.

Conclusion. The average size of *F. indicus* caught in the waters of Seneboy South Sorong is 16-49.6 mm CL for the males and 13.2-52.8 mm for the females. The sex ratio of *F. indicus* is unbalanced: 1:1.8, male against female. The spawning season lasts throughout the year with peaks in February and October. The catch is dominated by adults so that management sets the regulations for the minimum legal size. These results concluded that the arrest of *F. indicus* can be continued by arranging the arrest effort and stipulating a minimum legal size of 35 mm CL. This indicates that the level of exploitation of *F. indicus* in Seneboy waters is not yet optimal, but leads to overfishing. In order to maintain the sustainability of the *F. indicus* resources in the waters of Seneboy, West Papua, it is advisable (1) to reduce efforts with about 5% from the current level, by adjusting the mesh in such a manner that the retained catch size of *F. indicus* will be of at least 13.2 mm CL and (2) to suspend the *F. indicus* fishing operations in February and October.

Acknowledgements. This paper is part of the research activity "Biological Characteristics of Fisheries, Resource Habitat and Production Potential of Fish Resources in FMA 715 Tomini

Bay 2017* which was organized by the Muara Baru Jakarta Marine Fisheries Research Institute.

Conflict of interest. The authors declare no conflict of interest.

References

- Ansari H., Khodadadi M., Eskandari G., 2014 Growth and mortality parameters of Jingga shrimp (*Metapenaeus affinis* H. Milne Edwards, 1837) in western coast of Khoozestan Province. *Journal of Marine Biology* 6(23):1-11.
- Bachok Z., Mansor M. I., Noordin R. M., 2004 Diet composition and food habits of demersal and pelagic marine fishes from Terengganu Waters, East Coast of Peninsular Malaysia. *NAGA World Fish Center Quarterly* 27(3):41-47.
- Chan T. Y., 1998 Shrimps and shrimps. In: *FAO species identification guide for or fishery purposes. The living marine resources of the Western Central Pacific. Cephalopods, crustaceans, holothurians and sharks.* Carpenter K. E., Niem V. H. (eds), pp. 851-971, FAO, Rome.
- Chodriyah U., Suman A., 2017 [Some population parameters of white shrimp (*Penaeus merguensis demann*) in Tarakan Waters, North Kalimantan]. *Bawal Widya Research* 9(2):85-92. [In Indonesian].
- Crocos P. J., Park Y. C., Die D. J., Warburton K., Manson F., 2000 Reproductive dynamics of endeavor shrimps, *Metapenaeus endeavouri* and *M. ensis*, in Albatross Bay, Gulf of Carpentaria, Australia. *Marine Biology* 138:63-75.
- Devi S. L., 1987 Growth and population dynamics of three penaeid shrimps in the trawling grounds off Kakinada. *Indian Fish Journal* 34(2): 245-264.
- Dinh T. D., Moreau J., Van M. V., Phuong N. T., Toan V. T., 2010 Population dynamics of shrimps in littoral marine waters of the Mekong Delta, South of Vietnam. *Pakistan Journal of Biological Sciences* 13(14):683-690.
- Garcia S., 1988 Tropical penaeids shrimps. In: *Fish population dynamics.* Gulland J. A. (ed), pp. 219-249, John Wiley & Sons Ltd.
- Gayanilo F. C. Jr., Sparre P., Pauly D., 2005 FAO/ICLARM stock assessment tools II (FISAT II). User's guide. *FAO Computerized Information Series (Fisheries) No. 8*, FAO, Rome.
- Gerami M. H., Ghorbani R. S., Paighmabari Y., Momeni M., 2013 Reproductive season, maturation size (Lm50) and sex ratio of *Metapenaeus affinis* (Decapoda: Penaeidae) in Hormozgan shrimp fishing grounds, South of Iran. *International Journal of Aquatic Biology* 1(2):48-54.
- Gulland J. A., 1983 *Fish stock assessment. A manual of basic method.* FAO/Wiley Series on Food and Agriculture, Rome, 241 p.
- Hargiyatno I. T., Sumiono B., 2012 [Density of stock and biomass resources of tiger prawns (*Penaeus semisulcatus*) and dogol (*Metapenaeus endeavouri*) in Aru sub-area, Arafura Sea]. *Indonesian Fisheries Research Journal* 18(1):14-25. [In Indonesian].
- Hargiyatno I. T., Sumiono B., Suharyanto, 2013 [Capture rate, stock density and several aspects of the biology of the jermal shrimp (*Penaeus merguensis*) in Dolak waters, Arafura Sea]. *Bawal* 5(2):123-129. [In Indonesian].
- Hedianto D. A., Suryandari A., Tjahjo D. W., 2016 [Population dynamics and utilization status of *Penaeus monodon* shrimp in the waters of East Aceh, Province]. *Indonesian Fisheries Research Journal* 22(2):71-82. [In Indonesian].
- Kembaren D. D., Sumiono B., Suprpto, 2012 [Biology and growth parameter of white shrimp (*Penaeus merguensis*) in South Sulawesi]. In: [Exploitation status of fishery resources in Makasar Strait, Bone Bay, Flores Sea and Banda Sea]. Suman A., Wudianto, Sumiono B. (eds), IPB Press, Bogor, 300 p. [In Indonesian].
- Kembaren D. D., Ernawati T., 2015 [Population dynamics and estimated spawning potential ratios of white shrimp (*Penaeus merguensis* deMann, 1907) in the waters of the

- Cendrawasih Bay and its surroundings, Papua]. Indonesia Fisheries Resources Journal 21(3):201-210. [In Indonesian].
- Kembaren D. D., Suman A., 2013 Biology and population dynamics of white shrimp (*Penaeus merguensis*) in the Tarakan waters, East Borneo. Indonesian fisheries research journal. Research Center for Fisheries Management and Conservation of Fish Resources 19(2):99-105.
- King M. G., 1995 Fishery biology, assessment and management. Fishing News Books, United Kingdom, 341 p.
- Martosubroto P., 1978 [Spawning season and growth of Jerbung shrimp (*Penaeus merguensis* de Man) and dogol shrimp (*Metapenaeus ensis* de Haan) in the waters of Tanjung Karawang]. Proceedings of Seminar II on Shrimp Fisheries, pp. 7-20. [In Indonesian].
- Matthews G. A., 1982 Relative abundance and size distributions of commercially important shrimp during the 1981 Texas closure. Marine Fisheries Review 44:5-15.
- Melmammblessy F. D., 2011 [Size of the first time ripe white shrimp gonads (*Penaeus merguensis* de Man (1988) in Arafura Sea in Naukenjerai District, Merauke Regency]. Agribusiness and Fisheries Scientific Journal (AGRIKAN UMMU-Ternate) 4(2):75-81. [In Indonesian].
- Motoh H., 1981 Studies on the fisheries biology of the giant tiger shrimp, *Penaeus monodon*, in the Philippines. SEAFDEC Technology, Paper No. 7, 128 p.
- Naamin N., 1984 [Population dynamics of white shrimp (*Penaeus merguensis* de Man) in Arafura waters and alternative management]. PhD Thesis, Postgraduate Faculty, IPB Bogor, Indonesia, 277 p. [In Indonesian].
- Niamaimandi N., Arshad A. B., Daud S. K., Saed R. C., Kiabi B., 2007 Population dynamic of green tiger shrimp, *Penaeus semisulcatus* (De Haan) in Bushehr coastal waters, Persian Gulf. Fisheries Research 86:105-112.
- Ogbonna J. C., 2001 Reducing the impact of tropical shrimp trawling fisheries on the living marine resources through the adoption of environmentally friendly techniques and practices in Nigeria. Tropical shrimp fisheries and their impact on living resources. FAO Fisheries Circular, No. 974, Fii/C974, pp. 188-215.
- Olin M., Tiainen J., Rask M., Vinni M., Nyberg K., Lehtonen H., 2017 Effects of nonselective and size-selective fishing on perch populations in a small lake. Boreal Environment Research 22:137-155.
- Pauly D., 1983 Some simple methods for the assessment of tropical fish stocks. FAO Fisheries Technical Paper, 254 p.
- Pauly D., Ingles J., Neal R., 1984 Application to shrimp stocks of objective methods for the estimation of growth, mortality and recruitment-related parameters from length-frequency data (ELEFAN I and II). Penaeid shrimps-Their biology and management. Fishing News Books Ltd., Farnham-Surrey-England, pp. 220-234.
- Pillai S. L., Thirumilu P., 2013 Fishery and stock assessment of *Metapenaeus dobsoni* Miers, 1878) off Chennai. Indian Journal of Geo-Marine Sciences 42(4):448-452.
- Saputra S. W., 2005 Population dynamics of finger shrimps (*Metapenaeus elegans* De Mann 1907) and their management in the Segara Anakan lagoon of Cilacap, Central Java. PhD Thesis, Bogor Agricultural Institute, Bogor, 235 p.
- Saputra S. W., 2008 [Population dynamic of "dogol" shrimp (*Penaeus indicus* H. Milne Edwards 1837) in Segara Anakan Lagoon Cilacap Central Java]. Journal of Fisheries Sciences 10(2):213-222. [In Indonesian].
- Saputra S. W., Djuwito, Rutiyaningsih A., 2013 [Several aspects of the biology of the white shrimp (*Penaeus merguensis*) in the waters of the Cilacap coast, Java]. Journal of Management of Aquatic Resources 2(3):47-55. [In Indonesian].
- Saputra S. W., Subiyanto, 2007 [Population dynamics of white shrimp (*Penaeus merguensis* de Mann 1907) in the Segara Anakan Lagoon, Cilacap, Central Java]. Marine Science, Diponegoro University 12(3):157-166. [In Indonesian].

- Setyohadi D., Nugroho D., Lelono T. J., Wiadnya D. G. R., Martinus, 1999 [Biology and distribution of penaeid shrimp resources based on the catch in the Madura Strait]. Agricultural Research and Development Report, pp. 50-61. [In Indonesian].
- Sparre P., Venema S. C., 1992 Introduction to tropical fish stock assessment. FAO Fisheries Technical Paper, No. 306/1, Rome, 376 p.
- Suman A., 1992 Dynamics of dogol shrimp (*Metapenaeus ensis* de Haan) in the southern coastal waters of Java. Proceedings of the Seminar on Marine and Coastal Ecology I, pp. 64-71.
- Suman A., Boer M., 2005 Size at first maturity, spawning season and growth parameters of endeavor shrimp (*Metapenaeus ensis* de Haan) in Cilacap and adjacent waters. Indonesia Fisheries Resources Journal 11(2):65-71.
- Suman A., Hasanah A., Ernawati T., Pane A. R., 2017 The population dynamic of banana shrimp (*Penaeus merguensis* De Man) in Tanah Laut Waters, South Kalimantan. Indonesian Fisheries Research Journal 23(1):17-22.
- Suman A., Iskandar, Bachelor B., 1991 Biological, catching and economic aspects of shrimp fisheries in East Sumba Waters, East Nusa Tenggara. Marine Fisheries Journal 57:119-129.
- Suman A., Prisantoso B. I., Bintoro G., 2005 Population dynamic of endeavor shrimp (*Metapenaeus elegans*) in the waters of South Coast of Java. Indonesia Fisheries Resources Journal 13(1):49-54.
- Suman A., Sumiono B., Rijal M., 1988 Several aspects of the biology of the white shrimp (*Penaeus merguensis* De Man) in Panimbang waters, West Java. Indonesian Fisheries Research Journal 49:13-19.
- Suman A., Umar C., 2010 [Population dynamics of white shrimp (*Penaeus merguensis* de Mann) in Kotabaru waters, South Borneo]. Indonesia Fisheries Resources Journal 16(1):29-33. [In Indonesian].
- Susetiono, Setyono D., 1990 [Some information on the biology of white shrimp (*Penaeus merguensis* de Man) in the waters of East Seram Kufar]. Research Institute for Marine Resources Development, Ambon, 7 p. [In Indonesian].
- Teikwa E. D., Mgaya Y. D., 2003 Abundance and reproductive biology of the Penaeid shrimps of Bagamoyo coastal waters, Tanzania, Western Indian Ocean. Marine Science Journal 2(2):117-126.
- Tirtadanu, Ernawati T., 2016 [Biological studies of white shrimp (*Penaeus merguensis* de Man, 1888) in the waters of North Central Java]. Capture Fisheries Research Widya 8(2):109-116. [In Indonesian].
- Tirtadanu, Panggabean A. S., 2018 Catch rate and population parameters of banana shrimp *Penaeus merguensis* in Kaimana waters, West Papua, Indonesia. AACL Bioflux 11(4):1378-1387.
- Tirtadanu, Surapto, Suman A., 2017 [Distribution of length frequency, length-weight relationship, gonad maturity level and average size of first mature white shrimp gonads (*Penaeus merguensis*) in Kota Baru waters, South Kalimantan]. UNDER 9(3):145-152. [In Indonesian].
- Titrawani, Elvyra R., Sawalia R. U., 2013 A[nalysis of stomach contents of pleased fish (*Eleutheronema tetradactylum* Shaw) in Dumai waters]. Al-Kauniyah Journal of Biology 6(2):85-90. [In Indonesian].
- Walpole R. V. E., 1993 [Introduction to statistics B]. PT Gramedia, Jakarta, 321 p. [In Indonesian].
- Wedjatmiko, 2007 [Distribution and density of Mantis shrimp (*Carinosquilla spinosa*) in Arafura waters]. Indonesia Fishery Research Journal 13(1):61-69. [In Indonesian].
- Wedjatmiko, 2009 Catches and biological aspects of banana shrimp (*Penaeus* sp.) in the waters of West Aceh. Indonesia Fishery Research Journal 15(2):133-140.

- Wedjatmiko, Yulianti, 2003 [Several aspects of the biology of the white shrimp (*Penaeus merguensis*) in Mayangan Waters, North Coast of West Central Java]. *Journal of Indonesian Fisheries Research* 9(3):27-34. [In Indonesian].
- Wilson S. K., Fisher R., Pratchett M. S., Graham N. A. J., Dulvy N. K., Turner R. A., Cakacaka A., Polunin N. V. C., 2010 Habitat degradation and fishing effects on the size structure of coral reef fish communities. *Ecological Applications* 20(2):442-451.
- Yusuf H. N., Suman A., Hidayat T., 2017 [Some population parameters of banana shrimp (*Penaeus Indicus* H. Milne Edward, 1837) in Meulaboh Waters]. *Crustacean National Symposium Proceedings*, pp. 1-13. [In Indonesian].
- *** BRPL Research Institute of Marine Fisheries, 2016 [Fisheries research characteristics of biology, habitat and resource potential for production in FMA-715 (Tomini, Maluku Sea, Seram Sea, Halmahera Sea and the Berau Gulf)]. Ministry of Marine Fisheries, Indonesia, 206 p. [In Indonesian].
- *** DKP Sorong, 2017 [Annual report]. Sorong Marine and Fisheries Service, West Papua, 64 p. [In Indonesian].
- *** FAO, 2008 The state of world fisheries and aquaculture 2008. Food and Agriculture Organization of the United Nations, Rome, pp. 10-15.
- *** Seneboy, South Sorong Map, <https://www.mapsofworld.com/where-is/sorong.html>.

Received: 15 January 2021. Accepted: 21 April 2021. Published online: 04 May 2021.

Authors:

Helman Nur Yusuf, Research Institute for Marine Fisheries, Fishing capture, Jl. Raya Bogor KM. 47, Nanggewer Mekar, Cibinong, 16912 Bogor, West Jawa, Indonesia, e-mail: helmankkp183@gmail.com

All Suman, Research Institute for Marine Fisheries, Fishing capture, Jl. Raya Bogor KM. 47, Nanggewer Mekar, Cibinong, 16912 Bogor, West Jawa, Indonesia, e-mail: alisuman_62@yahoo.com

Achmad Suhermanto, Marine and Fisheries Polytechnic of Sorong, Faculty of Capture Fishing, Jl. Kapitan Pattimura Tanjung Kasuari Suprau, 98411 Sorong, West Papua, Indonesia, e-mail: ac.suhermanto.apsor@gmail.com

Adnal Yeka, Marine and Fisheries Polytechnic Pariaman, Faculty of Capture Fishing, Jl. Simpang Toboh V Koto, Kampung Dalam, Pariaman Tengah, 25514 Pariaman City, West Sumatera, Indonesia, e-mail: adnalyeka@gmail.com

Indra Syamson, Marine and Fisheries Polytechnic Pariaman, Faculty of Capture Fishing, Jl. Simpang Toboh V Koto, Kampung Dalam, Pariaman Tengah, 25514 Pariaman City, West Sumatera, Indonesia, e-mail: Indra syamson@gmail.com

Ratih Martia Rahmani, Marine and Fisheries Polytechnic Pariaman, Faculty of Capture Fishing, Jl. Simpang Toboh V Koto, Kampung Dalam, Pariaman Tengah, 25514 Pariaman City, West Sumatera, Indonesia, e-mail: ratihmartiarahmani@gmail.com

Yulia Fitri, Marine and Fisheries Polytechnic Pariaman, Faculty of Capture Fishing, Jl. Simpang Toboh V Koto, Kampung Dalam, Pariaman Tengah, 25514 Pariaman City, West Sumatera, Indonesia, e-mail: yulia_caffa@yahoo.co.id

Bongbongan Kusmedy, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, Jl. AUP Pasar Minggu, 12520 South Jakarta, Indonesia, e-mail: bkhutapea@gmail.com

Erick Nugraha, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, Jl. AUP Pasar Minggu, 12520 South Jakarta, Indonesia, e-mail: nugraha_eriq1@yahoo.co.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Yusuf H. N., Suman A., Suhermanto A., Yeka A., Syamson I., Rahmani R. M., Fitri Y., Kusmedy B., Nugraha E., 2021 The growth rate of banana shrimp (*Penaeus indicus*) in Seneboy Waters, South Sorong, Indonesia. *AAFL Bioflux* 14(3):1150-1162.

The growth rate of Fenneropenaeus indicus in Seneboy Waters, South Sorong, Papua, Indonesia

ORIGINALITY REPORT

12%

SIMILARITY INDEX

8%

INTERNET SOURCES

10%

PUBLICATIONS

4%

STUDENT PAPERS

PRIMARY SOURCES

- 1** Submitted to Universitas Muhammadiyah Purwokerto
Student Paper 2%
- 2** text-id.123dok.com
Internet Source 1%
- 3** ftp.academicjournals.org
Internet Source <1%
- 4** A Sofian, C Kusmana, A Fauzi, O Rusdiana. "The Carrying Capacity of Ecosystem Services of Mangrove Angke Kapuk Area, Jakarta Bay", IOP Conference Series: Earth and Environmental Science, 2019
Publication <1%
- 5** Paula S. Haynes, Patricia Browne, Liam Fullbrook, Conor T. Graham et al. "Growth in Nephrops norvegicus from a tag-recapture experiment", Scientific Reports, 2016
Publication <1%
- 6** Renato Luiz BOT NETO, Barbara Maichak de CARVALHO, Roberto SCHWARZ JÚNIOR, Henry <1%

Louis SPACH. "INSIGHTS ON THE INFLUENCE OF PHYLOGENETIC ON THE RELATIVE GROWTH USING *Atherinella brasiliensis* AS A TOOL", Boletim do Instituto de Pesca, 2021

Publication

7	www.inape.gub.uy Internet Source	<1 %
8	Wang, T., D. Huang, Y. Zhao, H. Wang, S. Hu, and J. Shen. "Age, growth and mortality of invasive sharpbelly, <i>Hemiculter leucisculus</i> (Basilewski, 1855) in Erhai Lake, China", <i>Journal of Applied Ichthyology</i> , 2013. Publication	<1 %
9	digital.ncdcr.gov Internet Source	<1 %
10	sportdocbox.com Internet Source	<1 %
11	dli.ejournal.unri.ac.id Internet Source	<1 %
12	ejournal.unmus.ac.id Internet Source	<1 %
13	www.mdpi.com Internet Source	<1 %
14	ia800204.us.archive.org Internet Source	<1 %

15

Submitted to Alexandru Ioan Cuza University
of Iasi

Student Paper

<1 %

16

www.waterboards.ca.gov

Internet Source

<1 %

17

Ana Paula Pereira Gomes, Karina Annes
Keunecke, Demarques Ribeiro da Silva,
Marcelo Vianna. "Modulating reproduction of
Penaeidae shrimps: ecological responses of
two sympatric species (Decapoda:
Dendrobranchiata) on south-eastern Brazilian
coast", Journal of the Marine Biological
Association of the United Kingdom, 2012

Publication

<1 %

18

B. A. Bluhm, T. Brey. "Age determination in
the Antarctic shrimp *Notocrangon antarcticus*
(Crustacea: Decapoda), using the
autofluorescent pigment lipofuscin", Marine
Biology, 2001

Publication

<1 %

19

C.E. Gabche, H.-U.P. Hockey. "Growth,
mortality and reproduction of *Sardinella*
maderensis (Lowe, 1841) in the artisanal
fisheries off Kribi, Cameroon", Fisheries
Research, 1995

Publication

<1 %

20

M. A. A. Pinheiro, G. Y. Hattori. " Growth of the speckled swimming crab, (Lamarck, 1818) (Crustacea, Brachyura, Portunidae), in Ubatuba (SP), Brazil ", Journal of Natural History, 2006

Publication

<1 %

21

Sri Puryono, Suryanti Suryanti. "Gastropod Diversity in Mangrove Forests of Mojo Village, Ulujami District, Pemalang Regency, Indonesia", Journal of Ecological Engineering, 2019

Publication

<1 %

22

ubm.opus.hbz-nrw.de

Internet Source

<1 %

23

123dok.com

Internet Source

<1 %

24

Ali Suman, Ap'idatul Hasanah, Tri Ernawati, Andina Ramadhani Putri Pane. "THE POPULATION DYNAMIC OF BANANA PRAWN (Penaeus merguensis de Man) IN TANAH LAUT WATERS, SOUTH KALIMANTAN", Indonesian Fisheries Research Journal, 2017

Publication

<1 %

25

Ali Suman, Duranta Diandria Kembaren, Muhammad Taufik. "BEBERAPA ASPEK BIOLOGI UDANG JERBUNG (Penaeus merguensis) DI PERAIRAN KEPULAUAN ARU

<1 %

DAN SEKITARNYA (LAUT ARAFURA) SEBAGAI
DASAR KEBIJAKAN PENGELOLAANNYA
SECARA BERKELANJUTAN", Jurnal Kebijakan
Perikanan Indonesia, 2022

Publication

26

Helman Nur Yusuf, Ali Suman, Thomas
Hidayat, Anthony Sisco Panggabean.
"PARAMETER POPULASI LOBSTER BAMBU
(Panulirus versicolor) DI PERAIRAN
SIMEULUE", BAWAL Widya Riset Perikanan
Tangkap, 2018

Publication

<1 %

27

Qiu, D.J.. "Investigation on the microstructure
and optical properties of c-axis oriented
nanocrystalline hexagonal BN films fabricated
by hot-filament chemical vapor deposition",
Thin Solid Films, 20050722

Publication

<1 %

28

Tien Thi Kieu Nguyen, Quang Minh Dinh, Nam
Sy Tran, Ton Huu Duc Nguyen. "Stock
assessment of two populations of
Glossogobius sparsipapillus (Osteichthyes,
Gobiidae) in the Mekong Delta", The Egyptian
Journal of Aquatic Research, 2021

Publication

<1 %

29

Tirtadanu Tirtadanu, Umi Chodriyah. "LAJU
TANGKAP, KARAKTERISTIK BIOLOGI DAN
STATUS PEMANFAATAN UDANG JERBUNG

<1 %

(*Penaeus merguensis* DE MANN, 1988) DAN UDANG DOGOL (*Metapenaeus affinis* H. MILNE EDWARDS, 1837) DI PERAIRAN CILACAP", Jurnal Penelitian Perikanan Indonesia, 2020

Publication

30

Umi Chodrijah, Ria Faizah. "BEBERAPA ASPEK BIOLOGI UDANG WINDU (*Penaeus monodon* (Fabricus, 1789) DI PERAIRAN TARAKAN, KALIMANTAN UTARA", BAWAL Widya Riset Perikanan Tangkap, 2018

Publication

31

Weyl, O.L.F.. "Management recommendations for *Copadichromis chrysonotus* (Pisces: Cichlidae) in Lake Malombe, Malawi, based on per-recruit analysis", Fisheries Research, 200502

Publication

32

doczz.fr
Internet Source

<1 %

33

ejournal2.undip.ac.id
Internet Source

<1 %

34

jmb.ahvaz.iau.ir
Internet Source

<1 %

35

magazines.ulbsibiu.ro
Internet Source

<1 %

ojs.unud.ac.id

36

Internet Source

<1 %

37

sigaa.ufra.edu.br

Internet Source

<1 %

38

www.frontiersin.org

Internet Source

<1 %

39

www.scialert.net

Internet Source

<1 %

40

Ali Suman, Duranta D Kembaren, Andina Ramadhani Putri Pane, Muhammad Taufik. "Status Stok Udang Jerbung (*Penaeus merguensis*) di Perairan Bengkalis dan Sekitarnya serta Kemungkinan Pengelolaannya Secara Berkelanjutan.", Jurnal Kebijakan Perikanan Indonesia, 2020

Publication

<1 %

41

P. A. A. T. Jayawardane. "Population dynamics of *Metapenaeus dobsoni* from the western coastal waters of Sri Lanka", Fisheries Management and Ecology, 6/2003

Publication

<1 %

Exclude quotes OnExclude matches OffExclude bibliography On