


• Preliminary acceptance 8

 • **Eni Kovacs** <ek.bioflux@gmail.com>
Kepada: erick nugraha

🖨 📧 📅 Kam, 17 Jun 2021 jam 23.54 ☆

Dear Dr. Nugraha,

We would like to inform you that after the preliminary evaluation your paper is qualified for processing (reviewing). However, the manuscript does not entirely respect the AACL formatting requirements. Please find the model on our website and replace each section with your text, **leaving the footer exactly as it is, including the page numbering**.

Before proceeding to the review please consider that a processing publication fee of 250 USD is required. The average duration of the publication process is 10 weeks, but it can be reduced in exchange of a priority tax of 50 USD (<http://www.bioflux.com.ro/journal/>).

In case the manuscript is not published, the author or his/her institution is reimbursed (exceptions are made in the cases of poor feedback from authors or withdrawal/rejection due to multiple submissions).

Concerning the payment procedure, you will find all the details bellow and I would like to kindly ask you to send me the scanned receipt of the payment in order to start the publishing process. Upon reception of your payment receipt we will send you the invoice.

1st payment option:
Beneficiary:
Bioflux SRL
City: Cluj-Napoca ,
Country: Romania, European Union;


SWIFT CODE of the bank: BTRLRO22

Account USD:
RO68BTRL01302202L28614XX

Bank:
BANCA TRANSILVANIA

Important! When bank transfer is used to pay a publication fee, please choose the right option, which is "US"! (among the three options you have: Ben, Our, US). The option "Us" means that **all the transfer fees are at your charge**. Please note that in case you do not specify the "Us" option, the payment will not be valid.

2nd payment option:



The Suitability of the Fishing Ground with the Forecast Map of the Fishing Ground (PPDPI) in Terms of Chlorophyll-a in the Savu Sea, Indonesia

¹Maman Hermawan, ²I Nyoman Radiarta, ¹I Gusti B. E. K. Bayu, ¹Erick Nugraha

¹ Faculty of Fishing Technology, Jakarta Technical University of Fisheries, Pasarminggu 12520, South Jakarta, Indonesia; ² Institute for Marine Research and Observation, Negara, Jembrana 82218, Bali, Indonesia; Corresponding author: E. Nugraha, nugraha_eriq1@yahoo.co.id

Abstract. The Savu Sea in East Nusa Tenggara Province is an area that is the location of fishing activities for most fishermen in East Nusa Tenggara. Behind all the potential possessed by the Savu Sea, the knowledge of the fishing ground is still conventional even though at this time various prediction maps of fishing ground have been issued which aim to help fishermen determine their fishing area when planning a voyage. This fishing ground forecast map is based on chlorophyll-a, sea surface temperature, salinity and current data obtained from NASA which is processed by the Marine Research and Observation Center which then displays potential fishing ground that have a box view with an accuracy level of 5 square miles. After that the forecast map of the fishing area will be compared with the fishing positions used by the fishermen. Through this scientific work, it is hoped that it will be able to become material for study by fishermen and become data validation by the agency that issued the map. The data collection method is a survey method and the data analysis method uses descriptive analysis and linear regression analysis. The results of the observation resulted in a map matching with the fishing position of the fishermen of 13.6% and the highest chlorophyll-a concentration of 1.29 mg m³ and the lowest 0.07 mg m³ where during the observation the West Flores area showed the highest concentration constantly. Partially chlorophyll-a has an effect of 62.2% on the catch, with the catch is dominated by *Katsuwonus pelamis* by 52%.

Keywords: MODIS, Sea Surface Temperature, Catch Composition, Skipjack tuna (*Katsuwonus pelamis*).

Introduction. Indonesia is the largest archipelagic country in the world, has a coastline length of 81,000 km² and a sea area of about 3,100,000 km² (Kusumastanto 1996; Darsono 1999; Latif et al 2019). An important information that needs to be known in terms of fisheries resource management, is determining fish growth patterns and catch period, so fisherman catching only feasible fish category (Nugraha et al 2020b). The province of East Nusa Tenggara has a sea area of 200,000 km² with a coastline of 5,700 km² (MMAF 2016). In the fisheries and marine sector, Savu Sea which has good fishery potential (Asagabaldan 2017).

A fishing ground is an area in the waters where there are many fish that are targeted to be caught and fishing gear can be operated technically and safely (Panjaitan 2014). The *Katsuwonus pelamis* is a fast swimming fish and has a greedy eating character and includes opportunistic predators with food types ranging from small fish (*Clupeidae* and *Engraulidae*), squids, crustaceans and Zooplankton (Tuli et al 2015). *K. pelamis* live in groups pelagically in coastal areas to the open sea (Nainggolan 2012). Schooling fish should be kept as close to the boat as possible so that they can be easily caught using pole and line fishing rods (Sudirman & Mallawa 2004). The determination of fishing ground can be expected from the condition of the waters where the habitat of a species. Water conditions are usually described by oceanographic parameters (Nugraha et al 2020a).

The characteristics of the Sea Surface Temperature (SST) of the *K. pelamis* fishing ground are in the range between 29.75 to 30.25°C (Zainuddin et al 2015; Nugraha et al 2020b; Zainuddin 2011). Meanwhile, the potential area has a characteristic chlorophyll-a concentration of around 0.125 mg m³. This condition is mainly seen in May. The catch of *K. pelamis* tends to decline outside of this range, as happened in April (SST conditions are highest in the Bone-Sea Flores bay). Likewise, when the chlorophyll-a concentration in the fishing ground has increased significantly, the catch tends to decrease. This means that

there is a certain range of values for the two oceanographic factors that are very suitable for the presence and abundance of *K. pelamis*. To knowing the distribution of chlorophyll-a can be a parameter in identifying the presence of fish in an area of water (Hasyim & Zainuddin 2008; Laurs *et al* 1984).

Forecast Maps of Fishing Ground (PPDPI) is the result of research conducted by *Institute for Marine Research and Observation* (IMRO) which is an effort to facilitate the search for fishing grounds in the form of internet information services based on fishing areas (Sugiari 2015; Panjaitan 2014). PPDPI have been produced and distributed since 2000 for fishing communities in Indonesia (Tambun *et al* 2018; Farda & Jatisworo 2019).

PPDPI was compiled using remote sensing data which is also widely used to detect water conditions such as the thermal front (Cayula & Cornillon 1992; Wibawa 2011; Jatisworo & Murdimanto 2013) and the concentration of chlorophyll-a (Seki *et al* 2002) which is one of the indicators of water fertility and school of fish. PPDPI preparation was carried out using SST and water fertility data (chlorophyll-a) generated from MODIS Aqua level 3 composite 3 daily with Hierarchical Data Format (HDF) (Komang *et al* 2015).

Map of prediction of fishing ground that already exists in Indonesia there are still many weaknesses including the spatial aspect of the point of forecasting far from the coast and the unclear type of fish predicted. The lack of understanding of the type of fish predicted, the risk of fishermen carrying the wrong fishing gear so that when reaching the forecasting location, fishermen can not catch fish that are abundant (Kunarso *et al* 2019).

Chlorophyll-a is an indicator of the abundance of phytoplankton in waters that play a role in the photosynthesis process (Huot *et al* 2007; Zhang & Han 2015). Phytoplankton contributes to a large extent to determine primary productivity in waters. Organic carbon production during photosynthesis is defined as primary productivity or Net Primary Productivity (Lee *et al* 2016).

Linear regression analysis was carried out on the model of the relationship between chlorophyll-a concentration and primary productivity. The results of this equation can be applied to satellite imagery so that it can help in monitoring water quality conditions (Nuzapril *et al* 2017). This abundance of phytoplankton will result in the creation of food chain clusters from phytoplankton to large pelagic fish schools (Sato *et al* 2015; Behrenfeld *et al* 2005).




Figure 1. Map of chlorophyll-a concentration (IMRO 2020).

Remote sensing data can show the variables used in determining fish catchment areas. One of the remote sensing images used to extract SST and chlorophyll-a information is a MODIS image (Nurman 2010; Hakim *et al* 2006).

Satellite imagery is one of the results of remote sensing technology. One of the technologies used is the Moderate-resolution Imaging Spectroradiometer (MODIS) which is one of the sensors belonging to the Earth Observing System (EOS) and is carried by two vehicles produced by the National Aeronautics and Space Administration (NASA), namely Terra and Aqua (<http://modis.gsfc.nasa.gov> 2007).

The MODIS sensor is a derivative of the Advanced Very High Resolution Radiometer (AVHRR) sensor, Sea-viewing Wide Field of View Sensor (SeaWiFS) and High Resolution Imaging Spectrometer (HRS) owned by EOS which has previously orbited, so it can be used to measure parameters that are in at sea level to the atmosphere, among others, measures the sea surface temperature, chlorophyll-a concentration, water vapor content and marine phenomena, such as the occurrence of upwelling, thermal fronts and others (<http://modis.gsfc.nasa.gov> 2007).

Material And Method

Description of the study sites. This research was conducted on a pole and line fishing boat operating in the Flores Sea, East Nusa Tenggara. For satellite image data retrieval is carried out at the Southeast Research and Ocean Observation Center (SeaCORM) Jembrana, Bali. The material used is sea water as an object for measuring oceanographic elements. The methods used in data collection are literature study, interview, documentation, observation.

Data analysis method

Regression analysis is a statistical analysis that studies the relationship between two or more variables. In linear regression analysis it is assumed that the form of a linear relationship in the parameters applies. The simplest linear regression module is linear regression with one independent variable (Pangestu 2016).

Linear regression is a statistical method used to model the relationship between the dependent variable (dependent; response; Y) and one or more independent variables (independent, predictor, X). If there are only one independent variable, it is called simple linear regression, whereas if there is more than one independent variable, it is called multiple linear regression. Regression analysis has at least three uses, namely for the purpose of description of the data phenomenon or the case being studied, for control purposes, and for predictive purposes.

Simple regression is carried out in stages, namely by connecting the independent variable (chlorophyll-a) with the dependent variable (tuna catch). In the use of Microsoft Office Excel to process data, namely by using data analysis then choosing data regression analysis (Kurniawan 2008). This simple regression analysis was conducted to determine the relationship between the dependent variable Y (*K. pelamis* catches) and the independent variable X (chlorophyll-a). The regression model used is:

$$Y = a + bX$$

Where:

X = Chlorophyll-a

Y = Pole and Line catch

a = constant (intercept), the value of a intersects the Y axis
The value of Y, when X = 0

b = regression coefficient

Chlorophyll-a image analysis

Determination of the fishing ground forecast can be predicted from the parameters of sea surface temperature and chlorophyll-a. The analysis of the distribution of sea surface temperature and chlorophyll-a can use level 1b and level 2 Modis images. The *Algorithm Theoretical Basic Document Modis* (ATBD) 25 is used to determine sea surface temperature and the Morel 4 Algorithm is used to determine chlorophyll-a. Modis image level 1b and 2 using the RMSE test calculation (Chavula 2012).

From the two *Root Mean Squared Error* (RMSE) tests carried out between the field data and the Modis image data level 1b and level 2, the better and more accurate RMSE value is obtained from the smallest RMSE sum value, namely the Level 2 Modis image with

an RMSE value of chlorophyll-a of 0.090315. So in determining the best fishing ground forecast is to use Level 2 Modis Image (Febriani & Sukojo 2016).

Results and Discussion

Fishing Ground. In this study we used pole and line fishing boat (figure 2) that operates in Savu sea, Flores, East Nusa Tenggara.



Figure 2. Pole and Line fishing boat

Composition and comparison of catches with the number of settings.

1. In November 2019 the ship made one trip with a catch of *K. pelamis* 1,020 kg.
2. In December 2019 the ship made 16 trips with a catch of 28,220 kg. The fishing season generally reaches a turning point in the middle of January and in December, according to the graph, the catch will decrease as the wind season has entered.
3. In January 2020 the catch obtained was 7,990 kg. January is generally the start of the wind season, so ships rarely go to sea.
4. In February 2020 the catches were dominated by *K. pelamis* and *T. albacares*. In February, fishermen said it was generally the peak of the wind season so that the catch was not good, this lasted until the middle or end of March.
5. In March 2020 the catches were dominated by *K. pelamis* and *T. albacares*, but unlike the previous month, this month, *C. hippurus* and *E. bipinnulata* were obtained. In March 2020 the fishermen began to increase the intensity of their trips, but taking into account the strength of the wind and the availability of bait.

The composition of the catch for 5 months can be seen in table 1 below:

Table 1

Comparison of catching with the number of settings

Period / Kg	November	December	January	February	March
Number of setting (times)	2	43	14	15	13
<i>K. pelamis</i>	1,020	15,300	2,890	6,290	7,480
<i>T. albacares</i>	0	9,520	2,040	5,440	4,250
<i>C. hippurus</i>	0	1,870	1,870	0	1,360
<i>E. bipinnulata</i>	0	1,530	1,190	0	1,870
Total	1,020	28,220	7,990	11,730	14,960
Catch rate per setting	510	656.28	570.71	782	1,150

or we can be seen in the figure 3 below:




Figure 3. Total catch (November 2019 to March 2020)

Based on the graph above, there is a decrease in the catch during the 4 months fishing. The lowest catch was obtained in January with a total catch of 7,990 kg consisting of *K. pelamis* 2,890 kg, *T. albacares* 2,040 kg and others were *E. bipinnulata* and *C. hippurus*. Meanwhile, the highest catch was obtained in December is 28,220 kg, dominated by *K. pelamis* with 15,300 kg and *T. albacares* 9,520 kg. In 2016, Kupang Regency was the largest supplier of *K. pelamis* in East Nusa Tenggara (MMAF 2018)

Catch Composition. If we see in Table 2 and figure 4 below, it is shown that *K. pelamis* is the most dominant catch that lands at Tenau port and fishing port around Kupang city, East Nusa Tenggara. This is in accordance with the statement and validating data from the Fisheries Service of the Province of East Nusa Tenggara that *K. pelamis* 52% is one of the main commodities produced in Kupang, followed by 33% *T. Albacares*, 7% *C. hippurus* and 8% *E. bipinnulata*.

Table 2

Catch rated per setting

Name of Fish	Amount (kg)	Catch Percentage (%)
<i>K. pelamis</i>	32,980	52
<i>T. albacares</i>	21,250	33
<i>C. hippurus</i>	4,590	7
<i>E. bipinnulata</i>	5,100	8
Total Catching	63,920	100
Catch rate per setting	734.71	1.14

The percentage Catching composition can see in Figure 4 below:




Figure 4. Catch composition diagram

Conformity between of the Fishing ground and PPDPI. Fishing is carried out on FADs that have been installed and with the position of FADs stored on the GPS of each ship. The following is a map of the fishing grounds with the positions of the FADs. Figure 5 below shows the fishing ground for 5 months of fishing:




Figure 5. Fishing ground

Forecast Map of Fishing Ground (PPDPI)

1. In November the PPDPI that can be used is 29 to 30 November 2019. On this map the potential fishing ground are around the northwest of Timor Island with an area of 150 NM.
2. In December, 15 files were issued with an interval of 2 days. At PDPPI 11 to 12 December 2019, the recommended fishing areas are south of Savu Island, northeast of Sumba Island and north of Rote Island, as well as many other areas with an area of more than 300 NM (IMRO 2019).
3. In January the PPDPI issued which corresponds to the time of the Catching trip, only 1 file out of 7 uploaded files. On the 31 December to 01 January map, the potential fishing grounds are around the southeast of Sumba Island which is one with the western area of Savu Island with an area of more than 300 NM (IMRO 2020).
4. In February the PPDPI uploaded as many as 10 files and there are 3 files that correspond to the time of the catching trip. One suitable file is on the 26 to 27 February map with the recommended fishing area just north of Sumba Island with an area of 300 NM. Other areas do not show signs of potential fishing grounds (IMRO 2020).
5. In March there were 4 files that were used because they matched the time of the catching trip. March should be the turning point of the monsoon season that occurs in East Nusa Tenggara but each region has a different time. One of the maps is the 16 to 17 March map, the areas that are recommended for fishing activities are in the south of Sumba Island and northeast of Sumba Island with an area of more than 300 NM, there is also one point with an area of 50 NM located on Semau and Kupang islands (IMRO 2020).




Figure 6. PPDPI period in November (1), Desember (2) (IMRO 2019); Januari (3), Februari (4), Maret (5) (IMRO 2020).

Compatibility between the Fishing Ground and PDPPI. The total trips was made and matched with the availability of PDPPI, there were only 17 files that matched the time of the fishing trip. For 4 months there were 17 map files forecasts of fishing ground used with a total of 44 settings (table 3).

Table 3

Compatibility positions of fishing ground with PPDPI


No	Setting	Position		Klorofil-a (mg m ⁻³)	Information
		Latitude	Longitude		
1	1	10 ⁰ 12' 27" S	121 ⁰ 40' 16" E	0.41	Inappropriate
	2	10 ⁰ 09' 43" S	121 ⁰ 37' 40" E	0.31	Inappropriate
2	1	09 ⁰ 56' 38" S	122 ⁰ 25' 03" E	0.3	Appropriate
	2	10 ⁰ 42' 09" S	121 ⁰ 50' 28" E	0.27	Inappropriate
	3	10 ⁰ 26' 28" S	122 ⁰ 47' 58" E	0.27	Appropriate
3	1	09 ⁰ 32' 11" S	123 ⁰ 20' 42" E	0.24	Inappropriate
	2	09 ⁰ 45' 37" S	123 ⁰ 13' 20" E	0.22	Inappropriate
4	1	10 ⁰ 04' 18" S	122 ⁰ 32' 21" E	0.27	Appropriate
	2	10 ⁰ 28' 31" S	122 ⁰ 41' 16" E	0.31	Inappropriate
	3	10 ⁰ 12' 31" S	122 ⁰ 05' 10" E	0.33	Appropriate
5	1	10 ⁰ 40' 10" S	122 ⁰ 15' 30" E	0.29	Inappropriate
	2	09 ⁰ 40' 33" S	122 ⁰ 55' 15" E	0.39	Inappropriate
6	1	09 ⁰ 31' 40" S	123 ⁰ 05' 09" E	0.21	Inappropriate

	2	09° 05' 45" S	123° 00' 41" E	0.28	Inappropriate
	3	09° 47' 16" S	122° 58' 34" E	0.13	Inappropriate
7	1	09° 40' 08" S	123° 22' 31" E	0.29	Inappropriate
	2	09° 50' 28" S	122° 53' 17" E	0.30	Inappropriate
8	1	10° 38' 15" S	121° 40' 30" E	0.41	Inappropriate
	2	10° 42' 09" S	121° 50' 28" E	0.38	Inappropriate
	3	10° 49' 25" S	121° 01' 51" E	-	Inappropriate
9	1	09° 32' 11" S	123° 20' 42" E	0.28	Inappropriate
	2	09° 45' 37" S	123° 13' 20" E	0.24	Inappropriate
	3	09° 54' 26" S	123° 01' 51" E	-	Inappropriate
10	1	10° 21' 15" S	122° 35' 07" E	0.45	Inappropriate
	2	10° 10' 35" S	122° 05' 37" E	-	Inappropriate
	3	10° 03' 47" S	122° 45' 10" E	0.54	Appropriate
11	1	09° 40' 36" S	122° 05' 15" E	0.8	Inappropriate
	2	09° 50' 17" S	121° 35' 20" E	1.1	Inappropriate
12	1	09° 30' 14" S	122° 50' 33" E	-	Inappropriate
	2	09° 21' 52" S	123° 00' 16" E	-	Inappropriate
13	1	10° 40' 16" S	123° 41' 40" E	-	Inappropriate
	2	10° 39' 13" S	123° 45' 36" E	-	Inappropriate
	3	10° 40' 41" S	123° 40' 22" E	-	Inappropriate
14	1	09° 50' 20" S	122° 10' 31" E	-	Inappropriate
	2	09° 18' 50" S	122° 46' 15" E	-	Inappropriate
	3	09° 30' 16" S	123° 01' 51" E	-	Inappropriate
15	1	10° 03' 15" S	122° 45' 17" E	0.31	Inappropriate
	2	10° 03' 40" S	121° 48' 29" E	-	Inappropriate
16	1	10° 08' 36" S	123° 07' 12" E	-	Inappropriate
	2	10° 15' 53" S	122° 46' 41" E	-	Inappropriate
	3	10° 03' 09" S	122° 53' 08" E	-	Inappropriate
17	1	09° 29' 21" S	122° 30' 14" E	0.6	Inappropriate
	2	09° 50' 15" S	122° 30' 35" E	0.54	Inappropriate
	3	09° 55' 23" S	121° 20' 54" E	0.49	Appropriate


The method used to determine the suitability of fishing grounds with PDPPI issued by IMRO uses a comparison method between the position of the fishing area that has been carried out with the position of the forecast map of the appropriate fishing area on that date. After being matched from 44 setting positions, there are 6 positions that are in accordance with PDPPI, so it can be concluded by calculating the suitable position points compared to the total number of positions carried out multiplied by 100%, the level of compatibility of the fishing ground of fishermen with PDPPI which has a current reference is obtained, chlorophyll-a and sea surface temperature have a match rate of 13.6%. Not all fishing trips can be used as a reference in determining suitability, because PDPPI cannot be uploaded every day because it is limited by the availability of images obtained from the NASA website.

With a match rate of 13.6%, it can be said that fishermen using FADs have sufficient accuracy in determining fishing grounds.


Concentration and Distribution of chlorophyll-a. Chlorophyll-a concentrations can be obtained by downloading data levels 1, 2 and 3 on the NASA website. Data needs in a longer time span such as months and years can use level 3 image data with better quality than others, but level 3 images only contain one specific information depending on what is needed. In this study, the basis used is a level 3 image which only requires chlorophyll-a information with a time span of months. In Figure 7 and Table 4 the following is image data for chlorophyll-a concentration level 3:




November 2019




December 2019



January 2020



February 2020



March 2020

Figure 7. Concentration of chlorophyll-a map (www.oceancolour.gsfc.nasa.gov)

By looking at the distribution map of chlorophyll-a concentrations (table 4) that has been obtained from November to March, it shows that around the west of Flores Island, Sumba Island and also Larantuka have very thick chlorophyll-a concentrations and it is indicated that a food chain occurs in the area.

Table 4

Concentration and Distribution of chlorophyll-a

Period	November	December	January	February	March
Concentration of chlorophyll-a (mg m ⁻³)					
a. Lowest	0.07	0.08	0.08	0.08	0.08
b. Highest	1.29	1.14	1.15	1.19	1.17
Average Value (mg m ⁻³)	0.2	0.3	0.2	0.2	0.2
Distribution Center	North of Kupang city, Sabu Island, and west of Flores	West of Flores	Kupang city, Sabu Island, Rote Island, and west of Flores	Kupang city, Rote Island, and west of Flores	Kupang city, west of Flores
Compatibility Level	Appropriate	Inappropriate	Appropriate	Appropriate	Inappropriate

Conclusion

Based on the results of the study it can be concluded:

1. Simultaneously, chlorophyll-a had an effect of 62.2% on the catch. The highest concentration of chlorophyll-a recorded was 1.29 mg m⁻³ and the lowest was 0.07 mg m⁻³.
2. The fishing area used by fishermen has a level of conformity with the forecast map for fishing areas of 13.6%.
3. Not all fishing trips can be used as a reference in determining the suitability of fishing grounds with PDPPI, because not every day PDPPI can be uploaded because it is limited by the availability of images obtained from the NASA website.

Acknowledgments. The authors would like to thank to Director and staff of Institute for Marine Research and Observation (IMRO), Agency for Marine and Fisheries Research and Development, Ministry of Marine Affairs and Fisheries, Indonesia. for the support to sharing and processing of the satellite data related to the distribution of SST and Chlorophyll-a in fishing ground and to all other parties for their help and contributions.

Conflict of interest. The authors declare no conflict of interest.

References.

- Asagabaldan M. A., 2017 Law and the Role of Lilifuk Amid Government Regimes: Legal Analysis and Sustainability of Local Wisdom in Kuanheun Village, West Kupang Regency, East Nusa Tenggara. *SABDA* 12(2):136-145.
- Behrenfeld M. J., Boss E., Siegel D. A., Shea D. M., 2005 Carbon-based ocean productivity and phytoplankton physiology from space. *Global Biogeochemical Cycles* 19(1):1-14.
- Cayula J. F., Cornillon P., 1992 Edge Detection Algorithm for SST Images. *Journal of Atmospheric and Oceanic Technology*, 9(1):67-80.
- Chavula G., Sungani H., Gondwe K., 2012 Mapping Potential Fishing Grounds in Lake Malawi Using AVHRR and MODIS Satellite Imagery. *International Journal of Geosciences* 3(03):650-658.
- Darsono P., 1999 Utilization of Marine Resources and Its Implications for Fishing Communities. *Oceana*, 24(4):1-9. [in Indonesia]
- Farda N. M., Jatisworo D., 2019 Mining Space-Time Patterns on Forecast Maps of Fishing Areas in Fisheries Management Areas (FMA) 712, 713, and 573. *Scientific Magazine GLOBE* 21(2):117-128. [in Indonesia].
- Febriani E. R., Sukojo B. M., 2016 Comparative Analysis of the Use of Level 1b and Level 2 MODIS Imagery in Determining Forecasting Fishing Areas (Case Study: South Coast of Blitar). *ITS Engineering Journal*, Surabaya 5(2):439-442. [in Indonesia]

- Hakim, D. M., Wikantika K., Widiadnyana N., Napitu A. M., Darmawan S., 2006 The Identification of Fishing Ground Area with MODIS Satellite Image (Case Study: South Coast of West Java) Proceeding. ITB Engineering Science 38B(2):147-158.
- Hasyim S., Zainuddin M., 2008 Predictions of Skipjack Fishing Areas Based on Oceanographic Conditions in the Waters of Takalar Regency and Surrounding Areas. Journal of Science and Technology 8(2):158-162. [in Indonesia]
- Huot Y., Babin M., Bruyant F., Grob C., Twardowski M. S., Claustre H., 2007 Does Chlorophyll-a Provide The Best Index Of Phytoplankton Biomass For Primary Productivity Studies?. Biogeosciences Discussions, Proxies Of Biomass For Primary production 4(2):707-745.
- Jatisworo D., Murdimanto A., 2013 Identification of thermal fronts in the Makassar Strait and Banda Sea. Proceedings of the National Symposium on Geoinformation Science III. Marine Research and Observation Institute. p. 8. [in Indonesian].
- Kunarso, Zainuri M., Sugianto D. N., Marwoto J., Hariyadi, Muslim, 2019 Prediction of Demersal Fishing Ground Associated with Coral Reefs in the Coastal Jepara Regency, Central Java, Indonesia Based on Sentinel 2a Imagery. Advances in Science, Technology and Engineering Systems (ASTES) Journal 4(6):263-269.
- Kurniawan D., 2008 Linear Regression. Academia Accelerating the world's research. Research Development Core Team (2008). R: A language and environment for statistical computing. (<http://ineddeni.wordpress.com>). p. 1-14 [in Indonesia].
- Kusumastanto T., 1996 An Evaluation Of Government Policies On The Development Of Fisheries In Indonesia. AGRIS, Fisheries Economy Bulletin 2(3):27-41 [in Indonesia].
- Latif B., Noor S. M., Sumardi J., Irwansyah, 2019 Sustainable development and sea protection: Trade on fish and fishery product. ICROEST. IOP Conf. Series: Earth and Environmental Science 343(1):1-6.
- Laurs R. M., Fiedler P. C., Montgomery D. R., 1984 Albacore Tuna Catch distribution relative to environmental features observed from satellites. Deep-Sea Research 31:1085-1099.
- Lee Y. J., Matrai P. A., Friedrichs M. A. M., Saba V. S., Aumont O., Babin M., Buitenhuis E. T., Chevallier M., Mora L., Dessert D., Dunne J. P., Ellingsen I. H., Feldman D., Frouin R., Gehlen M., Gorgues T., Ilyina T., Jin M., John J. G., Lawrence J., Manizza M., Menkes C. E., Perruche C., Fouest V. L., Popova E. E., Romanou A., Samuelsen A., Schwinger J., Séférian R., Stock C. A., Tjiputra J., Tremblay L. B., Ueyoshi K., Vichi M., Yool A., Zhang J., 2016 Net primary productivity estimates and environmental variables in the Arctic Ocean: An assessment of coupled physical-biogeochemical models. Journal of Geophysical Research: Oceans 121(12):8635-8669.
- Nainggolan C., 2012 Metode Penangkapan Ikan. Open University, Jakarta p. 1-61. [in Indonesia]
- Nugraha E., Gunawan R., Danapraja S., Yusrizal, Kusdinar A., Waluyo A. S., Hutajulu J., Prayitno H., Halim S., Sutisna D. H., 2020b The Sea Surface Temperature Effect On The Length And Size Of Skipjack Tuna (*Katsuwonus Pelamis*) Catches in the Banda Sea, Indonesia. AACL Bioflux 13(1):1-18.
- Nugraha E., Yudho G. S., Jaenudin A., Yusrizal, Kusmedy B., Kusnidar A., Husen E. S., 2020a Relationship Between Length And Weight Of Skipjack Tuna (*Katsuwonus Pelamis*) Purse Seine Catching in the Maluku Sea, Indonesia. AACL Bioflux 13(1):330-345.
- Nurman A., 2010 Utilization of Fashionable Data to Detect Fishing Grounds on the East and West Coasts of North Sumatra. Medan State University. Geography Journal 2(2):17-30. [in Indonesia]
- Nuzapril M., Susilo S. B., Panjaitan J. P., 2017 Relationship Between Chlorophyll-a Concentration With Primary Productivity Rate Using Landsat 8 Imagery. Journal of Fisheries and Marine Technology 8(1):105-114.
- Pangestu S., 2016 Simple Linear Regression. Applied Linear Model. Digital Library of a Open University of Indonesia p. 398 [in Indonesia]
- Panjaitan H., 2014 Utilization of Fishing Area Forecast Map. Fisheries Writing Book. Field. Access : ocean-wide-ki.blogspot.com [in Indonesia]

- Sato K., Matsuno K., Arima D., Abe Y., Yamaguchi A., 2015 Spatial and Temporal Changes in Zooplankton Abundance, Biovolume, and Size Spectra in the Neighboring Waters of Japan: Analyses Using An Optical Plankton Counter. *Zoological Studies. A Springer Open Journal* 54(18):1-15.
- Seki M. P., Polovina J. J., Kobayashi D. R., Bidigare R. R., Mitchum G. T., 2002 An oceanographic characterization of swordfish (*Xiphias gladius*) longline fishing grounds in the springtime subtropical North Pacific. *Fisheries Oceanography* 11(5):251-266.
- Sudirman, Mallawa A., 2012 *Fishing Techniques*. Rineka Cipta: Jakarta p. 168. [in Indonesia]
- Sugiari, Anshori M. A., Kusumawardani M., 2015 Coordinate Information System for Estimated Tuna Fishing grounds for Fishermen. *Journal of JARTEL State Polytechnic of Malang* 1(1):74-84. [in Indonesia]
- Suniada K. I., Susilo E., Hastuti A. W., 2015 Validation of Forecast Map of Fishing Grounds (PPDPI) in Java Sea Waters (FMA-RI 712). *Proceedings of the National Marine and Fisheries Science and Technology Forum* p. 87-96 [in Indonesia].
- Tambun R., Simbolon D., Wahyu R., Supartono, 2018 Potential Zones of Fishing Ground Based on Season at FMA RI 718. *Journal of Tropical Marine Science and Technology* 10(3):757-768 [in Indonesia]
- Tuli M., Menofatria B., Adrianto L., 2015 Analysis of Skipjack fish (*Katsuwonus pelamis*) resources in the waters of Pohuwato Regency, Gorontalo Province. *Marine Fisheries Journal* 6(2):109-117. [in Indonesia]
- Wibawa T. A., 2011 Utilization of oceanographic satellite data for prediction of potential fishing grounds for Bigeye Tuna (*T. obesus*) in the South Indian Ocean, Java and Bali. *SEGARA Journal* 7(1):29-41. [in Indonesia]
- Zainuddin M., 2011, Skipjack Tuna In Relation To Sea Surface Temperature And Chlorophyll-a Concentration Of Bone Bay Using Remotely Sensed Satellite Data. *Journal of Tropical Marine Science and Technology* 3(1):82-90.
- Zainuddin M., Safruddin, Farhum S. A., Nelwan A., Selamat M. B., Hidayat S., Sudirman, 2015 Characteristics of Potential Areas for Catching Skipjack Tuna in the Bone-Sea Bay of Flores based on Satellite Data on Sea Surface Temperature and Chlorophyll-a in the period January-June 2014 Makassar. *Journal of Science and Technology Utilization of fisheries resources. Hasanuddin University* 2(3):228-237. [in Indonesia]
- Zhang C., Han M., 2015 Mapping chlorophyll-a concentration in Laizhou Bay using Landsat 8 OLI data. e-proceedings of the 36th IAHR World Congress, The Hague, The Netherland p. 12-17.
- *** Institute for Marine Research and Observation (IMRO), 2019 Forecast Map of Fishing Ground. Bali. <http://bpol.litbang.kkp.go.id/> [in Indonesia]
- *** Institute for Marine Research and Observation (IMRO), 2020 Forecast Map of Fishing Ground. Bali. <http://bpol.litbang.kkp.go.id/> [in Indonesia]
- *** Ministry of Marine Affairs and Fisheries (MMAF), 2016 East Nusa Tenggara Provincial Annual Report 2016. East Nusa Tenggara Provincial Marine and Fisheries Service. [in Indonesia]
- *** Ministry of Marine Affairs and Fisheries (MMAF), 2018 East Nusa Tenggara Provincial Annual Report 2018. East Nusa Tenggara Provincial Marine and Fisheries Service. [in Indonesia]
- *** (<http://modis.gsfc.nasa.gov> 2007)
- *** (www.oceancolour.gsfc.nasa.gov)

Received: xxxxxxxx Accepted: xxxxxxxxxxxx Published online: xxxxxxxxx

Authors:

Maman Hermawan, Faculty of Fishing Technology, Jakarta Technical University of Fisheries, AUP Street Pasarminggu 12520, South Jakarta, Indonesia. e-mail: mhermawan60@gmail.com

I Nyoman Radiarta, Institute for Marine Research and Observation, Agency for Marine and Fisheries Research and Development Baru Street, Perancak, Negara, Jembrana 82251 Bali, Indonesia e-mail: radiarta@kkp.go.id
I Gusti Bagus Ega Krisna Bayu, Faculty of Fishing Technology, Jakarta Technical University of Fisheries, AUP Street Pasarminggu 12520, South Jakarta, Indonesia. e-mail: egabayu91@gmail.com
Erick Nugraha, Faculty of Fishing Technology, Jakarta Technical University of Fisheries, AUP Street Pasarminggu 12520, South Jakarta, Indonesia. e-mail: nugraha_eriq1@yahoo.co.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Hermawan M., Radiarta I. N., Bayu I. G. B. E. K., Nugraha E., 2021 The Suitability of the Fishing Ground with the *Forecast Map of the Fishing Ground (PPDPI)* in Terms of Chlorophyll-a in the Savu Sea, Indonesia. AACL Bioflux xxxxxxxxxx

← Kembali ↶ ↷ →

📁 Arsipkan 📁 Pindahkan 🗑️ Hapus 🛡️ Spam ⋮

🔍 ▲ ▼ ✕

• Preliminary acceptance 8

Yahoo/Terkiirim ☆

E • **Eni Kovacs** Dear Dr. Nugraha, We would like to inform you that after the preliminary evaluation your paper is qualified for processing (review 📧 Kam, 17 Jun 2021 jam 23.54 ☆

EN • **erick nugraha** <nugraha_eriq1@yahoo.co.id> Kepada: Eni Kovacs 📧 📧 Jum, 18 Jun 2021 jam 20.32 ☆

Dear Dr. Eni,
i have paid for this manuscript. (300 USD)

Thank You.


> Tampilkan pesan asli



WhatsAppjpeg

122.8kB

↶ ↷ → ⋮



~~The suitability of the fishing ground with the forecast map of the fishing ground (PPDPI) in terms of chlorophyll-a in the Savu Sea, Indonesia~~ Comparison between fisherman fishing ground and the forecast map of fishing ground (PPDPI) based on chlorophyll-a in the Savu Sea, Indonesia

¹Maman Hermawan, ²I Nyoman Radiarta, ¹I Gusti B. E. K. Bayu, ¹Erick Nugraha

¹ Faculty of Fishing Technology, Jakarta Technical University of Fisheries, Pasarminggu, South Jakarta, Indonesia; ² Institute for Marine Research and Observation, Negara, Jembrana, Bali, Indonesia. Corresponding author: E. Nugraha, nugraha_eriq1@yahoo.co.id

Abstract. The Savu Sea in East Nusa Tenggara Province is the location of fishing activities for most fishermen in East Nusa Tenggara. Despite its potential, the knowledge of the fishing ground is still conventional. Currently, predictive methods for trips planning are available, such as the fishing ground forecast maps based on the chlorophyll-a, sea surface temperature and salinity. The related data are provided by NASA, being processed by the Marine Research and Observation Center to display the potential fishing ground as a box view with an accuracy level of 5 square miles. Eventually, the forecast map of the fishing area will be compared with the fishing locations known by the fishermen, for validation. The data collection used a survey method and the data analysis used both a descriptive analysis and the linear regression. The forecast maps matched with the observed fishing locations at only 13.6%. The highest recorded chlorophyll-a concentration was of 1.29 mg m^{-3} (constantly observed in the West Flores) and the lowest was of 0.07 mg m^{-3} . The chlorophyll-a has an effect on 62.2% of the catch and 52% of the catch is composed of *Katsuwonus pelamis*.

Key Words: MODIS, sea surface temperature, catch composition, *Katsuwonus pelamis*.

Introduction. Indonesia is the largest archipelagic country in the world, having a coastline length of 81,000 km² and a sea area of about 3,100,000 km² (Kusumastanto 1996; Darsono 1999; Latif et al 2019). Managing the fisheries resource requires to determine the fish growth patterns and the catch period, in order to avoid over-exploitation (Nugraha et al 2020b). The province of East Nusa Tenggara has a sea area of 200,000 km² with a coastline of 5,700 km² (MMAF 2016). Savu Sea has a good potential for the fisheries and the whole marine sector (Asagabaldan 2017). ~~A fishing ground is an area in the waters with an abundance of fish that are targeted to be caught and where the fishing gear can be operated technically and safely (Panjaitan 2014).~~ Fishing area is an area where fish are caught in maximum amount and fishing gear can be operated economical. Assessment and mapping of fishing areas are needed in fishing effort (Nusantara et al 2014). The *Katsuwonus pelamis* is a fast swimming fish and has the greedy character of an opportunistic predator, with food types ranging from small fish (Clupeidae and Engraulidae), squids, crustaceans and zooplankton (Tuli et al 2015). *K. pelamis* is a pelagic fish living in groups, from the coastal areas to the open sea (Nainggolan 2012). Schooling fish should be kept as close to the boat as possible so that they can be easily caught using pole and line fishing rods (Sudirman & Mallawa 2004, 2012). The fishing grounds are determined by the waters' condition in the habitat of

Commented [A1]: Please reformulate to be clear: the idea is to confirm the chlorophyll maps-predicted fishing grounds via field observation and fishermen consultation.

Commented [WU2]: 2012 in the references.

Commented [A3R2]: Corrected thank you

a species. Water conditions are usually described by oceanographic parameters (Nugraha et al 2020a).


The Sea Surface Temperature (SST) of the *K. pelamis* fishing ground ranges from 29.75 to 30.25°C (Zainuddin et al 2015; Nugraha et al 2020b; Zainuddin 2011). Meanwhile, the potential area has a characteristic chlorophyll-a concentration of around 0.125 mg m⁻³. This condition is mainly seen in May. Outside of this range, the catch of *K. pelamis* tends to decline, as it happens in April (when the SST values in the Bone-Sea Flores bay are the highest). Likewise, when the chlorophyll-a concentration in the fishing ground significantly increase and the catch tends to decrease. This means that there is a certain range of values for the two oceanographic factors that are very suitable for the presence and abundance of *K. pelamis*. The distribution of chlorophyll-a can be a parameter used for identifying the presence of fish in a water area (Hasyim & Zainuddin 2008; Laurs et al 1984).

The Forecast Maps of Fishing Ground (PPDPI) is the result of a research conducted by the Institute for Marine Research and Observation (IMRO) in order to identify fishing grounds, based on an internet service of information on the fishing areas characteristics (Sugiari 2015). PPDPI have been produced for and distributed to the fishing communities of Indonesia, since 2000 (Tambun et al 2018; Farda & Jatisworo 2019). PDPPI was compiled using remote sensing data which is also widely used to detect water conditions such as the thermal front (Cayula & Cornillon 1992; Wibawa 2011; Jatisworo & Murdimanto 2013) and the concentration of chlorophyll-a (Seki et al 2002) which is one of the indicators of water fertility and school of fish. PDPPI preparation was carried out using SST and water fertility data (chlorophyll-a) generated from MODIS Aqua level 3 daily composites, in the Hierarchical Data Format (HDF) (Komang Suniada et al 2015)

The predictive maps of fishing grounds that already exists in Indonesia still have many weaknesses, including the spatial aspect of the points of forecasting located far from the coast and the uncertainty of the predicted type of fish. Thus, if inappropriate gear is selected for the fishing trip, the catch productivity is negatively affected, even in presence of abundant fish populations (Kunarso et al 2019).

Chlorophyll-a is an indicator of the abundance of phytoplankton in waters, playing a role in the photosynthesis process (Huot et al 2007; Zhang & Han 2015). Phytoplankton contributes to a large extent to determine primary productivity in waters. Organic carbon production during photosynthesis is defined as primary productivity or Net Primary Productivity (Lee et al 2016).

Linear regression analysis was carried out on the model of the relationship between chlorophyll-a concentration and primary productivity. The results of this equation can be applied to the satellite imagery so that it can help to monitor the water quality conditions (Nuzapril et al 2017). This abundance of phytoplankton will result in the creation of food chain clusters from phytoplankton to large pelagic fish schools (Sato et al 2015; Behrenfeld et al 2005).



Commented [Unknown A4]: In the whole manuscript, the authors have to replace "mg m³" by "mg m⁻³"

Commented [A5R4]: Thank You.

Commented [WU6]: Not listed in the references

Commented [A7R6]: Corrected : full named is Iwan Komang Suniada. Thank you.

Figure 1. Map of chlorophyll-a concentration (IMRO 2020).

Remote sensing data can show the variables used in determining fish catchment areas. Certain remote sensing images used to extract SST and chlorophyll-a information are provided by the MODIS program (Nurman 2010; Hakim et al 2006).

Satellite imagery is one of the applications of the remote sensing technology. The Moderate-resolution Imaging Spectroradiometer (MODIS) technology uses sensors belonging to the Earth Observing System (EOS), carried by two satellites produced by the National Aeronautics and Space Administration (NASA), namely Terra and Aqua (<http://modis.gsfc.nasa.gov> 2007).

The MODIS sensor is a derivative of the Advanced Very High Resolution Radiometer (AVHRR) sensor, Sea-viewing Wide Field-of-View Sensor (SeaWiFS) and High Resolution Imaging Spectrometer (HRS) owned by EOS, which has previously orbited, so it can be used to measure sea and atmosphere-level parameters (such as the sea surface temperature, chlorophyll-a concentration and water vapor content) and marine phenomena (such as the occurrence of upwelling, thermal fronts and others) (<http://modis.gsfc.nasa.gov> 2007).

Material and Method

Description of the study sites. This research was conducted on a pole and line fishing boat operating in the Flores Sea, East Nusa Tenggara. The satellite image data retrieval was carried out at the Southeast Research and Ocean Observation Center (SeaCORM) Jembrana, Bali. The material used consisted of sea water samples for measuring oceanographic elements. The methods used in data collection are literature study, interview, documentation and observation.

Data analysis method. Regression analysis is a statistical analysis that studies the relationship between two or more variables. In the linear regression analysis it is assumed that the this relationship is a linear combination, the simplest case consisting of one independent variable (Pangestu 2016).

Linear regression is a statistical method used to model the relationship between the dependent variable (dependent; response; Y) and one or more independent variables (independent, predictor, X). If there is only one independent variable, it is called simple linear regression, whereas if there is more than one independent variable, it is called multiple linear regression. Regression analysis has at least three use cases, namely for data description, for control and for predictive purposes.

Simple regression is carried out in stages, namely by connecting the independent variable (chlorophyll-a) with the dependent variable (tuna catch, using the data regression analysis function of Microsoft Office Excel (Kurniawan 2008). This simple regression analysis was conducted to determine the relationship between the dependent variable Y (*K. pelamis* catches) and the independent variable X (chlorophyll-a). The regression model used is:

$$Y = a + bX$$

Where:

X - chlorophyll-a;

Y - pole and line catch;

a - the constant (intercept) is the value of Y, when X = 0 which equals the value of a where it intersects the Y axis;

b - regression coefficient.

Chlorophyll-a image analysis. The fishing ground forecast can be predicted from the parameters of SST and chlorophyll-a. The analysis of the distribution of SST and chlorophyll-a can use level 1b and level 2 Modis images. The Algorithm Theoretical Basic Document Modis (ATBD) 25 is used to determine SST and the Morel 4 Algorithm is used

to determine the chlorophyll-a. The Modis image levels 1b and 2 use the RMSE test calculation (Chavula 2012).

From the two Root Mean Squared Error (RMSE) tests carried out between the field data and the Modis image data levels 1b and level 2, more accurate values were obtained from the smallest RMSE sum value, namely the Level 2 Modis image, with an RMSE value of chlorophyll-a of 0.090315. So the best fishing ground forecast can be determined through Level 2 Modis Images (Febriani & Sukojo 2016).

Results and Discussion

Fishing ground. In this study we used a pole and line fishing boat (Figure 2) that operates in Savu sea, Flores, East Nusa Tenggara.



Figure 2. Pole and line fishing boat.

Composition and comparison of catches with the number of settings.

1. In November 2019 the ship made one trip with a catch of *K. pelamis* of 1,020 kg.
2. In December 2019 the ship made 16 trips with a catch of 28,220 kg. The fishing season generally reaches a turning point in the middle of January and in December, according to the graph, the catch will decrease as the wind season arrives.
3. In January 2020 the catch obtained was of 7,990 kg. January is generally the start of the wind season, so ships rarely go to sea.
4. In February 2020 the catches were dominated by *K. pelamis* and *T. albacares*. According to the fishermen, February was generally the peak of the wind season, so that the catch was not good, until the middle or end of March.
5. In March 2020 the catches were dominated by *K. pelamis* and *T. albacares*, but unlike in the previous month, *C. hippurus* and *E. bipinnulata* were also obtained. In March 2020 the fishermen began to increase the intensity of their trips, but taking into account the strength of the wind and the availability of bait.

The composition of the catch for 5 months can be seen in Table 1 below:

Table 1

Comparison of catch volumes with the number of settings

Volumes / Period	November	December	January	February	March
Number of setting (times)	2	43	14	15	13
<i>K. pelamis</i> (kg)	1,020	15,300	2,890	6,290	7,480
<i>T. albacares</i> (kg)	0	9,520	2,040	5,440	4,250
<i>C. hippurus</i> (kg)	0	1,870	1,870	0	1,360
<i>E. bipinnulata</i> (kg)	0	1,530	1,190	0	1,870
Total (kg)	1,020	28,220	7,990	11,730	14,960
Catch rate per setting	510	656.28	570.71	782	1,150

(kg)

Catch frequencies can also be visualized in Figure 3, below:




Figure 3. Total catch (November 2019 to March 2020).

Based on the graph above, there is a decrease in the catch volumes during the 4 months of fishing. The lowest catch was obtained in January with a total of 7,990 kg, consisting of: *K. pelamis* (2,890 kg), *T. albacares* (2,040 kg) and others, such as *E. bipinnulata* and *C. hippurus*. Meanwhile, the highest catch was obtained in December is 28,220 kg, dominated by *K. pelamis* with 15,300 kg and *T. albacares* with 9,520 kg. In 2016, Kupang Regency was the largest supplier of *K. pelamis* in East Nusa Tenggara (MMAF 2018).

Catch composition. If we see in Table 2 and figure 4 below, it is shown that *K. pelamis* is the most dominant catch that lands at Tenau port and at the fishing port around Kupang city, East Nusa Tenggara. This is in accordance with the data from the Fisheries Service of the Province of East Nusa Tenggara, suggesting that *K. pelamis* is one of the main commodities produced in Kupang, with 52%, followed by *T. albacares*, with 33%, *C. hippurus*, with 7% and *E. bipinnulata*, with 8%.

Table 2

Catch rated per setting

Name of fish	Amount (kg)	Catch percentage (%)
<i>K. pelamis</i>	32,980	52
<i>T. albacares</i>	21,250	33
<i>C. hippurus</i>	4,590	7
<i>E. bipinnulata</i>	5,100	8
Total catching	63,920	100
Catch rate per setting	734.71	1.14

The catch compositions (in %) can be seen in Figure 4, below.

Commented [WU8]: Please reformulate, using scientific English.

Commented [A9R8]: Corrected. Thank you




Figure 4. Catch composition diagram.

Conformity between of the fishing ground and PPDPI. Fishing is carried out on FADs that have been installed, with the position of the FADs stored on the GPS of each ship. The following is a map of the fishing grounds with the positions of the FADs. Figure 5 below shows the fishing ground for 5 months of fishing:




Figure 5. Fishing ground.

Forecast map of fishing ground (PPDPI).

1. In November the PPDPI that can be used correspond to 29 to 30 November 2019. On this map, the potential fishing ground are around the northwest of Timor Island with an area of 150 NM.
2. In December, 15 files were issued with an interval of 2 days. At PDPPI 11 to 12 December 2019, the recommended fishing areas are south of Savu Island, northeast of Sumba Island and north of Rote Island, as well as many other areas with an area of more than 300 NM. In the PDPPI 11 to 12 December 2019, the fishing areas that recommended are in the south of Savu Island, northeast of Sumba Island and north of Rote Island, as well as many other areas with an area of more than 300 NM (IMRO 2019).
3. In January, only 1 file out of 7 uploaded PPDPI files issued corresponded to the time of the fishing trip. On the 31 December to 01 January map, the potential fishing grounds were located around the southeast of Sumba Island, west of Savu Island, with an area of more than 300 NM (IMRO 2020).

Commented [A10]: ? area unit, please.

Commented [A11R10]: Ya, NM or nautical mile is a unit of length used in air, marine, and space navigation, and for the definition of territorial waters.

Commented [A12]: Please reformulate

Commented [A13]: Please correct the unit

4. In February, as many as 10 PDPPI files were uploaded and 3 files corresponded to the time of the fishing trip. One of the suitable map files was issued on the 26 to 27 February recommending fishing zones located north of Sumba Island with an area of 300 NM. Other areas do not seem to be potential fishing grounds (IMRO 2020).
5. In March there were 4 files matching, at the time of the fishing trip. March should be the turning point of the monsoon season that occurs in East Nusa Tenggara, but each region has a different time. One of the maps is the 16 to 17 March map, the areas that are recommended for fishing activities are in the south of Sumba Island and northeast of Sumba Island, with an area of more than 300 NM. There was also one point with an area of 50 NM located on Semau and Kupang islands (IMRO 2020).

Commented [A14]: Please correct the surface unit




Figure 6. PDPPI period in November (1), December (2) (IMRO 2019); January (3), February (4), March (5) (IMRO 2020).

Compatibility between the fishing ground and PDPPI. Of all trips with available PDPPI, there were only 17 files that matched, at the time of the fishing trip. For the 4 months period, 17 maps of fishing ground forecasts were used, with a total of 44 settings (Table 3).

The method used to determine the suitability of fishing grounds with PDPPI, issued by IMRO, uses a comparison method between the position of the actual fishing area and the status of the same location on the forecast map, on that date. From 44 setting positions, 6 positions were in accordance with the PDPPI, which corresponds to a match rate of 13.6%, in terms of chlorophyll-a and SST. When the daily PDPPI images from the NASA website are not available, the fishing trips cannot be used as a reference in determining their suitability.

With a match rate of 13.6%, it can be said that fishermen using FADs have sufficient accuracy in determining fishing grounds

Table 3

Compatibility of the fishing ground with the PPDPI

No	Setting	Position		Chlorophyll-a (mg m ⁻³)	Information
		Latitude	Longitude		
1	1	10° 12' 27" S	121° 40' 16" E	0.41	Inappropriate
	2	10° 09' 43" S	121° 37' 40" E	0.31	Inappropriate
2	1	09° 56' 38" S	122° 25' 03" E	0.3	Appropriate
	2	10° 42' 09" S	121° 50' 28" E	0.27	Inappropriate
3	3	10° 26' 28" S	122° 47' 58" E	0.27	Appropriate
	1	09° 32' 11" S	123° 20' 42" E	0.24	Inappropriate
4	2	09° 45' 37" S	123° 13' 20" E	0.22	Inappropriate
	1	10° 04' 18" S	122° 32' 21" E	0.27	Appropriate
5	2	10° 28' 31" S	122° 41' 16" E	0.31	Inappropriate
	3	10° 12' 31" S	122° 05' 10" E	0.33	Appropriate
6	1	10° 40' 10" S	122° 15' 30" E	0.29	Inappropriate
	2	09° 40' 33" S	122° 55' 15" E	0.39	Inappropriate
7	1	09° 31' 40" S	123° 05' 09" E	0.21	Inappropriate
	2	09° 05' 45" S	123° 00' 41" E	0.28	Inappropriate
8	3	09° 47' 16" S	122° 58' 34" E	0.13	Inappropriate
	1	09° 40' 08" S	123° 22' 31" E	0.29	Inappropriate
9	2	09° 50' 28" S	122° 53' 17" E	0.30	Inappropriate
	1	10° 38' 15" S	121° 40' 30" E	0.41	Inappropriate
10	2	10° 42' 09" S	121° 50' 28" E	0.38	Inappropriate
	3	10° 49' 25" S	121° 01' 51" E	-	Inappropriate
11	1	09° 32' 11" S	123° 20' 42" E	0.28	Inappropriate
	2	09° 45' 37" S	123° 13' 20" E	0.24	Inappropriate
12	3	09° 54' 26" S	123° 01' 51" E	-	Inappropriate
	1	10° 21' 15" S	122° 35' 07" E	0.45	Inappropriate
13	2	10° 10' 35" S	122° 05' 37" E	-	Inappropriate
	3	10° 03' 47" S	122° 45' 10" E	0.54	Appropriate
14	1	09° 40' 36" S	122° 05' 15" E	0.8	Inappropriate
	2	09° 50' 17" S	121° 35' 20" E	1.1	Inappropriate
15	1	09° 30' 14" S	122° 50' 33" E	-	Inappropriate
	2	09° 21' 52" S	123° 00' 16" E	-	Inappropriate
16	1	10° 40' 16" S	123° 41' 40" E	-	Inappropriate
	2	10° 39' 13" S	123° 45' 36" E	-	Inappropriate
17	3	10° 40' 41" S	123° 40' 22" E	-	Inappropriate
	1	09° 50' 20" S	122° 10' 31" E	-	Inappropriate
18	2	09° 18' 50" S	122° 46' 15" E	-	Inappropriate
	3	09° 30' 16" S	123° 01' 51" E	-	Inappropriate
19	1	10° 03' 15" S	122° 45' 17" E	0.31	Inappropriate
	2	10° 03' 40" S	121° 48' 29" E	-	Inappropriate
20	1	10° 08' 36" S	123° 07' 12" E	-	Inappropriate
	2	10° 15' 53" S	122° 46' 41" E	-	Inappropriate
21	3	10° 03' 09" S	122° 53' 08" E	-	Inappropriate
	1	09° 29' 21" S	122° 30' 14" E	0.6	Inappropriate
22	2	09° 50' 15" S	122° 30' 35" E	0.54	Inappropriate
	3	09° 55' 23" S	121° 20' 54" E	0.49	Appropriate

Concentration and distribution of chlorophyll-a. Chlorophyll-a concentrations can be obtained by downloading data levels 1, 2 and 3 on the NASA website. Data needs in a longer time span such as months and years can use level 3 image data with better quality than others, but level 3 images only contain one specific information depending on what is needed. In this study, the basis used is a level 3 image which only requires chlorophyll-a information with a time span of months. In Figure 7 and Table 4 the following is image data for chlorophyll-a concentration level 3.

Chlorophyll-a concentrations can be obtained by downloading data levels 1, 2 and 3 from the NASA website. Data needs in a longer time span such as months and years can use level 3 image data with better quality than the others, but only containing one specific information, depending on what is needed. In this study, the basis used is a level 3 image which only requires chlorophyll-a information with a time span of months. Figure 7 images and Table 4 data correspond to level 3 chlorophyll-a concentration.




Figure 7. Concentration of chlorophyll-a map (www.oceancolour.gsfc.nasa.gov).

The distribution map of chlorophyll-a concentrations (Table 4) from November to March, shows elevated values around the west of Flores Island, Sumba Island and also Larantuka, suggesting that food chain occur in the area.

Table 4
Concentration and distribution of chlorophyll-a

<i>Period</i>	<i>November</i>	<i>December</i>	<i>January</i>	<i>February</i>	<i>March</i>
Concentration of chlorophyll-a (mg m ⁻³)					
a. Lowest	0.07	0.08	0.08	0.08	0.08
b. Highest	1.29	1.14	1.15	1.19	1.17
Average Value (mg m ⁻³)	0.2	0.3	0.2	0.2	0.2
Distribution center	North of Kupang city, Sabu Island, and west of Flores	West of Flores	Kupang city, Sabu Island, Rote Island, and west of Flores	Kupang city, Rote Island, and west of Flores	Kupang city, west of Flores
Compatibility level	Appropriate	Inappropriate	Appropriate	Appropriate	Inappropriate

Conclusions. Based on the results of the study it can be concluded: simultaneously, chlorophyll-a had an effect of 62.2% on the catch. The highest concentration of chlorophyll-a recorded was 1.29 mg m⁻³ and the lowest was 0.07 mg m⁻³; the fishing area used by fishermen has a level of conformity with the forecast map for fishing areas of 13.6%; Not all fishing trips can be used as a reference in determining the suitability of fishing grounds with PDPPI, because not every day PPDPI can be uploaded because it is limited by the availability of images obtained from the NASA website.

Acknowledgments. The authors would like to thank to Director and staff of Institute for Marine Research and Observation (IMRO), Agency for Marine and Fisheries Research and Development, Ministry of Marine Affairs and Fisheries, Indonesia. for the support to sharing and processing of the satellite data related to the distribution of SST and Chlorophyll-a in fishing ground and to all other parties for their help and contributions.

Conflict of interest. The authors declare no conflict of interest.

References

- Asagabaldan M. A., 2017 Law and the role of Lilifuk Amid Government Regimes: Legal analysis and sustainability of local wisdom in Kuanheun Village, West Kupang Regency, East Nusa Tenggara. *SABDA* 12(2):136-145.
- Behrenfeld M. J., Boss E., Siegel D. A., Shea D. M., 2005 Carbon-based ocean productivity and phytoplankton physiology from space. *Global Biogeochemical Cycles* 19(1):1-14.
- Cayula J. F., Cornillon P., 1992 Edge detection algorithm for SST Images. *Journal of Atmospheric and Oceanic Technology* 9(1):67-80.
- Chavula G., Sungani H., Gondwe K., 2012 Mapping potential fishing grounds in Lake Malawi using AVHRR and MODIS satellite imagery. *International Journal of Geosciences* 3(03):650-658.
- Darsono P., 1999 [Utilization of marine resources and its implications for fishing communities]. *Oceana* 24(4):1-9. [In Indonesian].

- Farda N. M., Jatisworo D., 2019 [Mining space-time patterns on forecast maps of fishing areas in fisheries management areas (FMA) 712, 713, and 573]. Scientific Magazine GLOBE 21(2):117-128. [In Indonesian].
- Febriani E. R., Sukojo B. M., 2016 [Comparative analysis of the use of level 1b and level 2 MODIS imagery in determining forecasting fishing areas (Case Study: South Coast of Blitar)]. ITS Engineering Journal, Surabaya 5(2):439-442. [In Indonesian].
- Hakim D. M., Wikantika K., Widiadnyana N., Napitu A. M., Darmawan S., 2006 The identification of fishing ground area with MODIS satellite image (Case Study: South Coast of West Java) Proceeding. ITB Engineering Science 38B(2):147-158.
- Hasyim S., Zainuddin M., 2008 Predictions of skipjack fishing areas based on oceanographic conditions in the waters of Takalar regency and surrounding areas. Journal of Science and Technology 8(2):158-162. [In Indonesian].
- Huot Y., Babin M., Bruyant F., Grob C., Twardowski M. S., Claustre H., 2007 Does chlorophyll-a provide the best index of phytoplankton biomass for primary productivity studies? Biogeosciences Discussions, Proxies of Biomass For Primary production 4(2):707-745.
- Jatisworo D., Murdimanto A., 2013 [Identification of thermal fronts in the Makassar Strait and Banda Sea]. Proceedings of the National Symposium on Geoinformation Science III. Marine Research and Observation Institute, 8 p. [In Indonesian].
- Kunarso, Zainuri M., Sugianto D. N., Marwoto J., Hariyadi, Muslim, 2019 Prediction of demersal fishing ground associated with coral reefs in the Coastal Jepara Regency, Central Java, Indonesia based on Sentinel 2a imagery. Advances in Science, Technology and Engineering Systems (ASTES) Journal 4(6):263-269.
- Kurniawan D., 2008 [Linear regression]. Academia accelerating the world's research. Research Development Core Team R: A language and environment for statistical computing, 14 p. [In Indonesian].
- Kusumastanto T., 1996 [An evaluation of government policies on the development of fisheries in Indonesia]. AGRIS, Fisheries Economy Bulletin 2(3):27-41. [In Indonesian].
- Latif B., Noor S. M., Sumardi J., Irwansyah, 2019 Sustainable development and sea protection: Trade on fish and fishery product. ICROEST. IOP Conference Series: Earth and Environmental Science 343(1):1-6.
- Laurs R. M., Fiedler P. C., Montgomery D. R., 1984 Albacore tuna catch distribution relative to environmental features observed from satellites. Deep-Sea Research 31:1085-1099.
- Lee Y. J., Matrai P. A., Friedrichs M. A. M., Saba V. S., Aumont O., Babin M., Buitenhuis E. T., Chevallier M., Mora L., Dessert D., Dunne J. P., Ellingsen I. H., Feldman D., Frouin R., Gehlen M., Gorgues T., Ilyina T., Jin M., John J. G., Lawrence J., Manizza M., Menkes C. E., Perruche C., Fouest V. L., Popova E. E., Romanou A., Samuelsen A., Schwinger J., Séférian R., Stock C. A., Tjiputra J., Tremblay L. B., Ueyoshi K., Vichi M., Yool A., Zhang J., 2016 Net primary productivity estimates and environmental variables in the Arctic Ocean: An assessment of coupled physical-biogeochemical models. Journal of Geophysical Research: Oceans 121(12):8635-8669.
- Nainggolan C., 2012 [Fishing methods]. Open University, Jakarta, 61 p. [In Indonesian].
- Nugraha E., Yudho G. S., Jaenudin A., Yusrizal, Kusmedy B., Kusnidar A., Husen E. S., 2020a Relationship between length and weight of skipjack tuna (*Katsuwonus pelamis*) purse seine catching in the Maluku Sea, Indonesia. AACL Bioflux 13(1):330-345.
- Nugraha E., Gunawan R., Danapraja S., Yusrizal, Kusdinar A., Waluyo A. S., Hutajulu J., Prayitno H., Halim S., Sutisna D. H., 2020b The sea surface temperature effect on the length and size of skipjack tuna (*Katsuwonus pelamis*) catches in the Banda Sea, Indonesia. AACL Bioflux 13(1):1-18.
- Nurman A., 2010 [Utilization of fashionable data to detect fishing grounds on the east and west coasts of North Sumatra]. Medan State University. Geography Journal 2(2):17-30. [In Indonesian].

- Nusantara R. A., Rosyid A., Boesono H., 2014 Analysis of differences in depth of fishing area towards fish composition on Danish seine in Rembang waters. *Journal of Fisheries Resources Utilization Management and Technology* 3(3):96-104 [In Indonesian].
- Nuzapril M., Susilo S. B., Panjaitan J. P., 2017 Relationship between chlorophyll-a concentration with primary productivity rate using Landsat 8 imagery. *Journal of Fisheries and Marine Technology* 8(1):105-114.
- Pangestu S., 2016 [Simple linear regression. Applied linear model]. Digital Library of a Open University of Indonesia, 398 p. [In Indonesian].
- Sato K., Matsuno K., Arima D., Abe Y., Yamaguchi A., 2015 Spatial and temporal changes in zooplankton abundance, biovolume, and size spectra in the neighboring waters of Japan: Analyses using an optical plankton counter. *Zoological Studies. A Springer Open Journal* 54(18):1-15.
- Seki M. P., Polovina J. J., Kobayashi D. R., Bidigare R. R., Mitchum G. T., 2002 An oceanographic characterization of swordfish (*Xiphias gladius*) longline fishing grounds in the springtime subtropical North Pacific. *Fisheries Oceanography* 11(5):251-266.
- Sudirman, Mallawa A., 2012 [Fishing techniques]. Rineka Cipta, Jakarta, Indonesia, 168 p. [In Indonesian].
- Sugiari, Anshori M. A., Kusumawardani M., 2015 [Coordinate information system for estimated tuna fishing grounds for fishermen]. *Journal of JARTEL State Polytechnic of Malang* 1(1):74-84. [In Indonesian].
- Suniada K. I., Susilo E., Hastuti A. W., 2015 [Validation of forecast map of fishing grounds (PPDPI) in Java Sea Waters (FMA-RI 712)]. *Proceedings of the National Marine and Fisheries Science and Technology Forum* 87-96. [In Indonesian].
- Tambun R., Simbolon D., Wahyu R., Supartono, 2018 [Potential zones of fishing ground based on season at FMA RI 718]. *Journal of Tropical Marine Science and Technology* 10(3):757-768. [In Indonesian].
- Tuli M., Mennofatria B., Adrianto L., 2015 [Analysis of skipjack fish (*Katsuwonus pelamis*) resources in the waters of Pohuwato Regency, Gorontalo Province]. *Marine Fisheries Journal* 6(2):109-117. [In Indonesian].
- Wibawa T. A., 2011 [Utilization of oceanographic satellite data for prediction of potential fishing grounds for bigeye tuna (*T. obesus*) in the South Indian Ocean, Java and Bali]. *SEGARA Journal* 7(1):29-41. [In Indonesian].
- Zainuddin M., 2011 Skipjack tuna in relation to sea surface temperature and chlorophyll-a concentration of Bone Bay using remotely sensed satellite data. *Journal of Tropical Marine Science and Technology* 3(1):82-90.
- Zainuddin M., Safruddin, Farhum S. A., Nelwan A., Selamat M. B., Hidayat S., Sudirman, 2015 [Characteristics of potential areas for catching skipjack tuna in the Bone-Sea Bay of Flores based on satellite data on sea surface temperature and chlorophyll-a in the period January-June 2014 Makassar]. *Journal of Science and Technology Utilization of fisheries resources. Hasanuddin University* 2(3):228-237. [In Indonesian].
- Zhang C., Han M., 2015 Mapping chlorophyll-a concentration in Laizhou Bay using Landsat 8 OLI data. e-proceedings of the 36th IAHR World Congress, The Hague, The Netherlands, pp. 12-17.
- *** Institute for Marine Research and Observation, IMRO, 2019 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian],
- *** Institute for Marine Research and Observation, IMRO, 2020 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2016 [East Nusa Tenggara Provincial annual report 2016]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2018 [East Nusa Tenggara Provincial annual report 2018]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** <http://modis.gsfc.nasa.gov> 2007

Commented [WU15]: Missing from the text

Commented [A16R15]: There is. Komang, Change with Suniada

*** www.oceancolour.gfsc.nasa.gov

Received: 2021. Accepted: 2021. Published online: 2021.

Authors:

Maman Hermawan, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: mhermawan60@gmail.com

I Nyoman Radiarta, Institute for Marine Research and Observation, Agency for Marine and Fisheries Research and Development Baru Street, Perancak, Negara, Jembrana 82251 Bali, Indonesia, e-mail: radiarta@kkp.go.id

I Gusti Bagus Ega Krisna Bayu, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: egabayu91@gmail.com

Erick Nugraha, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: nugraha_eriq1@yahoo.co.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Hermawan M., Radiarta I. N., Bayu I. G. B. E. K., Nugraha E., 2021 ~~The suitability of the fishing ground with the forecast map of the fishing ground (PPDPI) in terms of chlorophyll-a in the Savu Sea, Indonesia.~~ [Comparison between fisherman fishing ground and the forecast map of fishing ground \(PPDPI\) based on chlorophyll-a in the Savu Sea, Indonesia.](#) AACL Bioflux 14(x):

← Kembali ↶ ↷ →

📁 Arsipkan 📁 Pindahkan 🗑️ Hapus 🛡️ Spam ⋮

☰ ▲ ▼ ✕

 **Eni Kovacs** <ek.bioflux@gmail.com>
Kepada: erick.nugraha

🖨️ Sen, 21 Jun 2021 jam 22.13 ☆


Dear Dr. Nugraha,

The manuscript does still not have the footer, as in the model from our website. As mentioned before, please **leave the footer exactly as it is, including the page numbering.**

> Tampilkan pesan asli

--
Kind regards,
Editor AACL Bioflux
Eniko Kovacs

↶ ↷ → ⋮



The suitability of the fishing ground with the forecast map of the fishing ground (PPDPI) in terms of chlorophyll-a in the Savu Sea, Indonesia

¹Maman Hermawan, ²I Nyoman Radiarta, ¹I Gusti B. E. K. Bayu, ¹Erick Nugraha

¹ Faculty of Fishing Technology, Jakarta Technical University of Fisheries, Pasarminggu, South Jakarta, Indonesia; ² Institute for Marine Research and Observation, Negara, Jembrana, Bali, Indonesia. Corresponding author: E. Nugraha, nugraha_eriq1@yahoo.co.id

Abstract. The Savu Sea in East Nusa Tenggara Province is the location of fishing activities for most fishermen in East Nusa Tenggara. Despite its potential, the knowledge of the fishing ground is still conventional. Currently, predictive methods for trips planning are available, such as the fishing ground forecast maps based on the chlorophyll-a, sea surface temperature and salinity. The related data are provided by NASA, being processed by the Marine Research and Observation Center to display the potential fishing ground as a box view with an accuracy level of 5 square miles. Eventually, the forecast map of the fishing area will be compared with the fishing locations known by the fishermen, for validation. The data collection used a survey method and the data analysis used both a descriptive analysis and the linear regression. The forecast maps matched with the observed fishing locations at only 13.6%. The highest recorded chlorophyll-a concentration was of 1.29 mg m⁻³ (constantly observed in the West Flores) and the lowest was of 0.07 mg m⁻³. The chlorophyll-a has an effect on 62.2% of the catch and 52% of the catch is composed of *Katsuwonus pelamis*.

The Savu Sea in East Nusa Tenggara Province is an area that is the location of fishing activities for most fishermen in East Nusa Tenggara. Behind all the potential possessed by the Savu Sea, the knowledge of the fishing ground is still conventional even though at this time various prediction maps of fishing ground have been issued which aim to help fishermen determine their fishing area when planning a voyage. This fishing ground forecast map is based on chlorophyll-a, sea surface temperature, salinity and current data obtained from NASA which is processed by the Marine Research and Observation Center which then displays potential fishing ground that have a box view with an accuracy level of 5 square miles. After that the forecast map of the fishing area will be compared with the fishing positions used by the fishermen. Through this scientific work, it is hoped that it will be able to become material for study by fishermen and become data validation by the agency that issued the map. The data collection method is a survey method and the data analysis method uses descriptive analysis and linear regression analysis. The results of the observation resulted in a map matching with the fishing position of the fishermen of 13.6% and the highest chlorophyll-a concentration of 1.29 mg m⁻³ and the lowest 0.07 mg m⁻³ where during the observation the West Flores area showed the highest concentration constantly. Partially chlorophyll-a has an effect of 62.2% on the catch, with the catch is dominated by *Katsuwonus pelamis* by 52%.

Key Words: MODIS, sea surface temperature, catch composition, *Katsuwonus pelamis*.

Introduction. Indonesia is the largest archipelagic country in the world, having a coastline length of 81,000 km² and a sea area of about 3,100,000 km² (Kusumastanto 1996; Darsono 1999; Latif et al 2019). Managing the fisheries resource requires to determine the fish growth patterns and the catch period, in order to avoid over-exploitation (Nugraha et al 2020b). The province of East Nusa Tenggara has a sea area of 200,000 km² with a coastline of 5,700 km² (MMAF 2016). Savu Sea has a good potential for the fisheries and the whole marine sector (Asagabaldan 2017). Indonesia is the largest archipelagic country in the world, has a coastline length of 81,000 km² and a sea area of about 3,100,000 km² (Kusumastanto 1996; Darsono 1999; Latif et al 2019). An important information that needs to be known in terms of fisheries resource management, is determining fish growth patterns and catch period, so fisherman

Commented [A1]: Please reformulate to be clear: the idea is to confirm the chlorophyll maps-predicted fishing grounds via field observation and fishermen consultation.

catching only feasible fish category (Nugraha et al 2020b). The province of East Nusa Tenggara has a sea area of 200,000 km² with a coastline of 5,700 km² (MMAF 2016). In the fisheries and marine sector, Savu Sea which has good fishery potential (Asagabaldan 2017).

A fishing ground is an area in the waters with an abundance of fish that are targeted to be caught and where the fishing gear can be operated technically and safely (Panjaitan 2014). The *Katsuwonus pelamis* is a fast swimming fish and has the greedy character of an opportunistic predator, with food types ranging from small fish (Clupeidae and Engraulidae), squids, crustaceans and zooplankton (Tuli et al 2015). *K. pelamis* is a pelagic fish living in groups, from the coastal areas to the open sea (Nainggolan 2012). Schooling fish should be kept as close to the boat as possible so that they can be easily caught using pole and line fishing rods (Sudirman & Mallawa 2004). The fishing grounds are determined by the waters' condition in the habitat of a species. Water conditions are usually described by oceanographic parameters (Nugraha et al 2020a).

A fishing ground is an area in the waters where there are many fish that are targeted to be caught and fishing gear can be operated technically and safely (Panjaitan 2014). The *Katsuwonus pelamis* is a fast swimming fish and has a greedy eating character and includes opportunistic predators with food types ranging from small fish (Clupeidae and Engraulidae), squids, crustaceans and Zooplankton (Tuli et al 2015). *K. pelamis* live in groups pelagically in coastal areas to the open sea (Nainggolan 2012). Schooling fish should be kept as close to the boat as possible so that they can be easily caught using pole and line fishing rods (Sudirman & Mallawa 2004). The determination of fishing ground can be expected from the condition of the waters where the habitat of a species. Water conditions are usually described by oceanographic parameters (Nugraha et al 2020a).

The Sea Surface Temperature (SST) of the *K. pelamis* fishing ground ranges from 29.75 to 30.25°C (Zainuddin et al 2015; Nugraha et al 2020b; Zainuddin 2011). Meanwhile, the potential area has a characteristic chlorophyll-a concentration of around 0.125 mg m⁻³. This condition is mainly seen in May. Outside of this range, the catch of *K. pelamis* tends to decline, as it happens in April (when the SST values in the Bone-Sea Flores bay are the highest). Likewise, when the chlorophyll-a concentration in the fishing ground significantly increase and the catch tends to decrease. This means that there is a certain range of values for the two oceanographic factors that are very suitable for the presence and abundance of *K. pelamis*. The distribution of chlorophyll-a can be a parameter used for identifying the presence of fish in a water area (Hasyim & Zainuddin 2008; Laurs et al 1984).

The characteristics of the Sea Surface Temperature (SST) of the *K. pelamis* fishing ground are in the range between 29.75 to 30.25°C (Zainuddin et al 2015; Nugraha et al 2020b; Zainuddin 2011). Meanwhile, the potential area has a characteristic chlorophyll-a concentration of around 0.125 mg m⁻³. This condition is mainly seen in May. The catch of *K. pelamis* tends to decline outside of this range, as happened in April (SST conditions are highest in the Bone Sea Flores bay). Likewise, when the chlorophyll-a concentration in the fishing ground has increased significantly, the catch tends to decrease. This means that there is a certain range of values for the two oceanographic factors that are very suitable for the presence and abundance of *K. pelamis*. To knowing the distribution of chlorophyll-a can be a parameter in identifying the presence of fish in an area of water (Hasyim & Zainuddin 2008; Laurs et al 1984).

The Forecast Maps of Fishing Ground (PPDPI) is the result of a research conducted by the Institute for Marine Research and Observation (IMRO) in order to identify fishing grounds, based on an internet service of information on the fishing areas' characteristics (Sugiar 2015; Panjaitan 2014). PPDPI have been produced for and distributed to the fishing communities of Indonesia, since 2000 (Tambun et al 2018; Farda & Jatisworo 2019). Forecast Maps of Fishing Ground (PPDPI) is the result of research conducted by Institute for Marine Research and Observation (IMRO) which is an effort to facilitate the search for fishing grounds in the form of internet information services based on fishing areas (Sugiar 2015; Panjaitan 2014). PPDPI have been produced and distributed since 2000 for fishing communities in Indonesia (Tambun et al 2018; Farda & Jatisworo 2019).

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Commented [WU2]: 2012 in the references.

Formatted: Font: Not Italic

Commented [Unknown A3]: In the whole manuscript, the authors have to replace "mg m³" by "mg m⁻³"

PDPPI was compiled using remote sensing data which is also widely used to detect water conditions such as the thermal front (Cayula & Cornillon 1992; Wibawa 2011; Jatisworo & Murdimanto 2013) and the concentration of chlorophyll-a (Seki et al 2002) which is one of the indicators of water fertility and school of fish. [PDPPI preparation was carried out using SST and water fertility data \(chlorophyll-a\) generated from MODIS Aqua level 3 daily composites, in the Hierarchical Data Format \(HDF\) \(Komang et al 2015\)](#) ~~PDPPI preparation was carried out using SST and water fertility data (chlorophyll-a) generated from MODIS Aqua level 3 composite 3 daily with Hierarchical Data Format (HDF) (Komang et al 2015).~~

Commented [WU4]: Not listed in the references

[The predictive maps of fishing grounds that already exists in Indonesia still have many weaknesses, including the spatial aspect of the points of forecasting located far from the coast and the uncertainty of the predicted type of fish. Thus, if inappropriate gear is selected for the fishing trip, the catch productivity is negatively affected, even in presence of abundant fish populations \(Kunarso et al 2019\).](#)

~~Map of prediction of fishing ground that already exists in Indonesia there are still many weaknesses including the spatial aspect of the point of forecasting far from the coast and the unclear type of fish predicted. The lack of understanding of the type of fish predicted, the risk of fishermen carrying the wrong fishing gear so that when reaching the forecasting location, fishermen can not catch fish that are abundant (Kunarso et al 2019).~~

Chlorophyll-a is an indicator of the abundance of phytoplankton in waters, ~~that~~ [playing](#) a role in the photosynthesis process (Huot et al 2007; Zhang & Han 2015). Phytoplankton contributes to a large extent to determine primary productivity in waters. Organic carbon production during photosynthesis is defined as primary productivity or Net Primary Productivity (Lee et al 2016).

Linear regression analysis was carried out on the model of the relationship between chlorophyll-a concentration and primary productivity. The results of this equation can be applied to [the](#) satellite imagery so that it can help ~~in-to~~ [monitoring the](#) water quality conditions (Nuzapril et al 2017). This abundance of phytoplankton will result in the creation of food chain clusters from phytoplankton to large pelagic fish schools (Sato et al 2015; Behrenfeld et al 2005).




Figure 1. Map of chlorophyll-a concentration (IMRO 2020).

Remote sensing data can show the variables used in determining fish catchment areas. ~~Certain One of the~~ remote sensing images used to extract SST and chlorophyll-a information ~~are provided by their a~~ MODIS ~~program image~~ (Nurman 2010; Hakim et al 2006).

[Satellite imagery is one of the applications of the remote sensing technology. The Moderate-resolution Imaging Spectroradiometer \(MODIS\) technology uses sensors belonging to the Earth Observing System \(EOS\), carried by two satellites produced by the](#)

National Aeronautics and Space Administration (NASA), namely Terra and Aqua (<http://modis.gsfc.nasa.gov> 2007).

Satellite imagery is one of the results of remote sensing technology. One of the technologies used is the Moderate resolution Imaging Spectroradiometer (MODIS) which is one of the sensors belonging to the Earth Observing System (EOS) and is carried by two vehicles produced by the National Aeronautics and Space Administration (NASA), namely Terra and Aqua (<http://modis.gsfc.nasa.gov> 2007).

The MODIS sensor is a derivative of the Advanced Very High Resolution Radiometer (AVHRR) sensor, Sea-viewing Wide Field-of-View Sensor (SeaWiFS) and High Resolution Imaging Spectrometer (HRS) owned by EOS, which has previously orbited, so it can be used to measure [sea and atmosphere-level parameters \(such as the sea surface temperature, chlorophyll-a concentration and water vapor content\) and marine phenomena \(such as the occurrence of upwelling, thermal fronts and others\)](#) parameters that are in at sea level to the atmosphere, among others, measures the sea surface temperature, chlorophyll-a concentration, water vapor content and marine phenomena, such as the occurrence of upwelling, thermal fronts and others (<http://modis.gsfc.nasa.gov> 2007).

Material and Method

Description of the study sites. This research was conducted on a pole and line fishing boat operating in the Flores Sea, East Nusa Tenggara. [The satellite image data retrieval was carried out at the Southeast Research and Ocean Observation Center \(SeaCORM\) Jembrana, Bali.](#) ~~For satellite image data retrieval is carried out at the Southeast Research and Ocean Observation Center (SeaCORM) Jembrana, Bali.~~ The material used [consisted of is](#) sea water ~~samples as an object~~ for measuring oceanographic elements. The methods used in data collection are literature study, interview, documentation, ~~and~~ observation.

Data analysis method. [Regression analysis is a statistical analysis that studies the relationship between two or more variables. In the linear regression analysis it is assumed that the this relationship is a linear combination, the simplest case consisting of one independent variable \(Pangestu 2016\).](#) ~~Regression analysis is a statistical analysis that studies the relationship between two or more variables. In linear regression analysis it is assumed that the form of a linear relationship in the parameters applies. The simplest linear regression module is linear regression with one independent variable (Pangestu 2016).~~

Linear regression is a statistical method used to model the relationship between the dependent variable (dependent; response; Y) and one or more independent variables (independent, predictor, X). If there ~~are is~~ only one independent variable, it is called simple linear regression, whereas if there is more than one independent variable, it is called multiple linear regression. [Regression analysis has at least three use cases, namely for data description, for control and for predictive purposes.](#) ~~Regression analysis has at least three uses, namely for the purpose of description of the data phenomenon or the case being studied, for control purposes, and for predictive purposes.~~

Simple regression is carried out in stages, namely by connecting the independent variable (chlorophyll-a) with the dependent variable (tuna catch, [using the data regression analysis function of Microsoft Office Excel \(Kurniawan 2008\).](#)) ~~In the use of Microsoft Office Excel to process data, namely by using data analysis then choosing data regression analysis (Kurniawan 2008).~~ This simple regression analysis was conducted to determine the relationship between the dependent variable Y (*K. pelamis* catches) and the independent variable X (chlorophyll-a). The regression model used is:

$$Y = a + bX$$

Where:

X - chlorophyll-a;

Y - pole and line catch;

a - the constant (intercept) is the value of Y, when X = 0 which equals the value of a where it intersects the Y axis;
~~constant (intercept), the value of a intersects the Y axis;~~
~~—The value of Y, when X = 0~~
 b - regression coefficient.

Chlorophyll-a image analysis. The fishing ground forecast can be predicted from the parameters of sea surface temperature and chlorophyll-a. ~~Determination of the fishing ground forecast can be predicted from the parameters of sea surface temperature and chlorophyll-a.~~ The analysis of the distribution of sea surface temperature and chlorophyll-a can use level 1b and level 2 Modis images. The Algorithm Theoretical Basic Document Modis (ATBD) 25 is used to determine sea surface temperature and the Morel 4 Algorithm is used to determine the chlorophyll-a. The Modis image levels 1b and 2 ~~using use~~ the RMSE test calculation (Chavula 2012).

From the two Root Mean Squared Error (RMSE) tests carried out between the field data and the Modis image data levels 1b and level 2, ~~the better and~~ more accurate ~~RMSE~~ values ~~is were~~ obtained from the smallest RMSE sum value, namely the Level 2 Modis image, with an RMSE value of chlorophyll-a of 0.090315. So the best fishing ground forecast can be determined through Level 2 Modis Images ~~So in determining the best fishing ground forecast is to use Level 2 Modis Image~~ (Febriani & Sukojo 2016).

Results and Discussion

Fishing ground. In this study we used a pole and line fishing boat (Figure 2) that operates in Savu sea, Flores, East Nusa Tenggara.



Figure 2. Pole and line fishing boat (original).

Composition and comparison of catches with the number of settings.

1. In November 2019 the ship made one trip with a catch of *K. pelamis* of 1,020 kg.
2. In December 2019 the ship made 16 trips with a catch of 28,220 kg. The fishing season generally reaches a turning point in the middle of January and in December, according to the graph, the catch will decrease as the wind season ~~arrives~~ enters.
3. In January 2020 the catch obtained was of 7,990 kg. January is generally the start of the wind season, so ships rarely go to sea.
4. In February 2020 the catches were dominated by *K. pelamis* and *T. albacares*. According to the fishermen, February was generally the peak of the wind season, so that the catch was not good, until the middle or end of March. ~~In February, fishermen said it was generally the peak of the wind season so that the catch was not good, this lasted until the middle or end of March.~~

5. In March 2020 the catches were dominated by *K. pelamis* and *T. albacares*, but unlike in the previous month, this month, *C. hippurus* and *E. bipinnulata* were also obtained. In March 2020 the fishermen began to increase the intensity of their trips, but taking into account the strength of the wind and the availability of bait. The composition of the catch for 5 months can be seen in Table 1 below:

Table 1
Comparison of [catch volumes](#) with the number of settings

Volumes / Period (kg [±])	November	December	January	February	March
Number of setting (times)	2	43	14	15	13
<i>K. pelamis</i> (kg)	1,020	15,300	2,890	6,290	7,480
<i>T. albacares</i> (kg)	0	9,520	2,040	5,440	4,250
<i>C. hippurus</i> (kg)	0	1,870	1,870	0	1,360
<i>E. bipinnulata</i> (kg)	0	1,530	1,190	0	1,870
Total (kg)	1,020	28,220	7,990	11,730	14,960
Catch rate per setting (kg)	510	656.28	570.71	782	1,150

Catch frequencies can also be visualized in Figure 3, below ~~we can be seen in the figure 3 below:~~

Commented [WU5]: Please reformulate, using scientific English.




Figure 3. Total catch (November 2019 to March 2020).

Based on the graph above, there is a decrease in the catch volumes during the 4 months of fishing. The lowest catch was obtained in January with a total of 7,990 kg, catch consisting of: *K. pelamis* (2,890 kg), *T. albacares* (2,040 kg) and others, such as *E. bipinnulata* and *C. hippurus* 7,990 kg consisting of *K. pelamis* 2,890 kg, *T. albacares* 2,040 kg and others were *E. bipinnulata* and *C. hippurus*. Meanwhile, the highest catch was obtained in December is 28,220 kg, dominated by *K. pelamis* with 15,300 kg and *T. albacares* with 9,520 kg. In 2016, Kupang Regency was the largest supplier of *K. pelamis* in East Nusa Tenggara (MMAF 2018).

Catch composition. If we see in Table 2 and figure 4 below, it is shown that *K. pelamis* is the most dominant catch that lands at Tenau port and at the fishing port around Kupang city, East Nusa Tenggara. This is in accordance with the data from the Fisheries Service of the Province of East Nusa Tenggara, suggesting that *K. pelamis* is one of the main commodities produced in Kupang, with 52%, followed by *T. albacares*, with 33%, *C. hippurus*, with 7% and *E. bipinnulata*, with 8%. This is in accordance with the statement and validating data from the Fisheries Service of the Province of East Nusa Tenggara that *K. pelamis* 52% is one of the main commodities produced in Kupang, followed by 33% *T. albacares*, 7% *C. hippurus* and 8% *E. bipinnulata*.

Table 2

Catch rated- per setting

Name of fish	Amount (kg)	Catch percentage (%)
<i>K. pelamis</i>	32,980	52
<i>T. albacares</i>	21,250	33
<i>C. hippurus</i>	4,590	7
<i>E. bipinnulata</i>	5,100	8
Total catching	63,920	100
Catch rate per setting	734.71	1.14

The percentage-catching compositions (in %) can be seen in Figure 4 below.




Figure 4. Catch composition diagram.

Commented [WU6]: Please display the full name of the species.

Conformity between of the fishing ground and PPDPI. Fishing is carried out on FADs that have been installed and with the position of the FADs stored on the GPS of each ship. The following is a map of the fishing grounds with the positions of the FADs. Figure 5 below shows the fishing ground for 5 months of fishing:




Figure 5. Fishing ground.

Forecast map of fishing ground (PPDPI).

1. In November the PPDPI that can be used correspond to 29 to 30 November 2019. On this map, the potential fishing ground are around the northwest of Timor Island with an area of 150 NM.
2. In December, 15 files were issued with an interval of 2 days. At PDDPI 11 to 12 December 2019, the recommended fishing areas are south of Savu Island, northeast

Commented [A7]: ? area unit, please.

of Sumba Island and north of Rote Island, as well as many other areas with an area of more than 300 NM (IMRO 2019).

3. In January, only 1 file out of 7 uploaded PPDPI files issued corresponded to the time of the fishing trip. On the 31 December to 01 January map, the potential fishing grounds were located around the southeast of Sumba Island, west of Savu Island, with an area of more than 300 nM. In January the PPDPI issued which corresponds to the time of the Catching trip, only 1 file out of 7 uploaded files. On the 31 December to 01 January map, the potential fishing grounds are around the southeast of Sumba Island which is one with the western area of Savu Island with an area of more than 300 NM (IMRO 2020).
4. In February, as many as 10 PPDPI files were uploaded and 3 files corresponded to the time of the fishing trip. One of the suitable map files was issued on the 26 to 27 February recommending fishing zones located north of Sumba Island with an area of 300 nM. Other areas do not seem to be potential fishing grounds. In February the PPDPI uploaded as many as 10 files and there are 3 files that correspond to the time of the catching trip. One suitable file is on the 26 to 27 February map with the recommended fishing area just north of Sumba Island with an area of 300 NM. Other areas do not show signs of potential fishing grounds (IMRO 2020).
5. In March there were 4 files matching, at the time of the fishing trip. March should be the turning point of the monsoon season that occurs in East Nusa Tenggara, but each region has a different time. One of the maps is the 16 to 17 March map, the areas that are recommended for fishing activities are in the south of Sumba Island and northeast of Sumba Island, with an area of more than 300 NM. In March there were 4 files that were used because they matched the time of the catching trip. March should be the turning point of the monsoon season that occurs in East Nusa Tenggara but each region has a different time. One of the maps is the 16 to 17 March map, the areas that are recommended for fishing activities are in the south of Sumba Island and northeast of Sumba Island with an area of more than 300 NM. There is was also one point with an area of 50 NM located on Semau and Kupang islands (IMRO 2020).

Commented [A8]: Please reformulate

Commented [A9]: Please correct the unit

Commented [A10]: Please correct the surface unit




Figure 6. PDPPI period in November (1), December (2) (IMRO 2019); January (3), February (4), March (5) (IMRO 2020).

Compatibility between the fishing ground and PDPPI. [Of all trips with available PDPPI, there were only 17 files that matched, at the time of the fishing trip. For the 4 months period, 17 maps of fishing ground forecasts were used, with a total of 44 settings \(Table 3\).](#)

[The method used to determine the suitability of fishing grounds with PDPPI, issued by IMRO, uses a comparison method between the position of the actual fishing area and the status of the same location on the forecast map, on that date. From 44 setting positions, 6 positions were in accordance with the PDPPI, which corresponds to a match rate of 13.6%, in terms of chlorophyll-a and sea surface temperature. When the daily PDPPI images from the NASA website are not available, the fishing trips cannot be used as a reference in determining their suitability.](#)

[The total trips was made and matched with the availability of PDPPI, there were only 17 files that matched the time of the fishing trip. For 4 months there were 17 map files forecasts of fishing ground used with a total of 44 settings \(Table 3\):](#)

[The method used to determine the suitability of fishing grounds with PDPPI issued by IMRO uses a comparison method between the position of the fishing area that has been carried out with the position of the forecast map of the appropriate fishing area on that date. After being matched from 44 setting positions, there are 6 positions that are in accordance with PDPPI, so it can be concluded by calculating the suitable position points compared to the total number of positions carried out multiplied by 100%, the level of compatibility of the fishing ground of fishermen with PDPPI which has a current reference is obtained, chlorophyll a and sea surface temperature have a match rate of 13.6%. Not all fishing trips can be used as a reference in determining suitability, because PDPPI](#)

cannot be uploaded every day because it is limited by the availability of images obtained from the NASA website.

With a match rate of 13.6%, it can be said that fishermen using FADs have sufficient accuracy in determining fishing grounds

Table 3


Compatibility-positions of the fishing ground with the PPDI

No	Setting	Position		Chlorophyll-a (mg m ⁻³)	Information
		Latitude	Longitude		
1	1	10° 12' 27" S	121° 40' 16" E	0.41	Inappropriate
	2	10° 09' 43" S	121° 37' 40" E	0.31	Inappropriate
2	1	09° 56' 38" S	122° 25' 03" E	0.3	Appropriate
	2	10° 42' 09" S	121° 50' 28" E	0.27	Inappropriate
	3	10° 26' 28" S	122° 47' 58" E	0.27	Appropriate
3	1	09° 32' 11" S	123° 20' 42" E	0.24	Inappropriate
	2	09° 45' 37" S	123° 13' 20" E	0.22	Inappropriate
4	1	10° 04' 18" S	122° 32' 21" E	0.27	Appropriate
	2	10° 28' 31" S	122° 41' 16" E	0.31	Inappropriate
	3	10° 12' 31" S	122° 05' 10" E	0.33	Appropriate
5	1	10° 40' 10" S	122° 15' 30" E	0.29	Inappropriate
	2	09° 40' 33" S	122° 55' 15" E	0.39	Inappropriate
6	1	09° 31' 40" S	123° 05' 09" E	0.21	Inappropriate
	2	09° 05' 45" S	123° 00' 41" E	0.28	Inappropriate
	3	09° 47' 16" S	122° 58' 34" E	0.13	Inappropriate
7	1	09° 40' 08" S	123° 22' 31" E	0.29	Inappropriate
	2	09° 50' 28" S	122° 53' 17" E	0.30	Inappropriate
8	1	10° 38' 15" S	121° 40' 30" E	0.41	Inappropriate
	2	10° 42' 09" S	121° 50' 28" E	0.38	Inappropriate
	3	10° 49' 25" S	121° 01' 51" E	-	Inappropriate
9	1	09° 32' 11" S	123° 20' 42" E	0.28	Inappropriate
	2	09° 45' 37" S	123° 13' 20" E	0.24	Inappropriate
	3	09° 54' 26" S	123° 01' 51" E	-	Inappropriate
10	1	10° 21' 15" S	122° 35' 07" E	0.45	Inappropriate
	2	10° 10' 35" S	122° 05' 37" E	-	Inappropriate
	3	10° 03' 47" S	122° 45' 10" E	0.54	Appropriate
11	1	09° 40' 36" S	122° 05' 15" E	0.8	Inappropriate
	2	09° 50' 17" S	121° 35' 20" E	1.1	Inappropriate
12	1	09° 30' 14" S	122° 50' 33" E	-	Inappropriate
	2	09° 21' 52" S	123° 00' 16" E	-	Inappropriate
13	1	10° 40' 16" S	123° 41' 40" E	-	Inappropriate
	2	10° 39' 13" S	123° 45' 36" E	-	Inappropriate
	3	10° 40' 41" S	123° 40' 22" E	-	Inappropriate
14	1	09° 50' 20" S	122° 10' 31" E	-	Inappropriate
	2	09° 18' 50" S	122° 46' 15" E	-	Inappropriate
	3	09° 30' 16" S	123° 01' 51" E	-	Inappropriate
15	1	10° 03' 15" S	122° 45' 17" E	0.31	Inappropriate
	2	10° 03' 40" S	121° 48' 29" E	-	Inappropriate
16	1	10° 08' 36" S	123° 07' 12" E	-	Inappropriate
	2	10° 15' 53" S	122° 46' 41" E	-	Inappropriate
	3	10° 03' 09" S	122° 53' 08" E	-	Inappropriate
17	1	09° 29' 21" S	122° 30' 14" E	0.6	Inappropriate
	2	09° 50' 15" S	122° 30' 35" E	0.54	Inappropriate
	3	09° 55' 23" S	121° 20' 54" E	0.49	Appropriate

Concentration and distribution of chlorophyll-a. Chlorophyll-a concentrations can be obtained by downloading data levels 1, 2 and 3 on the NASA website. Data needs in a longer time span such as months and years can use level 3 image data with better

quality than others, but level 3 images only contain one specific information depending on what is needed. In this study, the basis used is a level 3 image which only requires chlorophyll-a information with a time span of months. In Figure 7 and Table 4 the following is image data for chlorophyll-a concentration level 3.

[Chlorophyll-a concentrations can be obtained by downloading data levels 1, 2 and 3 from the NASA website. Data needs in a longer time span such as months and years can use level 3 image data with better quality than the others, but only containing one specific information, depending on what is needed. In this study, the basis used is a level 3 image which only requires chlorophyll-a information with a time span of months. Figure 7 images and Table 4 data correspond to level 3 chlorophyll-a concentration.](#)



March 2020

Figure 7. Concentration of chlorophyll-a map (www.oceancolour.gsfc.nasa.gov).

[The distribution map of chlorophyll-a concentrations \(Table 4\) from November to March, shows elevated values around the west of Flores Island, Sumba Island and also Larantuka, suggesting that food chain occur in the area.](#)

[By looking at the distribution map of chlorophyll-a concentrations \(Table 4\) that has been obtained from November to March, it shows that around the west of Flores Island, Sumba Island and also Larantuka have very thick chlorophyll a concentrations and it is indicated that a food chain occurs in the area.](#)

Table 4

Concentration and distribution of chlorophyll-a

<i>Period</i>	<i>November</i>	<i>December</i>	<i>January</i>	<i>February</i>	<i>March</i>
Concentration of chlorophyll-a (mg m ⁻³)					
a. Lowest	0.07	0.08	0.08	0.08	0.08
b. Highest	1.29	1.14	1.15	1.19	1.17
Average Value (mg m ⁻³)	0.2	0.3	0.2	0.2	0.2
Distribution center	North of Kupang city, Sabu Island, and west of Flores	West of Flores	Kupang city, Sabu Island, Rote Island, and west of Flores	Kupang city, Rote Island, and west of Flores	Kupang city, west of Flores
Compatibility level	Appropriate	Inappropriate	Appropriate	Appropriate	Inappropriate

Conclusions. Based on the results of the study it can be concluded: simultaneously, chlorophyll-a had an effect of 62.2% on the catch. The highest concentration of chlorophyll-a recorded was 1.29 mg m⁻³ and the lowest was 0.07 mg m⁻³; the fishing area used by fishermen has a level of conformity with the forecast map for fishing areas of 13.6%; Not all fishing trips can be used as a reference in determining the suitability of fishing grounds with PDPPI, because not every day PDPPI can be uploaded because it is limited by the availability of images obtained from the NASA website.

Acknowledgments. The authors would like to thank to Director and staff of Institute for Marine Research and Observation (IMRO), Agency for Marine and Fisheries Research and Development, Ministry of Marine Affairs and Fisheries, Indonesia. for the support to sharing and processing of the satellite data related to the distribution of SST and Chlorophyll-a in fishing ground and to all other parties for their help and contributions.

Conflict of interest. The authors declare no conflict of interest.

References

Asagabaldan M. A., 2017 Law and the role of Lilifuk Amid Government Regimes: Legal analysis and sustainability of local wisdom in Kuanheun Village, West Kupang Regency, East Nusa Tenggara. SABDA 12(2):136-145.

- Behrenfeld M. J., Boss E., Siegel D. A., Shea D. M., 2005 Carbon-based ocean productivity and phytoplankton physiology from space. *Global Biogeochemical Cycles* 19(1):1-14.
- Cayula J. F., Cornillon P., 1992 Edge detection algorithm for SST Images. *Journal of Atmospheric and Oceanic Technology* 9(1):67-80.
- Chavula G., Sungani H., Gondwe K., 2012 Mapping potential fishing grounds in Lake Malawi using AVHRR and MODIS satellite imagery. *International Journal of Geosciences* 3(03):650-658.
- Darsono P., 1999 [Utilization of marine resources and its implications for fishing communities]. *Oceana* 24(4):1-9. [In Indonesian].
- Farda N. M., Jatisworo D., 2019 [Mining space-time patterns on forecast maps of fishing areas in fisheries management areas (FMA) 712, 713, and 573]. *Scientific Magazine GLOBE* 21(2):117-128. [In Indonesian].
- Febriani E. R., Sukojo B. M., 2016 [Comparative analysis of the use of level 1b and level 2 MODIS imagery in determining forecasting fishing areas (Case Study: South Coast of Blitar)]. *ITS Engineering Journal, Surabaya* 5(2):439-442. [In Indonesian].
- Hakim D. M., Wikantika K., Widiadnyana N., Napitu A. M., Darmawan S., 2006 The identification of fishing ground area with MODIS satellite image (Case Study: South Coast of West Java) *Proceeding. ITB Engineering Science* 38B(2):147-158.
- Hasyim S., Zainuddin M., 2008 Predictions of skipjack fishing areas based on oceanographic conditions in the waters of Takalar regency and surrounding areas. *Journal of Science and Technology* 8(2):158-162. [In Indonesian].
- Huot Y., Babin M., Bruyant F., Grob C., Twardowski M. S., Claustre H., 2007 Does chlorophyll-a provide the best index of phytoplankton biomass for primary productivity studies? *Biogeosciences Discussions, Proxies of Biomass For Primary production* 4(2):707-745.
- Jatisworo D., Murdimanto A., 2013 [Identification of thermal fronts in the Makassar Strait and Banda Sea]. *Proceedings of the National Symposium on Geoinformation Science III. Marine Research and Observation Institute*, 8 p. [In Indonesian].
- Kunarso, Zainuri M., Sugianto D. N., Marwoto J., Hariyadi, Muslim, 2019 Prediction of demersal fishing ground associated with coral reefs in the Coastal Jepara Regency, Central Java, Indonesia based on Sentinel 2a imagery. *Advances in Science, Technology and Engineering Systems (ASTES) Journal* 4(6):263-269.
- Kurniawan D., 2008 [Linear regression]. *Academia accelerating the world's research. Research Development Core Team R: A language and environment for statistical computing*, 14 p. [In Indonesian].
- Kusumastanto T., 1996 [An evaluation of government policies on the development of fisheries in Indonesia]. *AGRIS, Fisheries Economy Bulletin* 2(3):27-41. [In Indonesian].
- Latif B., Noor S. M., Sumardi J., Irwansyah, 2019 Sustainable development and sea protection: Trade on fish and fishery product. *ICROEST. IOP Conference Series: Earth and Environmental Science* 343(1):1-6.
- Laurs R. M., Fiedler P. C., Montgomery D. R., 1984 Albacore tuna catch distribution relative to environmental features observed from satellites. *Deep-Sea Research* 31:1085-1099.
- Lee Y. J., Matrai P. A., Friedrichs M. A. M., Saba V. S., Aumont O., Babin M., Buitenhuis E. T., Chevallier M., Mora L., Dessert D., Dunne J. P., Ellingsen I. H., Feldman D., Frouin R., Gehlen M., Gorgues T., Ilyina T., Jin M., John J. G., Lawrence J., Manizza M., Menkes C. E., Perruche C., Fouest V. L., Popova E. E., Romanou A., Samuelson A., Schwinger J., Séférian R., Stock C. A., Tjiputra J., Tremblay L. B., Ueyoshi K., Vichi M., Yool A., Zhang J., 2016 Net primary productivity estimates and environmental variables in the Arctic Ocean: An assessment of coupled physical-biogeochemical models. *Journal of Geophysical Research: Oceans* 121(12):8635-8669.
- Nainggolan C., 2012 [Fishing methods]. *Open University, Jakarta*, 61 p. [In Indonesian].

- Nugraha E., Gunawan R., Danapraja S., Yusrizal, Kusdinar A., Waluyo A. S., Hutajulu J., Prayitno H., Halim S., Sutisna D. H., 2020b The sea surface temperature effect on the length and size of skipjack tuna (*Katsuwonus pelamis*) catches in the Banda Sea, Indonesia. *AAFL Bioflux* 13(1):1-18.
- Nugraha E., Yudho G. S., Jaenudin A., Yusrizal, Kusmedy B., Kusnidar A., Husen E. S., 2020a Relationship between length and weight of skipjack tuna (*Katsuwonus Pelamis*) purse seine catching in the Maluku Sea, Indonesia. *AAFL Bioflux* 13(1):330-345.
- Nurman A., 2010 [Utilization of fashionable data to detect fishing grounds on the east and west coasts of North Sumatra]. Medan State University. *Geography Journal* 2(2):17-30. [In Indonesian].
- Nuzapril M., Susilo S. B., Panjaitan J. P., 2017 Relationship between chlorophyll-a concentration with primary productivity rate using Landsat 8 imagery. *Journal of Fisheries and Marine Technology* 8(1):105-114.
- Pangestu S., 2016 [Simple linear regression. Applied linear model]. Digital Library of a Open University of Indonesia, 398 p. [In Indonesian].
- Panjaitan H., 2014 [Utilization of fishing area forecast map]. fisheries writing book. ocean-wide-ki.blogspot.com [In Indonesian].
- Sato K., Matsuno K., Arima D., Abe Y., Yamaguchi A., 2015 Spatial and temporal changes in zooplankton abundance, biovolume, and size spectra in the neighboring waters of Japan: Analyses using an optical plankton counter. *Zoological Studies. A Springer Open Journal* 54(18):1-15.
- Seki M. P., Polovina J. J., Kobayashi D. R., Bidigare R. R., Mitchum G. T., 2002 An oceanographic characterization of swordfish (*Xiphias gladius*) longline fishing grounds in the springtime subtropical North Pacific. *Fisheries Oceanography* 11(5):251-266.
- Sudirman, Mallawa A., 2012 [Fishing techniques]. Rineka Cipta, Jakarta, Indonesia, 168 p. [In Indonesian].
- Sugiari, Anshori M. A., Kusumawardani M., 2015 [Coordinate information system for estimated tuna fishing grounds for fishermen]. *Journal of JARTEL State Polytechnic of Malang* 1(1):74-84. [In Indonesian].
- Suniada K. I., Susilo E., Hastuti A. W., 2015 [Validation of forecast map of fishing grounds (PPDPI) in Java Sea Waters (FMA-RI 712)]. *Proceedings of the National Marine and Fisheries Science and Technology Forum* 87-96. [In Indonesian].
- Tambun R., Simbolon D., Wahju R., Supartono, 2018 [Potential zones of fishing ground based on season at FMA RI 718]. *Journal of Tropical Marine Science and Technology* 10(3):757-768. [In Indonesian].
- Tuli M., Mennofatria B., Adrianto L., 2015 [Analysis of skipjack fish (*Katsuwonus pelamis*) resources in the waters of Pohuwato Regency, Gorontalo Province]. *Marine Fisheries Journal* 6(2):109-117. [In Indonesian].
- Wibawa T. A., 2011 [Utilization of oceanographic satellite data for prediction of potential fishing grounds for bigeye tuna (*T. obesus*) in the South Indian Ocean, Java and Bali]. *SEGARA Journal* 7(1):29-41. [In Indonesian].
- Zainuddin M., 2011 Skipjack tuna in relation to sea surface temperature and chlorophyll-a concentration of Bone Bay using remotely sensed satellite data. *Journal of Tropical Marine Science and Technology* 3(1):82-90.
- Zainuddin M., Safruddin, Farhum S. A., Nelwan A., Selamat M. B., Hidayat S., Sudirman, 2015 [Characteristics of potential areas for catching skipjack tuna in the Bone-Sea Bay of Flores based on satellite data on sea surface temperature and chlorophyll-a in the period January-June 2014 Makassar]. *Journal of Science and Technology Utilization of fisheries resources. Hasanuddin University* 2(3):228-237. [In Indonesian].
- Zhang C., Han M., 2015 Mapping chlorophyll-a concentration in Laizhou Bay using Landsat 8 OLI data. e-proceedings of the 36th IAHR World Congress, The Hague, The Netherlands, pp. 12-17.
- *** Institute for Marine Research and Observation, IMRO, 2019 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian],

Commented [WU11]: No of pages.

Commented [WU12]: Missing from the text

- *** Institute for Marine Research and Observation, IMRO, 2020 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2016 [East Nusa Tenggara Provincial annual report 2016]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2018 [East Nusa Tenggara Provincial annual report 2018]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** <http://modis.gsfc.nasa.gov> 2007
- *** www.oceancolour.gsfc.nasa.gov

Received: 2021. Accepted: 2021. Published online: 2021.

Authors:

Maman Hermawan, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street

Pasarminggu 12520, South Jakarta, Indonesia, e-mail: mhermawan60@gmail.com

I Nyoman Radiarta, Institute for Marine Research and Observation, Agency for Marine and Fisheries Research

and Development Baru Street, Perancak, Negara, Jembrana 82251 Bali, Indonesia, e-mail: radiarta@kkp.go.id

I Gusti Bagus Ega Krisna Bayu, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP

Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: egabayu91@gmail.com

Erick Nugraha, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu

12520, South Jakarta, Indonesia, e-mail: nugraha_eriq1@yahoo.co.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Hermawan M., Radiarta I. N., Bayu I. G. B. E. K., Nugraha E., 2021 The suitability of the fishing ground with the forecast map of the fishing ground (PPDPI) in terms of chlorophyll-a in the Savu Sea, Indonesia. *AAFL Bioflux* 14(x):

← Kembali ↩ ↶ ↷

📁 Arsipkan 📁 Pindahkan 🗑 Hapus 🛡 Spam ⋮

☰ ▲ ▼ ✕

• Revision 4

Yahoo/Terkirim ☆

 **Eni Kovacs** <ek.bioflux@gmail.com>
Kepada: erick nugraha

🖨 📧 Min, 18 Jul 2021 jam 00.21 ☆

Dear Dr. Nugraha,

Regarding your manuscript submitted to AACL Bioflux, the editorial team has some requests prior to acceptance (please see the attachment).

Please note: Always operate corrections/additions (or deletions) in the manuscript we are sending you, by **highlighting with a bright color** (for an easy identification). We never work on the manuscript you send back, just identify the corrections and operate them on our document (to avoid any undesirable accidental operations like changed page set up, otherwise the editors have to start all the work from the beginning, and we cannot ask them to re-check every manuscript word by word to identify unmarked modifications).


Thank you for understanding!

--
Kind regards,
Editor AACL Bioflux
Eniko Kovacs



Nugraha_PP....doc
1.6MB

↩ ↶ ↷ ⋮



The suitability of the fishing ground with the forecast map of the fishing ground (PPDPI) in terms of chlorophyll-a in the Savu Sea, Indonesia

¹Maman Hermawan, ²I Nyoman Radiarta, ¹I Gusti B. E. K. Bayu, ¹Erick Nugraha

¹ Faculty of Fishing Technology, Jakarta Technical University of Fisheries, Pasarminggu, South Jakarta, Indonesia; ² Institute for Marine Research and Observation, Negara, Jembrana, Bali, Indonesia. Corresponding author: E. Nugraha, nugraha_eriq1@yahoo.co.id

Abstract. The Savu Sea in East Nusa Tenggara Province is the location of fishing activities for most fishermen in East Nusa Tenggara. Despite its potential, the knowledge of the fishing ground is still conventional. Currently, predictive methods for trips planning are available, such as the fishing ground forecast maps based on the chlorophyll-a, sea surface temperature and salinity. The related data are provided by NASA, being processed by the Marine Research and Observation Center to display the potential fishing ground as a box view with an accuracy level of 5 square miles. Eventually, the forecast map of the fishing area will be compared with the fishing locations known by the fishermen, for validation. The data collection used a survey method and the data analysis used both a descriptive analysis and the linear regression. The forecast maps matched with the observed fishing locations at only 13.6%. The highest recorded chlorophyll-a concentration was of 1.29 mg m⁻³ (constantly observed in the West Flores) and the lowest was of 0.07 mg m⁻³. The chlorophyll-a has an effect on 62.2% of the catch and 52% of the catch is composed of *Katsuwonus pelamis*.

The Savu Sea in East Nusa Tenggara Province is an area that is the location of fishing activities for most fishermen in East Nusa Tenggara. Behind all the potential possessed by the Savu Sea, the knowledge of the fishing ground is still conventional even though at this time various prediction maps of fishing ground have been issued which aim to help fishermen determine their fishing area when planning a voyage. This fishing ground forecast map is based on chlorophyll-a, sea surface temperature, salinity and current data obtained from NASA which is processed by the Marine Research and Observation Center which then displays potential fishing ground that have a box view with an accuracy level of 5 square miles. After that the forecast map of the fishing area will be compared with the fishing positions used by the fishermen. Through this scientific work, it is hoped that it will be able to become material for study by fishermen and become data validation by the agency that issued the map. The data collection method is a survey method and the data analysis method uses descriptive analysis and linear regression analysis. The results of the observation resulted in a map matching with the fishing position of the fishermen of 13.6% and the highest chlorophyll-a concentration of 1.29 mg m⁻³ and the lowest 0.07 mg m⁻³ where during the observation the West Flores area showed the highest concentration constantly. Partially chlorophyll-a has an effect of 62.2% on the catch, with the catch is dominated by *Katsuwonus pelamis* by 52%.

Key Words: MODIS, sea surface temperature, catch composition, *Katsuwonus pelamis*.

Introduction. Indonesia is the largest archipelagic country in the world, having a coastline length of 81,000 km² and a sea area of about 3,100,000 km² (Kusumastanto 1996; Darsono 1999; Latif et al 2019). Managing the fisheries resource requires to determine the fish growth patterns and the catch period, in order to avoid over-exploitation (Nugraha et al 2020b). The province of East Nusa Tenggara has a sea area of 200,000 km² with a coastline of 5,700 km² (MMAF 2016). Savu Sea has a good potential for the fisheries and the whole marine sector, (Asagabaldan 2017). Indonesia is the largest archipelagic country in the world, has a coastline length of 81,000 km² and a sea area of about 3,100,000 km² (Kusumastanto 1996; Darsono 1999; Latif et al 2019). An important information that needs to be known in terms of fisheries resource management, is determining fish growth patterns and catch period, so fisherman

Commented [A1]: Please reformulate to be clear: the idea is to confirm the chlorophyll maps-predicted fishing grounds via field observation and fishermen consultation.

catching only feasible fish category (Nugraha et al 2020b). The province of East Nusa Tenggara has a sea area of 200,000 km² with a coastline of 5,700 km² (MMAF 2016). In the fisheries and marine sector, Savu Sea which has good fishery potential (Asagabaldan 2017).

A fishing ground is an area in the waters with an abundance of fish that are targeted to be caught and where the fishing gear can be operated technically and safely (Panjaitan 2014). The *Katsuwonus pelamis* is a fast swimming fish and has the greedy character of an opportunistic predator, with food types ranging from small fish (Clupeidae and Engraulidae), squids, crustaceans and zooplankton (Tuli et al 2015). *K. pelamis* is a pelagic fish living in groups, from the coastal areas to the open sea (Nainggolan 2012). Schooling fish should be kept as close to the boat as possible so that they can be easily caught using pole and line fishing rods (Sudirman & Mallawa 2004). The fishing grounds are determined by the waters' condition in the habitat of a species. Water conditions are usually described by oceanographic parameters (Nugraha et al 2020a).

A fishing ground is an area in the waters where there are many fish that are targeted to be caught and fishing gear can be operated technically and safely (Panjaitan 2014). The *Katsuwonus pelamis* is a fast swimming fish and has a greedy eating character and includes opportunistic predators with food types ranging from small fish (Clupeidae and Engraulidae), squids, crustaceans and Zooplankton (Tuli et al 2015). *K. pelamis* live in groups pelagically in coastal areas to the open sea (Nainggolan 2012). Schooling fish should be kept as close to the boat as possible so that they can be easily caught using pole and line fishing rods (Sudirman & Mallawa 2004). The determination of fishing ground can be expected from the condition of the waters where the habitat of a species. Water conditions are usually described by oceanographic parameters (Nugraha et al 2020a).

The Sea Surface Temperature (SST) of the *K. pelamis* fishing ground ranges from 29.75 to 30.25°C (Zainuddin et al 2015; Nugraha et al 2020b; Zainuddin 2011). Meanwhile, the potential area has a characteristic chlorophyll-a concentration of around 0.125 mg m⁻³. This condition is mainly seen in May. Outside of this range, the catch of *K. pelamis* tends to decline, as it happens in April (when the SST values in the Bone-Sea Flores bay are the highest). Likewise, when the chlorophyll-a concentration in the fishing ground significantly increase and the catch tends to decrease. This means that there is a certain range of values for the two oceanographic factors that are very suitable for the presence and abundance of *K. pelamis*. The distribution of chlorophyll-a can be a parameter used for identifying the presence of fish in a water area (Hasyim & Zainuddin 2008; Laurs et al 1984).

The characteristics of the Sea Surface Temperature (SST) of the *K. pelamis* fishing ground are in the range between 29.75 to 30.25°C (Zainuddin et al 2015; Nugraha et al 2020b; Zainuddin 2011). Meanwhile, the potential area has a characteristic chlorophyll-a concentration of around 0.125 mg m⁻³. This condition is mainly seen in May. The catch of *K. pelamis* tends to decline outside of this range, as happened in April (SST conditions are highest in the Bone Sea Flores bay). Likewise, when the chlorophyll-a concentration in the fishing ground has increased significantly, the catch tends to decrease. This means that there is a certain range of values for the two oceanographic factors that are very suitable for the presence and abundance of *K. pelamis*. To knowing the distribution of chlorophyll-a can be a parameter in identifying the presence of fish in an area of water (Hasyim & Zainuddin 2008; Laurs et al 1984).

The Forecast Maps of Fishing Ground (PPDPI) is the result of a research conducted by the Institute for Marine Research and Observation (IMRO) in order to identify fishing grounds, based on an internet service of information on the fishing areas' characteristics (Sugiar 2015; Panjaitan 2014). PPDPI have been produced for and distributed to the fishing communities of Indonesia, since 2000 (Tambun et al 2018; Farda & Jatisworo 2019). Forecast Maps of Fishing Ground (PPDPI) is the result of research conducted by Institute for Marine Research and Observation (IMRO) which is an effort to facilitate the search for fishing grounds in the form of internet information services based on fishing areas (Sugiar 2015; Panjaitan 2014). PPDPI have been produced and distributed since 2000 for fishing communities in Indonesia (Tambun et al 2018; Farda & Jatisworo 2019).

Formatted: Font: Not Italic

Formatted: Font: Not Italic

Commented [WU2]: 2012 in the references.

Formatted: Font: Not Italic

Commented [Unknown A3]: In the whole manuscript, the authors have to replace "mg m³" by "mg m⁻³"

PDPPI was compiled using remote sensing data which is also widely used to detect water conditions such as the thermal front (Cayula & Cornillon 1992; Wibawa 2011; Jatisworo & Murdimanto 2013) and the concentration of chlorophyll-a (Seki et al 2002) which is one of the indicators of water fertility and school of fish. [PDPPI preparation was carried out using SST and water fertility data \(chlorophyll-a\) generated from MODIS Aqua level 3 daily composites, in the Hierarchical Data Format \(HDF\) \(Komang et al 2015\)](#)~~PDPPI preparation was carried out using SST and water fertility data (chlorophyll-a) generated from MODIS Aqua level 3 composite 3 daily with Hierarchical Data Format (HDF) (Komang et al 2015).~~

Commented [WU4]: Not listed in the references

[The predictive maps of fishing grounds that already exists in Indonesia still have many weaknesses, including the spatial aspect of the points of forecasting located far from the coast and the uncertainty of the predicted type of fish. Thus, if inappropriate gear is selected for the fishing trip, the catch productivity is negatively affected, even in presence of abundant fish populations \(Kunarso et al 2019\).](#)

~~Map of prediction of fishing ground that already exists in Indonesia there are still many weaknesses including the spatial aspect of the point of forecasting far from the coast and the unclear type of fish predicted. The lack of understanding of the type of fish predicted, the risk of fishermen carrying the wrong fishing gear so that when reaching the forecasting location, fishermen can not catch fish that are abundant (Kunarso et al 2019).~~

Chlorophyll-a is an indicator of the abundance of phytoplankton in waters, ~~that~~ [playing](#) a role in the photosynthesis process (Huot et al 2007; Zhang & Han 2015). Phytoplankton contributes to a large extent to determine primary productivity in waters. Organic carbon production during photosynthesis is defined as primary productivity or Net Primary Productivity (Lee et al 2016).

Linear regression analysis was carried out on the model of the relationship between chlorophyll-a concentration and primary productivity. The results of this equation can be applied to [the](#) satellite imagery so that it can help ~~in-to~~ [monitoring the](#) water quality conditions (Nuzapril et al 2017). This abundance of phytoplankton will result in the creation of food chain clusters from phytoplankton to large pelagic fish schools (Sato et al 2015; Behrenfeld et al 2005).




Figure 1. Map of chlorophyll-a concentration (IMRO 2020).

Remote sensing data can show the variables used in determining fish catchment areas. ~~Certain~~ [One of the](#) remote sensing images used to extract SST and chlorophyll-a information ~~are provided by theis a~~ [MODIS program image](#) (Nurman 2010; Hakim et al 2006).

[Satellite imagery is one of the applications of the remote sensing technology. The Moderate-resolution Imaging Spectroradiometer \(MODIS\) technology uses sensors belonging to the Earth Observing System \(EOS\), carried by two satellites produced by the](#)

National Aeronautics and Space Administration (NASA), namely Terra and Aqua (<http://modis.gsfc.nasa.gov> 2007).

Satellite imagery is one of the results of remote sensing technology. One of the technologies used is the Moderate resolution Imaging Spectroradiometer (MODIS) which is one of the sensors belonging to the Earth Observing System (EOS) and is carried by two vehicles produced by the National Aeronautics and Space Administration (NASA), namely Terra and Aqua (<http://modis.gsfc.nasa.gov> 2007).

The MODIS sensor is a derivative of the Advanced Very High Resolution Radiometer (AVHRR) sensor, Sea-viewing Wide Field-of-View Sensor (SeaWiFS) and High Resolution Imaging Spectrometer (HRS) owned by EOS, which has previously orbited, so it can be used to measure [sea and atmosphere-level parameters \(such as the sea surface temperature, chlorophyll-a concentration and water vapor content\) and marine phenomena \(such as the occurrence of upwelling, thermal fronts and others\)](#) parameters that are in at sea level to the atmosphere, among others, measures the sea surface temperature, chlorophyll-a concentration, water vapor content and marine phenomena, such as the occurrence of upwelling, thermal fronts and others (<http://modis.gsfc.nasa.gov> 2007).

Material and Method

Description of the study sites. This research was conducted on a pole and line fishing boat operating in the Flores Sea, East Nusa Tenggara. [The satellite image data retrieval was carried out at the Southeast Research and Ocean Observation Center \(SeaCORM\) Jembrana, Bali.](#) ~~For satellite image data retrieval is carried out at the Southeast Research and Ocean Observation Center (SeaCORM) Jembrana, Bali.~~ The material used [consisted of is](#) sea water ~~samples as an object~~ for measuring oceanographic elements. The methods used in data collection are literature study, interview, documentation, ~~and~~ observation.

Data analysis method. [Regression analysis is a statistical analysis that studies the relationship between two or more variables. In the linear regression analysis it is assumed that the this relationship is a linear combination, the simplest case consisting of one independent variable \(Pangestu 2016\).](#) ~~Regression analysis is a statistical analysis that studies the relationship between two or more variables. In linear regression analysis it is assumed that the form of a linear relationship in the parameters applies. The simplest linear regression module is linear regression with one independent variable (Pangestu 2016).~~

Linear regression is a statistical method used to model the relationship between the dependent variable (dependent; response; Y) and one or more independent variables (independent, predictor, X). If there ~~are is~~ only one independent variable, it is called simple linear regression, whereas if there is more than one independent variable, it is called multiple linear regression. [Regression analysis has at least three use cases, namely for data description, for control and for predictive purposes.](#) ~~Regression analysis has at least three uses, namely for the purpose of description of the data phenomenon or the case being studied, for control purposes, and for predictive purposes.~~

Simple regression is carried out in stages, namely by connecting the independent variable (chlorophyll-a) with the dependent variable (tuna catch, [using the data regression analysis function of Microsoft Office Excel \(Kurniawan 2008\).](#)) ~~In the use of Microsoft Office Excel to process data, namely by using data analysis then choosing data regression analysis (Kurniawan 2008).~~ This simple regression analysis was conducted to determine the relationship between the dependent variable Y (*K. pelamis* catches) and the independent variable X (chlorophyll-a). The regression model used is:

$$Y = a + bX$$

Where:

X - chlorophyll-a;

Y - pole and line catch;

a - the constant (intercept) is the value of Y, when X = 0 which equals the value of a where it intersects the Y axis;
~~constant (intercept), the value of a intersects the Y axis;~~
~~The value of Y, when X = 0~~
 b - regression coefficient.

Chlorophyll-a image analysis. ~~The fishing ground forecast can be predicted from the parameters of sea surface temperature and chlorophyll-a. Determination of the fishing ground forecast can be predicted from the parameters of sea surface temperature and chlorophyll-a.~~ The analysis of the distribution of sea surface temperature and chlorophyll-a can use level 1b and level 2 Modis images. The Algorithm Theoretical Basic Document Modis (ATBD) 25 is used to determine sea surface temperature and the Morel 4 Algorithm is used to determine the chlorophyll-a. The Modis image levels 1b and 2 ~~using use~~ the RMSE test calculation (Chavula 2012).

From the two Root Mean Squared Error (RMSE) tests carried out between the field data and the Modis image data levels 1b and level 2, ~~the better and~~ more accurate ~~RMSE~~ values ~~is were~~ obtained from the smallest RMSE sum value, namely the Level 2 Modis image, with an RMSE value of chlorophyll-a of 0.090315. So the best fishing ground forecast can be determined through Level 2 Modis Images~~So in determining the best fishing ground forecast is to use Level 2 Modis Image~~ (Febriani & Sukojo 2016).

Results and Discussion

Fishing ground. In this study we used a pole and line fishing boat (Figure 2) that operates in Savu sea, Flores, East Nusa Tenggara.



Figure 2. Pole and line fishing boat (original).

Composition and comparison of catches with the number of settings.

1. In November 2019 the ship made one trip with a catch of *K. pelamis* of 1,020 kg.
2. In December 2019 the ship made 16 trips with a catch of 28,220 kg. The fishing season generally reaches a turning point in the middle of January and in December, according to the graph, the catch will decrease as the wind season ~~arrives~~ has entered.
3. In January 2020 the catch obtained was of 7,990 kg. January is generally the start of the wind season, so ships rarely go to sea.
4. In February 2020 the catches were dominated by *K. pelamis* and *T. albacares*. According to the fishermen, February was generally the peak of the wind season, so that the catch was not good, until the middle or end of March. ~~In February, fishermen said it was generally the peak of the wind season so that the catch was not good, this lasted until the middle or end of March.~~

5. In March 2020 the catches were dominated by *K. pelamis* and *T. albacares*, but unlike in the previous month, this month, *C. hippurus* and *E. bipinnulata* were also obtained. In March 2020 the fishermen began to increase the intensity of their trips, but taking into account the strength of the wind and the availability of bait. The composition of the catch for 5 months can be seen in Table 1 below:

Table 1
Comparison of [catch volumes](#) with the number of settings

Volumes / Period (kg [±])	November	December	January	February	March
Number of setting (times)	2	43	14	15	13
<i>K. pelamis</i> (kg)	1,020	15,300	2,890	6,290	7,480
<i>T. albacares</i> (kg)	0	9,520	2,040	5,440	4,250
<i>C. hippurus</i> (kg)	0	1,870	1,870	0	1,360
<i>E. bipinnulata</i> (kg)	0	1,530	1,190	0	1,870
Total (kg)	1,020	28,220	7,990	11,730	14,960
Catch rate per setting (kg)	510	656.28	570.71	782	1,150

Catch frequencies can also be visualized in Figure 3, below ~~we can be seen in the figure 3-below:~~

Commented [WU5]: Please reformulate, using scientific English.




Figure 3. Total catch (November 2019 to March 2020).

Based on the graph above, there is a decrease in the catch volumes during the 4 months of fishing. The lowest catch was obtained in January with a total of 7,990 kg, catch consisting of: *K. pelamis* (2,890 kg), *T. albacares* (2,040 kg) and others, such as *E. bipinnulata* and *C. hippurus*. Meanwhile, the highest catch was obtained in December is 28,220 kg, dominated by *K. pelamis* with 15,300 kg and *T. albacares* with 9,520 kg. In 2016, Kupang Regency was the largest supplier of *K. pelamis* in East Nusa Tenggara (MMAF 2018).

Catch composition. If we see in Table 2 and figure 4 below, it is shown that *K. pelamis* is the most dominant catch that lands at Tenau port and at the fishing port around Kupang city, East Nusa Tenggara. This is in accordance with the data from the Fisheries Service of the Province of East Nusa Tenggara, suggesting that *K. pelamis* is one of the main commodities produced in Kupang, with 52%, followed by *T. albacares*, with 33%, *C. hippurus*, with 7% and *E. bipinnulata*, with 8%. This is in accordance with the statement and validating data from the Fisheries Service of the Province of East Nusa Tenggara that *K. pelamis* 52% is one of the main commodities produced in Kupang, followed by 33% *T. albacares*, 7% *C. hippurus* and 8% *E. bipinnulata*.

Table 2

Catch rated- per setting

Name of fish	Amount (kg)	Catch percentage (%)
<i>K. pelamis</i>	32,980	52
<i>T. albacares</i>	21,250	33
<i>C. hippurus</i>	4,590	7
<i>E. bipinnulata</i>	5,100	8
Total catching	63,920	100
Catch rate per setting	734.71	1.14

The percentage-catching compositions (in %) can be seen in Figure 4 below.




Figure 4. Catch composition diagram.

Commented [WU6]: Please display the full name of the species.

Conformity between of the fishing ground and PPDPI. Fishing is carried out on FADs that have been installed and with the position of the FADs stored on the GPS of each ship. The following is a map of the fishing grounds with the positions of the FADs. Figure 5 below shows the fishing ground for 5 months of fishing:




Figure 5. Fishing ground.

Forecast map of fishing ground (PPDPI).

1. In November the PPDPI that can be used correspond to 29 to 30 November 2019. On this map, the potential fishing ground are around the northwest of Timor Island with an area of 150 NM.
2. In December, 15 files were issued with an interval of 2 days. At PDPPI 11 to 12 December 2019, the recommended fishing areas are south of Savu Island, northeast

Commented [A7]: ? area unit, please.

of Sumba Island and north of Rote Island, as well as many other areas with an area of more than 300 NM (IMRO 2019).

3. In January, only 1 file out of 7 uploaded PPDPI files issued corresponded to the time of the fishing trip. On the 31 December to 01 January map, the potential fishing grounds were located around the southeast of Sumba Island, west of Savu Island, with an area of more than 300 nM. In January the PPDPI issued which corresponds to the time of the Catching trip, only 1 file out of 7 uploaded files. On the 31 December to 01 January map, the potential fishing grounds are around the southeast of Sumba Island which is one with the western area of Savu Island with an area of more than 300 NM (IMRO 2020).
4. In February, as many as 10 PPDPI files were uploaded and 3 files corresponded to the time of the fishing trip. One of the suitable map files was issued on the 26 to 27 February recommending fishing zones located north of Sumba Island with an area of 300 nM. Other areas do not seem to be potential fishing grounds. In February the PPDPI uploaded as many as 10 files and there are 3 files that correspond to the time of the catching trip. One suitable file is on the 26 to 27 February map with the recommended fishing area just north of Sumba Island with an area of 300 NM. Other areas do not show signs of potential fishing grounds (IMRO 2020).
5. In March there were 4 files matching, at the time of the fishing trip. March should be the turning point of the monsoon season that occurs in East Nusa Tenggara, but each region has a different time. One of the maps is the 16 to 17 March map, the areas that are recommended for fishing activities are in the south of Sumba Island and northeast of Sumba Island, with an area of more than 300 NM. In March there were 4 files that were used because they matched the time of the catching trip. March should be the turning point of the monsoon season that occurs in East Nusa Tenggara but each region has a different time. One of the maps is the 16 to 17 March map, the areas that are recommended for fishing activities are in the south of Sumba Island and northeast of Sumba Island with an area of more than 300 NM. There is was also one point with an area of 50 NM located on Semau and Kupang islands (IMRO 2020).

Commented [A8]: Please reformulate

Commented [A9]: Please correct the unit

Commented [A10]: Please correct the surface unit




Figure 6. PDPPI period in November (1), December (2) (IMRO 2019); January (3), February (4), March (5) (IMRO 2020).

Compatibility between the fishing ground and PDPPI. [Of all trips with available PDPPI, there were only 17 files that matched, at the time of the fishing trip. For the 4 months period, 17 maps of fishing ground forecasts were used, with a total of 44 settings \(Table 3\).](#)

[The method used to determine the suitability of fishing grounds with PDPPI, issued by IMRO, uses a comparison method between the position of the actual fishing area and the status of the same location on the forecast map, on that date. From 44 setting positions, 6 positions were in accordance with the PDPPI, which corresponds to a match rate of 13.6%, in terms of chlorophyll-a and sea surface temperature. When the daily PDPPI images from the NASA website are not available, the fishing trips cannot be used as a reference in determining their suitability.](#)

[The total trips was made and matched with the availability of PDPPI, there were only 17 files that matched the time of the fishing trip. For 4 months there were 17 map files forecasts of fishing ground used with a total of 44 settings \(Table 3\):](#)

[The method used to determine the suitability of fishing grounds with PDPPI issued by IMRO uses a comparison method between the position of the fishing area that has been carried out with the position of the forecast map of the appropriate fishing area on that date. After being matched from 44 setting positions, there are 6 positions that are in accordance with PDPPI, so it can be concluded by calculating the suitable position points compared to the total number of positions carried out multiplied by 100%, the level of compatibility of the fishing ground of fishermen with PDPPI which has a current reference is obtained, chlorophyll a and sea surface temperature have a match rate of 13.6%. Not all fishing trips can be used as a reference in determining suitability, because PDPPI](#)

cannot be uploaded every day because it is limited by the availability of images obtained from the NASA website.

With a match rate of 13.6%, it can be said that fishermen using FADs have sufficient accuracy in determining fishing grounds

Table 3


Compatibility positions of the fishing ground with the PPDI

No	Setting	Position		Chlorophyll-a (mg m ⁻³)	Information
		Latitude	Longitude		
1	1	10° 12' 27" S	121° 40' 16" E	0.41	Inappropriate
	2	10° 09' 43" S	121° 37' 40" E	0.31	Inappropriate
2	1	09° 56' 38" S	122° 25' 03" E	0.3	Appropriate
	2	10° 42' 09" S	121° 50' 28" E	0.27	Inappropriate
3	3	10° 26' 28" S	122° 47' 58" E	0.27	Appropriate
	1	09° 32' 11" S	123° 20' 42" E	0.24	Inappropriate
4	2	09° 45' 37" S	123° 13' 20" E	0.22	Inappropriate
	1	10° 04' 18" S	122° 32' 21" E	0.27	Appropriate
5	2	10° 28' 31" S	122° 41' 16" E	0.31	Inappropriate
	3	10° 12' 31" S	122° 05' 10" E	0.33	Appropriate
6	1	10° 40' 10" S	122° 15' 30" E	0.29	Inappropriate
	2	09° 40' 33" S	122° 55' 15" E	0.39	Inappropriate
7	1	09° 31' 40" S	123° 05' 09" E	0.21	Inappropriate
	2	09° 05' 45" S	123° 00' 41" E	0.28	Inappropriate
8	3	09° 47' 16" S	122° 58' 34" E	0.13	Inappropriate
	1	09° 40' 08" S	123° 22' 31" E	0.29	Inappropriate
9	2	09° 50' 28" S	122° 53' 17" E	0.30	Inappropriate
	1	10° 38' 15" S	121° 40' 30" E	0.41	Inappropriate
10	2	10° 42' 09" S	121° 50' 28" E	0.38	Inappropriate
	3	10° 49' 25" S	121° 01' 51" E	-	Inappropriate
11	1	09° 32' 11" S	123° 20' 42" E	0.28	Inappropriate
	2	09° 45' 37" S	123° 13' 20" E	0.24	Inappropriate
12	3	09° 54' 26" S	123° 01' 51" E	-	Inappropriate
	1	10° 21' 15" S	122° 35' 07" E	0.45	Inappropriate
13	2	10° 10' 35" S	122° 05' 37" E	-	Inappropriate
	3	10° 03' 47" S	122° 45' 10" E	0.54	Appropriate
14	1	09° 40' 36" S	122° 05' 15" E	0.8	Inappropriate
	2	09° 50' 17" S	121° 35' 20" E	1.1	Inappropriate
15	1	09° 30' 14" S	122° 50' 33" E	-	Inappropriate
	2	09° 21' 52" S	123° 00' 16" E	-	Inappropriate
16	1	10° 40' 16" S	123° 41' 40" E	-	Inappropriate
	2	10° 39' 13" S	123° 45' 36" E	-	Inappropriate
17	3	10° 40' 41" S	123° 40' 22" E	-	Inappropriate
	1	09° 50' 20" S	122° 10' 31" E	-	Inappropriate
18	2	09° 18' 50" S	122° 46' 15" E	-	Inappropriate
	3	09° 30' 16" S	123° 01' 51" E	-	Inappropriate
19	1	10° 03' 15" S	122° 45' 17" E	0.31	Inappropriate
	2	10° 03' 40" S	121° 48' 29" E	-	Inappropriate
20	1	10° 08' 36" S	123° 07' 12" E	-	Inappropriate
	2	10° 15' 53" S	122° 46' 41" E	-	Inappropriate
21	3	10° 03' 09" S	122° 53' 08" E	-	Inappropriate
	1	09° 29' 21" S	122° 30' 14" E	0.6	Inappropriate
22	2	09° 50' 15" S	122° 30' 35" E	0.54	Inappropriate
	3	09° 55' 23" S	121° 20' 54" E	0.49	Appropriate

Concentration and distribution of chlorophyll-a. Chlorophyll-a concentrations can be obtained by downloading data levels 1, 2 and 3 on the NASA website. Data needs in a longer time span such as months and years can use level 3 image data with better

quality than others, but level 3 images only contain one specific information depending on what is needed. In this study, the basis used is a level 3 image which only requires chlorophyll-a information with a time span of months. In Figure 7 and Table 4 the following is image data for chlorophyll-a concentration level 3.

[Chlorophyll-a concentrations can be obtained by downloading data levels 1, 2 and 3 from the NASA website. Data needs in a longer time span such as months and years can use level 3 image data with better quality than the others, but only containing one specific information, depending on what is needed. In this study, the basis used is a level 3 image which only requires chlorophyll-a information with a time span of months. Figure 7 images and Table 4 data correspond to level 3 chlorophyll-a concentration.](#)



March 2020

Figure 7. Concentration of chlorophyll-a map (www.oceancolour.gsfc.nasa.gov).

The distribution map of chlorophyll-a concentrations (Table 4) from November to March, shows elevated values around the west of Flores Island, Sumba Island and also Larantuka, suggesting that food chain occur in the area.

By looking at the distribution map of chlorophyll-a concentrations (Table 4) that has been obtained from November to March, it shows that around the west of Flores Island, Sumba Island and also Larantuka have very thick chlorophyll-a concentrations and it is indicated that a food chain occurs in the area.

Table 4

Concentration and distribution of chlorophyll-a

<i>Period</i>	<i>November</i>	<i>December</i>	<i>January</i>	<i>February</i>	<i>March</i>
Concentration of chlorophyll-a (mg m ⁻³)					
a. Lowest	0.07	0.08	0.08	0.08	0.08
b. Highest	1.29	1.14	1.15	1.19	1.17
Average Value (mg m ⁻³)	0.2	0.3	0.2	0.2	0.2
Distribution center	North of Kupang city, Sabu Island, and west of Flores	West of Flores	Kupang city, Sabu Island, Rote Island, and west of Flores	Kupang city, Rote Island, and west of Flores	Kupang city, west of Flores
Compatibility level	Appropriate	Inappropriate	Appropriate	Appropriate	Inappropriate

Conclusions. Based on the results of the study it can be concluded: simultaneously, chlorophyll-a had an effect of 62.2% on the catch. The highest concentration of chlorophyll-a recorded was 1.29 mg m⁻³ and the lowest was 0.07 mg m⁻³; the fishing area used by fishermen has a level of conformity with the forecast map for fishing areas of 13.6%; Not all fishing trips can be used as a reference in determining the suitability of fishing grounds with PDPPI, because not every day PDPPI can be uploaded because it is limited by the availability of images obtained from the NASA website.

Acknowledgments. The authors would like to thank to Director and staff of Institute for Marine Research and Observation (IMRO), Agency for Marine and Fisheries Research and Development, Ministry of Marine Affairs and Fisheries, Indonesia. for the support to sharing and processing of the satellite data related to the distribution of SST and Chlorophyll-a in fishing ground and to all other parties for their help and contributions.

Conflict of interest. The authors declare no conflict of interest.

References

Asagabaldan M. A., 2017 Law and the role of Lilifuk Amid Government Regimes: Legal analysis and sustainability of local wisdom in Kuanheun Village, West Kupang Regency, East Nusa Tenggara. SABDA 12(2):136-145.

- Behrenfeld M. J., Boss E., Siegel D. A., Shea D. M., 2005 Carbon-based ocean productivity and phytoplankton physiology from space. *Global Biogeochemical Cycles* 19(1):1-14.
- Cayula J. F., Cornillon P., 1992 Edge detection algorithm for SST Images. *Journal of Atmospheric and Oceanic Technology* 9(1):67-80.
- Chavula G., Sungani H., Gondwe K., 2012 Mapping potential fishing grounds in Lake Malawi using AVHRR and MODIS satellite imagery. *International Journal of Geosciences* 3(03):650-658.
- Darsono P., 1999 [Utilization of marine resources and its implications for fishing communities]. *Oceana* 24(4):1-9. [In Indonesian].
- Farda N. M., Jatisworo D., 2019 [Mining space-time patterns on forecast maps of fishing areas in fisheries management areas (FMA) 712, 713, and 573]. *Scientific Magazine GLOBE* 21(2):117-128. [In Indonesian].
- Febriani E. R., Sukojo B. M., 2016 [Comparative analysis of the use of level 1b and level 2 MODIS imagery in determining forecasting fishing areas (Case Study: South Coast of Blitar)]. *ITS Engineering Journal, Surabaya* 5(2):439-442. [In Indonesian].
- Hakim D. M., Wikantika K., Widiadnyana N., Napitu A. M., Darmawan S., 2006 The identification of fishing ground area with MODIS satellite image (Case Study: South Coast of West Java) *Proceeding. ITB Engineering Science* 38B(2):147-158.
- Hasyim S., Zainuddin M., 2008 Predictions of skipjack fishing areas based on oceanographic conditions in the waters of Takalar regency and surrounding areas. *Journal of Science and Technology* 8(2):158-162. [In Indonesian].
- Huot Y., Babin M., Bruyant F., Grob C., Twardowski M. S., Claustre H., 2007 Does chlorophyll-a provide the best index of phytoplankton biomass for primary productivity studies? *Biogeosciences Discussions, Proxies of Biomass For Primary production* 4(2):707-745.
- Jatisworo D., Murdimanto A., 2013 [Identification of thermal fronts in the Makassar Strait and Banda Sea]. *Proceedings of the National Symposium on Geoinformation Science III. Marine Research and Observation Institute*, 8 p. [In Indonesian].
- Kunarso, Zainuri M., Sugianto D. N., Marwoto J., Hariyadi, Muslim, 2019 Prediction of demersal fishing ground associated with coral reefs in the Coastal Jepara Regency, Central Java, Indonesia based on Sentinel 2a imagery. *Advances in Science, Technology and Engineering Systems (ASTES) Journal* 4(6):263-269.
- Kurniawan D., 2008 [Linear regression]. *Academia accelerating the world's research. Research Development Core Team R: A language and environment for statistical computing*, 14 p. [In Indonesian].
- Kusumastanto T., 1996 [An evaluation of government policies on the development of fisheries in Indonesia]. *AGRIS, Fisheries Economy Bulletin* 2(3):27-41. [In Indonesian].
- Latif B., Noor S. M., Sumardi J., Irwansyah, 2019 Sustainable development and sea protection: Trade on fish and fishery product. *ICROEST. IOP Conference Series: Earth and Environmental Science* 343(1):1-6.
- Laurs R. M., Fiedler P. C., Montgomery D. R., 1984 Albacore tuna catch distribution relative to environmental features observed from satellites. *Deep-Sea Research* 31:1085-1099.
- Lee Y. J., Matrai P. A., Friedrichs M. A. M., Saba V. S., Aumont O., Babin M., Buitenhuis E. T., Chevallier M., Mora L., Dessert D., Dunne J. P., Ellingsen I. H., Feldman D., Frouin R., Gehlen M., Gorgues T., Ilyina T., Jin M., John J. G., Lawrence J., Manizza M., Menkes C. E., Perruche C., Fouest V. L., Popova E. E., Romanou A., Samuelson A., Schwinger J., Séférian R., Stock C. A., Tjiputra J., Tremblay L. B., Ueyoshi K., Vichi M., Yool A., Zhang J., 2016 Net primary productivity estimates and environmental variables in the Arctic Ocean: An assessment of coupled physical-biogeochemical models. *Journal of Geophysical Research: Oceans* 121(12):8635-8669.
- Nainggolan C., 2012 [Fishing methods]. *Open University, Jakarta*, 61 p. [In Indonesian].

- Nugraha E., Gunawan R., Danapraja S., Yusrizal, Kusdinar A., Waluyo A. S., Hutajulu J., Prayitno H., Halim S., Sutisna D. H., 2020b The sea surface temperature effect on the length and size of skipjack tuna (*Katsuwonus pelamis*) catches in the Banda Sea, Indonesia. *AACL Bioflux* 13(1):1-18.
- Nugraha E., Yudho G. S., Jaenudin A., Yusrizal, Kusmedy B., Kusnidar A., Husen E. S., 2020a Relationship between length and weight of skipjack tuna (*Katsuwonus Pelamis*) purse seine catching in the Maluku Sea, Indonesia. *AACL Bioflux* 13(1):330-345.
- Nurman A., 2010 [Utilization of fashionable data to detect fishing grounds on the east and west coasts of North Sumatra]. Medan State University. *Geography Journal* 2(2):17-30. [In Indonesian].
- Nuzapril M., Susilo S. B., Panjaitan J. P., 2017 Relationship between chlorophyll-a concentration with primary productivity rate using Landsat 8 imagery. *Journal of Fisheries and Marine Technology* 8(1):105-114.
- Pangestu S., 2016 [Simple linear regression. Applied linear model]. Digital Library of a Open University of Indonesia, 398 p. [In Indonesian].
- Panjaitan H., 2014 [Utilization of fishing area forecast map]. fisheries writing book. ocean-wide-ki.blogspot.com [In Indonesian].
- Sato K., Matsuno K., Arima D., Abe Y., Yamaguchi A., 2015 Spatial and temporal changes in zooplankton abundance, biovolume, and size spectra in the neighboring waters of Japan: Analyses using an optical plankton counter. *Zoological Studies. A Springer Open Journal* 54(18):1-15.
- Seki M. P., Polovina J. J., Kobayashi D. R., Bidigare R. R., Mitchum G. T., 2002 An oceanographic characterization of swordfish (*Xiphias gladius*) longline fishing grounds in the springtime subtropical North Pacific. *Fisheries Oceanography* 11(5):251-266.
- Sudirman, Mallawa A., 2012 [Fishing techniques]. Rineka Cipta, Jakarta, Indonesia, 168 p. [In Indonesian].
- Sugiari, Anshori M. A., Kusumawardani M., 2015 [Coordinate information system for estimated tuna fishing grounds for fishermen]. *Journal of JARTEL State Polytechnic of Malang* 1(1):74-84. [In Indonesian].
- Suniada K. I., Susilo E., Hastuti A. W., 2015 [Validation of forecast map of fishing grounds (PPDPI) in Java Sea Waters (FMA-RI 712)]. *Proceedings of the National Marine and Fisheries Science and Technology Forum* 87-96. [In Indonesian].
- Tambun R., Simbolon D., Wahju R., Supartono, 2018 [Potential zones of fishing ground based on season at FMA RI 718]. *Journal of Tropical Marine Science and Technology* 10(3):757-768. [In Indonesian].
- Tuli M., Mennofatria B., Adrianto L., 2015 [Analysis of skipjack fish (*Katsuwonus pelamis*) resources in the waters of Pohuwato Regency, Gorontalo Province]. *Marine Fisheries Journal* 6(2):109-117. [In Indonesian].
- Wibawa T. A., 2011 [Utilization of oceanographic satellite data for prediction of potential fishing grounds for bigeye tuna (*T. obesus*) in the South Indian Ocean, Java and Bali]. *SEGARA Journal* 7(1):29-41. [In Indonesian].
- Zainuddin M., 2011 Skipjack tuna in relation to sea surface temperature and chlorophyll-a concentration of Bone Bay using remotely sensed satellite data. *Journal of Tropical Marine Science and Technology* 3(1):82-90.
- Zainuddin M., Safruddin, Farhum S. A., Nelwan A., Selamat M. B., Hidayat S., Sudirman, 2015 [Characteristics of potential areas for catching skipjack tuna in the Bone-Sea Bay of Flores based on satellite data on sea surface temperature and chlorophyll-a in the period January-June 2014 Makassar]. *Journal of Science and Technology Utilization of fisheries resources. Hasanuddin University* 2(3):228-237. [In Indonesian].
- Zhang C., Han M., 2015 Mapping chlorophyll-a concentration in Laizhou Bay using Landsat 8 OLI data. e-proceedings of the 36th IAHR World Congress, The Hague, The Netherlands, pp. 12-17.
- *** Institute for Marine Research and Observation, IMRO, 2019 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian],

Commented [WU11]: No of pages.

Commented [WU12]: Missing from the text

- *** Institute for Marine Research and Observation, IMRO, 2020 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2016 [East Nusa Tenggara Provincial annual report 2016]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2018 [East Nusa Tenggara Provincial annual report 2018]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** <http://modis.gsfc.nasa.gov> 2007
- *** www.oceancolour.gsfc.nasa.gov

Received: 2021. Accepted: 2021. Published online: 2021.

Authors:

Maman Hermawan, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street

Pasarminggu 12520, South Jakarta, Indonesia, e-mail: mhermawan60@gmail.com

I Nyoman Radiarta, Institute for Marine Research and Observation, Agency for Marine and Fisheries Research

and Development Baru Street, Perancak, Negara, Jembrana 82251 Bali, Indonesia, e-mail: radiarta@kkp.go.id

I Gusti Bagus Ega Krisna Bayu, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP

Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: egabayu91@gmail.com

Erick Nugraha, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu

12520, South Jakarta, Indonesia, e-mail: nugraha_eriq1@yahoo.co.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Hermawan M., Radiarta I. N., Bayu I. G. B. E. K., Nugraha E., 2021 The suitability of the fishing ground with the forecast map of the fishing ground (PPDPI) in terms of chlorophyll-a in the Savu Sea, Indonesia. *AAFL Bioflux* 14(x):

← Kembali ↶ ↷ →

📁 Arsipkan 📁 Pindahkan 🗑️ Hapus 🛡️ Spam ⋮

☰ ▲ ▼ ✕



erick nugraha <nugraha_eriq1@yahoo.co.id>
Kepada: Eni Kovacs

🖨️ 🔗 Min, 18 Jul 2021 jam 11.16 ☆

Dear Dr. Eni,
I send You edited Manuscript.


thank you

> Tampilkan pesan asli



Nugraha_PP...doc
1.5MB

↶ ↷ → ⋮



~~The suitability of the fishing ground with the forecast map of the fishing ground (PPDPI) in terms of chlorophyll-a in the Savu Sea, Indonesia~~ Comparison between fisherman fishing ground and the forecast map of fishing ground (PPDPI) based on chlorophyll-a in the Savu Sea, Indonesia

¹Maman Hermawan, ²I Nyoman Radiarta, ¹I Gusti B. E. K. Bayu, ¹Erick Nugraha

¹ Faculty of Fishing Technology, Jakarta Technical University of Fisheries, Pasarminggu, South Jakarta, Indonesia; ² Institute for Marine Research and Observation, Negara, Jembrana, Bali, Indonesia. Corresponding author: E. Nugraha, nugraha_eriq1@yahoo.co.id

Abstract. The Savu Sea in East Nusa Tenggara Province is the location of fishing activities for most fishermen in East Nusa Tenggara. Despite its potential, the knowledge of the fishing ground is still conventional. Currently, predictive methods for trips planning are available, such as the fishing ground forecast maps based on the chlorophyll-a, sea surface temperature and salinity. The related data are provided by NASA, being processed by the Marine Research and Observation Center to display the potential fishing ground as a box view with an accuracy level of 5 square miles. Eventually, the forecast map of the fishing area will be compared with the fishing locations known by the fishermen, for validation. The data collection used a survey method and the data analysis used both a descriptive analysis and the linear regression. The forecast maps matched with the observed fishing locations at only 13.6%. The highest recorded chlorophyll-a concentration was of 1.29 mg m^{-3} (constantly observed in the West Flores) and the lowest was of 0.07 mg m^{-3} . The chlorophyll-a has an effect on 62.2% of the catch and 52% of the catch is composed of *Katsuwonus pelamis*.

Key Words: MODIS, sea surface temperature, catch composition, *Katsuwonus pelamis*.

Introduction. Indonesia is the largest archipelagic country in the world, having a coastline length of 81,000 km² and a sea area of about 3,100,000 km² (Kusumastanto 1996; Darsono 1999; Latif et al 2019). Managing the fisheries resource requires to determine the fish growth patterns and the catch period, in order to avoid over-exploitation (Nugraha et al 2020b). The province of East Nusa Tenggara has a sea area of 200,000 km² with a coastline of 5,700 km² (MMAF 2016). Savu Sea has a good potential for the fisheries and the whole marine sector (Asagabaldan 2017). ~~A fishing ground is an area in the waters with an abundance of fish that are targeted to be caught and where the fishing gear can be operated technically and safely (Panjaitan 2014).~~ Fishing area is an area where fish are caught in maximum amount and fishing gear can be operated economical. Assessment and mapping of fishing areas are needed in fishing effort (Nusantara et al 2014). The *Katsuwonus pelamis* is a fast swimming fish and has the greedy character of an opportunistic predator, with food types ranging from small fish (Clupeidae and Engraulidae), squids, crustaceans and zooplankton (Tuli et al 2015). *K. pelamis* is a pelagic fish living in groups, from the coastal areas to the open sea (Nainggolan 2012). Schooling fish should be kept as close to the boat as possible so that they can be easily caught using pole and line fishing rods (Sudirman & Mallawa 2004, 2012). The fishing grounds are determined by the waters' condition in the habitat of

Commented [A1]: Please reformulate to be clear: the idea is to confirm the chlorophyll maps-predicted fishing grounds via field observation and fishermen consultation.

Commented [WU2]: 2012 in the references.

Commented [A3R2]: Corrected thank you

a species. Water conditions are usually described by oceanographic parameters (Nugraha et al 2020a).


The Sea Surface Temperature (SST) of the *K. pelamis* fishing ground ranges from 29.75 to 30.25°C (Zainuddin et al 2015; Nugraha et al 2020b; Zainuddin 2011). Meanwhile, the potential area has a characteristic chlorophyll-a concentration of around 0.125 mg m⁻³. This condition is mainly seen in May. Outside of this range, the catch of *K. pelamis* tends to decline, as it happens in April (when the SST values in the Bone-Sea Flores bay are the highest). Likewise, when the chlorophyll-a concentration in the fishing ground significantly increase and the catch tends to decrease. This means that there is a certain range of values for the two oceanographic factors that are very suitable for the presence and abundance of *K. pelamis*. The distribution of chlorophyll-a can be a parameter used for identifying the presence of fish in a water area (Hasyim & Zainuddin 2008; Laurs et al 1984).

The Forecast Maps of Fishing Ground (PPDPI) is the result of a research conducted by the Institute for Marine Research and Observation (IMRO) in order to identify fishing grounds, based on an internet service of information on the fishing areas characteristics (Sugiari 2015). PPDPI have been produced for and distributed to the fishing communities of Indonesia, since 2000 (Tambun et al 2018; Farda & Jatisworo 2019). PDPPI was compiled using remote sensing data which is also widely used to detect water conditions such as the thermal front (Cayula & Cornillon 1992; Wibawa 2011; Jatisworo & Murdimanto 2013) and the concentration of chlorophyll-a (Seki et al 2002) which is one of the indicators of water fertility and school of fish. PDPPI preparation was carried out using SST and water fertility data (chlorophyll-a) generated from MODIS Aqua level 3 daily composites, in the Hierarchical Data Format (HDF) (Komang Suniada et al 2015)

The predictive maps of fishing grounds that already exists in Indonesia still have many weaknesses, including the spatial aspect of the points of forecasting located far from the coast and the uncertainty of the predicted type of fish. Thus, if inappropriate gear is selected for the fishing trip, the catch productivity is negatively affected, even in presence of abundant fish populations (Kunarso et al 2019).

Chlorophyll-a is an indicator of the abundance of phytoplankton in waters, playing a role in the photosynthesis process (Huot et al 2007; Zhang & Han 2015). Phytoplankton contributes to a large extent to determine primary productivity in waters. Organic carbon production during photosynthesis is defined as primary productivity or Net Primary Productivity (Lee et al 2016).

Linear regression analysis was carried out on the model of the relationship between chlorophyll-a concentration and primary productivity. The results of this equation can be applied to the satellite imagery so that it can help to monitor the water quality conditions (Nuzapril et al 2017). This abundance of phytoplankton will result in the creation of food chain clusters from phytoplankton to large pelagic fish schools (Sato et al 2015; Behrenfeld et al 2005).



Commented [Unknown A4]: In the whole manuscript, the authors have to replace "mg m³" by "mg m⁻³"

Commented [A5R4]: Thank You.

Commented [WU6]: Not listed in the references

Commented [A7R6]: Corrected : full named is Iwan Komang Suniada. Thank you.

Figure 1. Map of chlorophyll-a concentration (IMRO 2020).

Remote sensing data can show the variables used in determining fish catchment areas. Certain remote sensing images used to extract SST and chlorophyll-a information are provided by the MODIS program (Nurman 2010; Hakim et al 2006).

Satellite imagery is one of the applications of the remote sensing technology. The Moderate-resolution Imaging Spectroradiometer (MODIS) technology uses sensors belonging to the Earth Observing System (EOS), carried by two satellites produced by the National Aeronautics and Space Administration (NASA), namely Terra and Aqua (<http://modis.gsfc.nasa.gov> 2007).

The MODIS sensor is a derivative of the Advanced Very High Resolution Radiometer (AVHRR) sensor, Sea-viewing Wide Field-of-View Sensor (SeaWiFS) and High Resolution Imaging Spectrometer (HRS) owned by EOS, which has previously orbited, so it can be used to measure sea and atmosphere-level parameters (such as the sea surface temperature, chlorophyll-a concentration and water vapor content) and marine phenomena (such as the occurrence of upwelling, thermal fronts and others) (<http://modis.gsfc.nasa.gov> 2007).

Material and Method

Description of the study sites. This research was conducted on a pole and line fishing boat operating in the Flores Sea, East Nusa Tenggara. The satellite image data retrieval was carried out at the Southeast Research and Ocean Observation Center (SeaCORM) Jembrana, Bali. The material used consisted of sea water samples for measuring oceanographic elements. The methods used in data collection are literature study, interview, documentation and observation.

Data analysis method. Regression analysis is a statistical analysis that studies the relationship between two or more variables. In the linear regression analysis it is assumed that the this relationship is a linear combination, the simplest case consisting of one independent variable (Pangestu 2016).

Linear regression is a statistical method used to model the relationship between the dependent variable (dependent; response; Y) and one or more independent variables (independent, predictor, X). If there is only one independent variable, it is called simple linear regression, whereas if there is more than one independent variable, it is called multiple linear regression. Regression analysis has at least three use cases, namely for data description, for control and for predictive purposes.

Simple regression is carried out in stages, namely by connecting the independent variable (chlorophyll-a) with the dependent variable (tuna catch, using the data regression analysis function of Microsoft Office Excel (Kurniawan 2008). This simple regression analysis was conducted to determine the relationship between the dependent variable Y (*K. pelamis* catches) and the independent variable X (chlorophyll-a). The regression model used is:

$$Y = a + bX$$

Where:

X - chlorophyll-a;

Y - pole and line catch;

a - the constant (intercept) is the value of Y, when X = 0 which equals the value of a where it intersects the Y axis;

b - regression coefficient.

Chlorophyll-a image analysis. The fishing ground forecast can be predicted from the parameters of SST and chlorophyll-a. The analysis of the distribution of SST and chlorophyll-a can use level 1b and level 2 Modis images. The Algorithm Theoretical Basic Document Modis (ATBD) 25 is used to determine SST and the Morel 4 Algorithm is used

to determine the chlorophyll-a. The Modis image levels 1b and 2 use the RMSE test calculation (Chavula 2012).

From the two Root Mean Squared Error (RMSE) tests carried out between the field data and the Modis image data levels 1b and level 2, more accurate values were obtained from the smallest RMSE sum value, namely the Level 2 Modis image, with an RMSE value of chlorophyll-a of 0.090315. So the best fishing ground forecast can be determined through Level 2 Modis Images (Febriani & Sukojo 2016).

Results and Discussion

Fishing ground. In this study we used a pole and line fishing boat (Figure 2) that operates in Savu sea, Flores, East Nusa Tenggara.



Figure 2. Pole and line fishing boat.

Composition and comparison of catches with the number of settings.

1. In November 2019 the ship made one trip with a catch of *K. pelamis* of 1,020 kg.
2. In December 2019 the ship made 16 trips with a catch of 28,220 kg. The fishing season generally reaches a turning point in the middle of January and in December, according to the graph, the catch will decrease as the wind season arrives.
3. In January 2020 the catch obtained was of 7,990 kg. January is generally the start of the wind season, so ships rarely go to sea.
4. In February 2020 the catches were dominated by *K. pelamis* and *T. albacares*. According to the fishermen, February was generally the peak of the wind season, so that the catch was not good, until the middle or end of March.
5. In March 2020 the catches were dominated by *K. pelamis* and *T. albacares*, but unlike in the previous month, *C. hippurus* and *E. bipinnulata* were also obtained. In March 2020 the fishermen began to increase the intensity of their trips, but taking into account the strength of the wind and the availability of bait.

The composition of the catch for 5 months can be seen in Table 1 below:

Table 1

Comparison of catch volumes with the number of settings

Volumes / Period	November	December	January	February	March
Number of setting (times)	2	43	14	15	13
<i>K. pelamis</i> (kg)	1,020	15,300	2,890	6,290	7,480
<i>T. albacares</i> (kg)	0	9,520	2,040	5,440	4,250
<i>C. hippurus</i> (kg)	0	1,870	1,870	0	1,360
<i>E. bipinnulata</i> (kg)	0	1,530	1,190	0	1,870
Total (kg)	1,020	28,220	7,990	11,730	14,960
Catch rate per setting	510	656.28	570.71	782	1,150

(kg)

Catch frequencies can also be visualized in Figure 3, below:




Figure 3. Total catch (November 2019 to March 2020).

Based on the graph above, there is a decrease in the catch volumes during the 4 months of fishing. The lowest catch was obtained in January with a total of 7,990 kg, consisting of: *K. pelamis* (2,890 kg), *T. albacares* (2,040 kg) and others, such as *E. bipinnulata* and *C. hippurus*. Meanwhile, the highest catch was obtained in December is 28,220 kg, dominated by *K. pelamis* with 15,300 kg and *T. albacares* with 9,520 kg. In 2016, Kupang Regency was the largest supplier of *K. pelamis* in East Nusa Tenggara (MMAF 2018).

Catch composition. If we see in Table 2 and figure 4 below, it is shown that *K. pelamis* is the most dominant catch that lands at Tenau port and at the fishing port around Kupang city, East Nusa Tenggara. This is in accordance with the data from the Fisheries Service of the Province of East Nusa Tenggara, suggesting that *K. pelamis* is one of the main commodities produced in Kupang, with 52%, followed by *T. albacares*, with 33%, *C. hippurus*, with 7% and *E. bipinnulata*, with 8%.

Table 2

Catch rated per setting

Name of fish	Amount (kg)	Catch percentage (%)
<i>K. pelamis</i>	32,980	52
<i>T. albacares</i>	21,250	33
<i>C. hippurus</i>	4,590	7
<i>E. bipinnulata</i>	5,100	8
Total catching	63,920	100
Catch rate per setting	734.71	1.14

The catch compositions (in %) can be seen in Figure 4, below.

Commented [WU8]: Please reformulate, using scientific English.

Commented [A9R8]: Corrected. Thank you




Figure 4. Catch composition diagram.

Conformity between of the fishing ground and PPDPI. Fishing is carried out on FADs that have been installed, with the position of the FADs stored on the GPS of each ship. The following is a map of the fishing grounds with the positions of the FADs. Figure 5 below shows the fishing ground for 5 months of fishing:




Figure 5. Fishing ground.

Forecast map of fishing ground (PPDPI).

1. In November the PPDPI that can be used correspond to 29 to 30 November 2019. On this map, the potential fishing ground are around the northwest of Timor Island with an area of 150 NM.
2. In December, 15 files were issued with an interval of 2 days. At PDPPI 11 to 12 December 2019, the recommended fishing areas are south of Savu Island, northeast of Sumba Island and north of Rote Island, as well as many other areas with an area of more than 300 NM. In the PDPPI 11 to 12 December 2019, the fishing areas that recommended are in the south of Savu Island, northeast of Sumba Island and north of Rote Island, as well as many other areas with an area of more than 300 NM (IMRO 2019).
3. In January, only 1 file out of 7 uploaded PPDPI files issued corresponded to the time of the fishing trip. On the 31 December to 01 January map, the potential fishing grounds were located around the southeast of Sumba Island, west of Savu Island, with an area of more than 300 NM (IMRO 2020).

Commented [A10]: ? area unit, please.

Commented [A11R10]: Ya, NM or nautical mile is a unit of length used in air, marine, and space navigation, and for the definition of territorial waters.

Commented [A12]: Please reformulate

Commented [A13]: Please correct the unit

4. In February, as many as 10 PDPPI files were uploaded and 3 files corresponded to the time of the fishing trip. One of the suitable map files was issued on the 26 to 27 February recommending fishing zones located north of Sumba Island with an area of 300 NM. Other areas do not seem to be potential fishing grounds (IMRO 2020).
5. In March there were 4 files matching, at the time of the fishing trip. March should be the turning point of the monsoon season that occurs in East Nusa Tenggara, but each region has a different time. One of the maps is the 16 to 17 March map, the areas that are recommended for fishing activities are in the south of Sumba Island and northeast of Sumba Island, with an area of more than 300 NM. There was also one point with an area of 50 NM located on Semau and Kupang islands (IMRO 2020).

Commented [A14]: Please correct the surface unit




Figure 6. PDPPI period in November (1), December (2) (IMRO 2019); January (3), February (4), March (5) (IMRO 2020).

Compatibility between the fishing ground and PDPPI. Of all trips with available PDPPI, there were only 17 files that matched, at the time of the fishing trip. For the 4 months period, 17 maps of fishing ground forecasts were used, with a total of 44 settings (Table 3).

The method used to determine the suitability of fishing grounds with PDPPI, issued by IMRO, uses a comparison method between the position of the actual fishing area and the status of the same location on the forecast map, on that date. From 44 setting positions, 6 positions were in accordance with the PDPPI, which corresponds to a match rate of 13.6%, in terms of chlorophyll-a and SST. When the daily PDPPI images from the NASA website are not available, the fishing trips cannot be used as a reference in determining their suitability.

With a match rate of 13.6%, it can be said that fishermen using FADs have sufficient accuracy in determining fishing grounds

Table 3

Compatibility of the fishing ground with the PPDPI

No	Setting	Position		Chlorophyll-a (mg m ⁻³)	Information
		Latitude	Longitude		
1	1	10° 12' 27" S	121° 40' 16" E	0.41	Inappropriate
	2	10° 09' 43" S	121° 37' 40" E	0.31	Inappropriate
2	1	09° 56' 38" S	122° 25' 03" E	0.3	Appropriate
	2	10° 42' 09" S	121° 50' 28" E	0.27	Inappropriate
3	3	10° 26' 28" S	122° 47' 58" E	0.27	Appropriate
	1	09° 32' 11" S	123° 20' 42" E	0.24	Inappropriate
4	2	09° 45' 37" S	123° 13' 20" E	0.22	Inappropriate
	1	10° 04' 18" S	122° 32' 21" E	0.27	Appropriate
5	2	10° 28' 31" S	122° 41' 16" E	0.31	Inappropriate
	3	10° 12' 31" S	122° 05' 10" E	0.33	Appropriate
6	1	10° 40' 10" S	122° 15' 30" E	0.29	Inappropriate
	2	09° 40' 33" S	122° 55' 15" E	0.39	Inappropriate
7	1	09° 31' 40" S	123° 05' 09" E	0.21	Inappropriate
	2	09° 05' 45" S	123° 00' 41" E	0.28	Inappropriate
8	3	09° 47' 16" S	122° 58' 34" E	0.13	Inappropriate
	1	09° 40' 08" S	123° 22' 31" E	0.29	Inappropriate
9	2	09° 50' 28" S	122° 53' 17" E	0.30	Inappropriate
	1	10° 38' 15" S	121° 40' 30" E	0.41	Inappropriate
10	2	10° 42' 09" S	121° 50' 28" E	0.38	Inappropriate
	3	10° 49' 25" S	121° 01' 51" E	-	Inappropriate
11	1	09° 32' 11" S	123° 20' 42" E	0.28	Inappropriate
	2	09° 45' 37" S	123° 13' 20" E	0.24	Inappropriate
12	3	09° 54' 26" S	123° 01' 51" E	-	Inappropriate
	1	10° 21' 15" S	122° 35' 07" E	0.45	Inappropriate
13	2	10° 10' 35" S	122° 05' 37" E	-	Inappropriate
	3	10° 03' 47" S	122° 45' 10" E	0.54	Appropriate
14	1	09° 40' 36" S	122° 05' 15" E	0.8	Inappropriate
	2	09° 50' 17" S	121° 35' 20" E	1.1	Inappropriate
15	1	09° 30' 14" S	122° 50' 33" E	-	Inappropriate
	2	09° 21' 52" S	123° 00' 16" E	-	Inappropriate
16	1	10° 40' 16" S	123° 41' 40" E	-	Inappropriate
	2	10° 39' 13" S	123° 45' 36" E	-	Inappropriate
17	3	10° 40' 41" S	123° 40' 22" E	-	Inappropriate
	1	09° 50' 20" S	122° 10' 31" E	-	Inappropriate
18	2	09° 18' 50" S	122° 46' 15" E	-	Inappropriate
	3	09° 30' 16" S	123° 01' 51" E	-	Inappropriate
19	1	10° 03' 15" S	122° 45' 17" E	0.31	Inappropriate
	2	10° 03' 40" S	121° 48' 29" E	-	Inappropriate
20	1	10° 08' 36" S	123° 07' 12" E	-	Inappropriate
	2	10° 15' 53" S	122° 46' 41" E	-	Inappropriate
21	3	10° 03' 09" S	122° 53' 08" E	-	Inappropriate
	1	09° 29' 21" S	122° 30' 14" E	0.6	Inappropriate
22	2	09° 50' 15" S	122° 30' 35" E	0.54	Inappropriate
	3	09° 55' 23" S	121° 20' 54" E	0.49	Appropriate

Concentration and distribution of chlorophyll-a. Chlorophyll-a concentrations can be obtained by downloading data levels 1, 2 and 3 on the NASA website. Data needs in a longer time span such as months and years can use level 3 image data with better quality than others, but level 3 images only contain one specific information depending on what is needed. In this study, the basis used is a level 3 image which only requires chlorophyll-a information with a time span of months. In Figure 7 and Table 4 the following is image data for chlorophyll-a concentration level 3.

Chlorophyll-a concentrations can be obtained by downloading data levels 1, 2 and 3 from the NASA website. Data needs in a longer time span such as months and years can use level 3 image data with better quality than the others, but only containing one specific information, depending on what is needed. In this study, the basis used is a level 3 image which only requires chlorophyll-a information with a time span of months. Figure 7 images and Table 4 data correspond to level 3 chlorophyll-a concentration.




Figure 7. Concentration of chlorophyll-a map (www.oceancolour.gsfc.nasa.gov).

The distribution map of chlorophyll-a concentrations (Table 4) from November to March, shows elevated values around the west of Flores Island, Sumba Island and also Larantuka, suggesting that food chain occur in the area.

Table 4

Concentration and distribution of chlorophyll-a

<i>Period</i>	<i>November</i>	<i>December</i>	<i>January</i>	<i>February</i>	<i>March</i>
Concentration of chlorophyll-a (mg m ⁻³)					
a. Lowest	0.07	0.08	0.08	0.08	0.08
b. Highest	1.29	1.14	1.15	1.19	1.17
Average Value (mg m ⁻³)	0.2	0.3	0.2	0.2	0.2
Distribution center	North of Kupang city, Sabu Island, and west of Flores	West of Flores	Kupang city, Sabu Island, Rote Island, and west of Flores	Kupang city, Rote Island, and west of Flores	Kupang city, west of Flores
Compatibility level	Appropriate	Inappropriate	Appropriate	Appropriate	Inappropriate

Conclusions. Based on the results of the study it can be concluded: simultaneously, chlorophyll-a had an effect of 62.2% on the catch. The highest concentration of chlorophyll-a recorded was 1.29 mg m⁻³ and the lowest was 0.07 mg m⁻³; the fishing area used by fishermen has a level of conformity with the forecast map for fishing areas of 13.6%; Not all fishing trips can be used as a reference in determining the suitability of fishing grounds with PDPPI, because not every day PPDPI can be uploaded because it is limited by the availability of images obtained from the NASA website.

Acknowledgments. The authors would like to thank to Director and staff of Institute for Marine Research and Observation (IMRO), Agency for Marine and Fisheries Research and Development, Ministry of Marine Affairs and Fisheries, Indonesia. for the support to sharing and processing of the satellite data related to the distribution of SST and Chlorophyll-a in fishing ground and to all other parties for their help and contributions.

Conflict of interest. The authors declare no conflict of interest.

References

- Asagabaldan M. A., 2017 Law and the role of Lilifuk Amid Government Regimes: Legal analysis and sustainability of local wisdom in Kuanheun Village, West Kupang Regency, East Nusa Tenggara. *SABDA* 12(2):136-145.
- Behrenfeld M. J., Boss E., Siegel D. A., Shea D. M., 2005 Carbon-based ocean productivity and phytoplankton physiology from space. *Global Biogeochemical Cycles* 19(1):1-14.
- Cayula J. F., Cornillon P., 1992 Edge detection algorithm for SST Images. *Journal of Atmospheric and Oceanic Technology* 9(1):67-80.
- Chavula G., Sungani H., Gondwe K., 2012 Mapping potential fishing grounds in Lake Malawi using AVHRR and MODIS satellite imagery. *International Journal of Geosciences* 3(03):650-658.
- Darsono P., 1999 [Utilization of marine resources and its implications for fishing communities]. *Oceana* 24(4):1-9. [In Indonesian].

- Farda N. M., Jatisworo D., 2019 [Mining space-time patterns on forecast maps of fishing areas in fisheries management areas (FMA) 712, 713, and 573]. Scientific Magazine GLOBE 21(2):117-128. [In Indonesian].
- Febriani E. R., Sukojo B. M., 2016 [Comparative analysis of the use of level 1b and level 2 MODIS imagery in determining forecasting fishing areas (Case Study: South Coast of Blitar)]. ITS Engineering Journal, Surabaya 5(2):439-442. [In Indonesian].
- Hakim D. M., Wikantika K., Widiadnyana N., Napitu A. M., Darmawan S., 2006 The identification of fishing ground area with MODIS satellite image (Case Study: South Coast of West Java) Proceeding. ITB Engineering Science 38B(2):147-158.
- Hasyim S., Zainuddin M., 2008 Predictions of skipjack fishing areas based on oceanographic conditions in the waters of Takalar regency and surrounding areas. Journal of Science and Technology 8(2):158-162. [In Indonesian].
- Huot Y., Babin M., Bruyant F., Grob C., Twardowski M. S., Claustre H., 2007 Does chlorophyll-a provide the best index of phytoplankton biomass for primary productivity studies? Biogeosciences Discussions, Proxies of Biomass For Primary production 4(2):707-745.
- Jatisworo D., Murdimanto A., 2013 [Identification of thermal fronts in the Makassar Strait and Banda Sea]. Proceedings of the National Symposium on Geoinformation Science III. Marine Research and Observation Institute, 8 p. [In Indonesian].
- Kunarso, Zainuri M., Sugianto D. N., Marwoto J., Hariyadi, Muslim, 2019 Prediction of demersal fishing ground associated with coral reefs in the Coastal Jepara Regency, Central Java, Indonesia based on Sentinel 2a imagery. Advances in Science, Technology and Engineering Systems (ASTES) Journal 4(6):263-269.
- Kurniawan D., 2008 [Linear regression]. Academia accelerating the world's research. Research Development Core Team R: A language and environment for statistical computing, 14 p. [In Indonesian].
- Kusumastanto T., 1996 [An evaluation of government policies on the development of fisheries in Indonesia]. AGRIS, Fisheries Economy Bulletin 2(3):27-41. [In Indonesian].
- Latif B., Noor S. M., Sumardi J., Irwansyah, 2019 Sustainable development and sea protection: Trade on fish and fishery product. ICROEST. IOP Conference Series: Earth and Environmental Science 343(1):1-6.
- Laurs R. M., Fiedler P. C., Montgomery D. R., 1984 Albacore tuna catch distribution relative to environmental features observed from satellites. Deep-Sea Research 31:1085-1099.
- Lee Y. J., Matrai P. A., Friedrichs M. A. M., Saba V. S., Aumont O., Babin M., Buitenhuis E. T., Chevallier M., Mora L., Dessert D., Dunne J. P., Ellingsen I. H., Feldman D., Frouin R., Gehlen M., Gorgues T., Ilyina T., Jin M., John J. G., Lawrence J., Manizza M., Menkes C. E., Perruche C., Fouest V. L., Popova E. E., Romanou A., Samuelsen A., Schwinger J., Séférian R., Stock C. A., Tjiputra J., Tremblay L. B., Ueyoshi K., Vichi M., Yool A., Zhang J., 2016 Net primary productivity estimates and environmental variables in the Arctic Ocean: An assessment of coupled physical-biogeochemical models. Journal of Geophysical Research: Oceans 121(12):8635-8669.
- Nainggolan C., 2012 [Fishing methods]. Open University, Jakarta, 61 p. [In Indonesian].
- Nugraha E., Yudho G. S., Jaenudin A., Yusrizal, Kusmedy B., Kusnidar A., Husen E. S., 2020a Relationship between length and weight of skipjack tuna (*Katsuwonus pelamis*) purse seine catching in the Maluku Sea, Indonesia. AACL Bioflux 13(1):330-345.
- Nugraha E., Gunawan R., Danapraja S., Yusrizal, Kusdinar A., Waluyo A. S., Hutajulu J., Prayitno H., Halim S., Sutisna D. H., 2020b The sea surface temperature effect on the length and size of skipjack tuna (*Katsuwonus pelamis*) catches in the Banda Sea, Indonesia. AACL Bioflux 13(1):1-18.
- Nurman A., 2010 [Utilization of fashionable data to detect fishing grounds on the east and west coasts of North Sumatra]. Medan State University. Geography Journal 2(2):17-30. [In Indonesian].

- Nusantara R. A., Rosyid A., Boesono H., 2014 Analysis of differences in depth of fishing area towards fish composition on Danish seine in Rembang waters. *Journal of Fisheries Resources Utilization Management and Technology* 3(3):96-104 [In Indonesian].
- Nuzapril M., Susilo S. B., Panjaitan J. P., 2017 Relationship between chlorophyll-a concentration with primary productivity rate using Landsat 8 imagery. *Journal of Fisheries and Marine Technology* 8(1):105-114.
- Pangestu S., 2016 [Simple linear regression. Applied linear model]. Digital Library of a Open University of Indonesia, 398 p. [In Indonesian].
- Sato K., Matsuno K., Arima D., Abe Y., Yamaguchi A., 2015 Spatial and temporal changes in zooplankton abundance, biovolume, and size spectra in the neighboring waters of Japan: Analyses using an optical plankton counter. *Zoological Studies. A Springer Open Journal* 54(18):1-15.
- Seki M. P., Polovina J. J., Kobayashi D. R., Bidigare R. R., Mitchum G. T., 2002 An oceanographic characterization of swordfish (*Xiphias gladius*) longline fishing grounds in the springtime subtropical North Pacific. *Fisheries Oceanography* 11(5):251-266.
- Sudirman, Mallawa A., 2012 [Fishing techniques]. Rineka Cipta, Jakarta, Indonesia, 168 p. [In Indonesian].
- Sugiari, Anshori M. A., Kusumawardani M., 2015 [Coordinate information system for estimated tuna fishing grounds for fishermen]. *Journal of JARTEL State Polytechnic of Malang* 1(1):74-84. [In Indonesian].
- Suniada K. I., Susilo E., Hastuti A. W., 2015 [Validation of forecast map of fishing grounds (PPDPI) in Java Sea Waters (FMA-RI 712)]. *Proceedings of the National Marine and Fisheries Science and Technology Forum* 87-96. [In Indonesian].
- Tambun R., Simbolon D., Wahyu R., Supartono, 2018 [Potential zones of fishing ground based on season at FMA RI 718]. *Journal of Tropical Marine Science and Technology* 10(3):757-768. [In Indonesian].
- Tuli M., Mennofatria B., Adrianto L., 2015 [Analysis of skipjack fish (*Katsuwonus pelamis*) resources in the waters of Pohuwato Regency, Gorontalo Province]. *Marine Fisheries Journal* 6(2):109-117. [In Indonesian].
- Wibawa T. A., 2011 [Utilization of oceanographic satellite data for prediction of potential fishing grounds for bigeye tuna (*T. obesus*) in the South Indian Ocean, Java and Bali]. *SEGARA Journal* 7(1):29-41. [In Indonesian].
- Zainuddin M., 2011 Skipjack tuna in relation to sea surface temperature and chlorophyll-a concentration of Bone Bay using remotely sensed satellite data. *Journal of Tropical Marine Science and Technology* 3(1):82-90.
- Zainuddin M., Safruddin, Farhum S. A., Nelwan A., Selamat M. B., Hidayat S., Sudirman, 2015 [Characteristics of potential areas for catching skipjack tuna in the Bone-Sea Bay of Flores based on satellite data on sea surface temperature and chlorophyll-a in the period January-June 2014 Makassar]. *Journal of Science and Technology Utilization of fisheries resources. Hasanuddin University* 2(3):228-237. [In Indonesian].
- Zhang C., Han M., 2015 Mapping chlorophyll-a concentration in Laizhou Bay using Landsat 8 OLI data. e-proceedings of the 36th IAHR World Congress, The Hague, The Netherlands, pp. 12-17.
- *** Institute for Marine Research and Observation, IMRO, 2019 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian],
- *** Institute for Marine Research and Observation, IMRO, 2020 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2016 [East Nusa Tenggara Provincial annual report 2016]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2018 [East Nusa Tenggara Provincial annual report 2018]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** <http://modis.gsfc.nasa.gov> 2007

Commented [WU15]: Missing from the text

Commented [A16R15]: There is. Komang, Change with Suniada

*** www.oceancolour.gfsc.nasa.gov

Received: 2021. Accepted: 2021. Published online: 2021.

Authors:

Maman Hermawan, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: mhermawan60@gmail.com

I Nyoman Radiarta, Institute for Marine Research and Observation, Agency for Marine and Fisheries Research and Development Baru Street, Perancak, Negara, Jembrana 82251 Bali, Indonesia, e-mail: radiarta@kkp.go.id

I Gusti Bagus Ega Krisna Bayu, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: egabayu91@gmail.com

Erick Nugraha, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: nugraha_eriq1@yahoo.co.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Hermawan M., Radiarta I. N., Bayu I. G. B. E. K., Nugraha E., 2021 ~~The suitability of the fishing ground with the forecast map of the fishing ground (PPDPI) in terms of chlorophyll-a in the Savu Sea, Indonesia.~~ [Comparison between fisherman fishing ground and the forecast map of fishing ground \(PPDPI\) based on chlorophyll-a in the Savu Sea, Indonesia.](#) AACL Bioflux 14(x):

← Kembali ↶ ↷ →

📁 Arsipkan 📁 Pindahkan 🗑️ Hapus 🛡️ Spam ⋮

☰ ▲ ▼ ✕

E • **Eni Kovacs** Dear Dr.Nugraha, Have you received the our previous e-mail regarding the formatting of your manuscript? "As mentioned in the prelimi" Sen, 21 Jun 2021 jam 14.17 ☆

EN • **erick nugraha** <nugraha_eriq1@yahoo.co.id> Kepada: Eni Kovacs 📄 🔗 Sen, 21 Jun 2021 jam 21.55 ☆

Dear Dr. Eni


I send you corrected of manuscript journal.
thank you.

> Tampilkan pesan asli



The suitabi....docx
13MB

↶ ↷ → ⋮



Identifying fishing grounds in the Savu Sea, Indonesia: fishermen's experience compared with the chlorophyll-a forecast maps

¹Maman Hermawan, ²I Nyoman Radiarta, ¹I Gusti B. E. K. Bayu, ¹Erick Nugraha

¹ Faculty of Fishing Technology, Jakarta Technical University of Fisheries, Pasarminggu, South Jakarta, Indonesia; ² Institute for Marine Research and Observation, Negara, Jembrana, Bali, Indonesia. Corresponding author: E. Nugraha, nugraha_eriq1@yahoo.co.id

Abstract. The Savu Sea in East Nusa Tenggara Province is the location of fishing activities for most fishermen in East Nusa Tenggara. Despite its potential, the knowledge of the fishing ground is still conventional. Currently, predictive methods for trips planning are available, such as the fishing ground forecast maps based on the chlorophyll-a, sea surface temperature and salinity. The related data are provided by NASA, being processed by the Marine Research and Observation Center to display the potential fishing ground as a box view with an accuracy level of 5 square miles. Eventually, the forecast map of the fishing area will be compared with the fishing locations known by the fishermen, for validation. The data collection used a survey method and the data analysis used both a descriptive analysis and the linear regression. The forecast maps matched with the observed fishing locations at only 13.6%. The highest recorded chlorophyll-a concentration was of 1.29 mg m⁻³ (constantly observed in the West Flores) and the lowest was of 0.07 mg m⁻³. The chlorophyll-a has an effect on 62.2% of the catch and 52% of the catch is composed of *Katsuwonus pelamis*

Key Words: MODIS, sea surface temperature, catch composition, *Katsuwonus pelamis*.

Introduction. Indonesia is the largest archipelagic country in the world, having a coastline length of 81,000 km² and a sea area of about 3,100,000 km² (Kusumastanto 1996; Darsono 1999; Latif et al 2019). Managing the fisheries resource requires determining the fish growth patterns and the catch period, in order to avoid over-exploitation (Nugraha et al 2020b). The province of East Nusa Tenggara has a sea area of 200,000 km² with a coastline of 5,700 km² (MMAF 2016). Savu Sea has a good potential for fisheries and the whole marine sector (Asagabaldan 2017). A fishing area is an area where fish are caught in maximum amount and fishing gear can be operated economically. Assessment and mapping of fishing areas are needed (Nusantara et al 2014). The *Katsuwonus pelamis* is a fast swimming fish and has the greedy character of an opportunistic predator, with food types ranging from small fish (Clupeidae and Engraulidae), squids, crustaceans and zooplankton (Tuli et al 2015). *K. pelamis* is a pelagic fish living in groups, from the coastal areas to the open sea (Nainggolan 2012). Schooling fish should be kept as close to the boat as possible so that they can be easily caught using pole and line fishing rods (Sudirman & Mallawa 2012). The fishing grounds are determined by the waters' condition in the habitat of a species. Water conditions are usually described by oceanographic parameters (Nugraha et al 2020a).

The Sea Surface Temperature (SST) of the *K. pelamis* fishing ground ranges from 29.75 to 30.25°C (Zainuddin et al 2015; Nugraha et al 2020b; Zainuddin 2011). Meanwhile, the potential area has a characteristic chlorophyll-a concentration of around 0.125 mg m⁻³. This condition is mainly seen in May. Outside of this range, the catch of *K. pelamis* tends to decline, as it happens in April (when the SST values in the Bone-Sea Flores bay are the highest). Likewise, when the chlorophyll-a concentration in the fishing ground significantly increase and the catch tends to decrease. This means that there is a

certain range of values for the two oceanographic factors that are very suitable for the presence and abundance of *K. pelamis*. The distribution of chlorophyll-a can be a parameter used for identifying the presence of fish in a water area (Hasyim & Zainuddin 2008; Laurs et al 1984).

The Forecast Maps of Fishing Ground (PPDPI) is the result of a research conducted by the Institute for Marine Research and Observation (IMRO) in order to identify fishing grounds, based on an internet service of information on the fishing areas' characteristics (Sugiari 2015; Panjaitan 2014). PPDPI have been produced for and distributed to the fishing communities of Indonesia, since 2000 (Tambun et al 2018; Farda & Jatisworo 2019). PPDPI was compiled using remote sensing data which is also widely used to detect water conditions such as the thermal front (Cayula & Cornillon 1992; Wibawa 2011; Jatisworo & Murdimanto 2013) and the concentration of chlorophyll-a (Seki et al 2002) which is one of the indicators of water fertility and school of fish. PPDPI preparation was carried out using SST and water fertility data (chlorophyll-a) generated from MODIS Aqua level 3 daily composites, in the Hierarchical Data Format (HDF) (Suniada et al 2015).

The predictive maps of fishing grounds that already exists in Indonesia still have many weaknesses, including the spatial aspect of the points of forecasting located far from the coast and the uncertainty of the predicted type of fish. Thus, if inappropriate gear is selected for the fishing trip, the catch productivity is negatively affected, even in presence of abundant fish populations (Kunarso et al 2019).

Chlorophyll-a is an indicator of the abundance of phytoplankton in waters, playing a role in the photosynthesis process (Huot et al 2007; Zhang & Han 2015). Phytoplankton contributes to a large extent to determine primary productivity in waters. Organic carbon production during photosynthesis is defined as primary productivity or Net Primary Productivity (Lee et al 2016).

Linear regression analysis was carried out on the model of the relationship between chlorophyll-a concentration and primary productivity. The results of this equation can be applied to the satellite imagery so that it can help to monitor the water quality conditions (Nuzapril et al 2017). This abundance of phytoplankton will result in the creation of food chain clusters from phytoplankton to large pelagic fish schools (Sato et al 2015; Behrenfeld et al 2005).




Figure 1. Map of chlorophyll-a concentration (IMRO 2020).

Remote sensing data can show the variables used in determining fish catchment areas. Certain remote sensing images used to extract SST and chlorophyll-a information are provided by the MODIS program (Nurman 2010; Hakim et al 2006).

Satellite imagery is one of the applications of the remote sensing technology. The Moderate-resolution Imaging Spectroradiometer (MODIS) technology uses sensors belonging to the Earth Observing System (EOS), carried by two satellites produced by the National Aeronautics and Space Administration (NASA), namely Terra and Aqua (<http://modis.gsfc.nasa.gov> 2007).

The MODIS sensor is a derivative of the Advanced Very High Resolution Radiometer (AVHRR) sensor, Sea-viewing Wide Field-of-View Sensor (SeaWiFS) and High Resolution Imaging Spectrometer (HRS) owned by EOS, which has previously orbited, so it can be used to measure sea and atmosphere-level parameters (such as the sea surface temperature, chlorophyll-a concentration and water vapor content) and marine phenomena (such as the occurrence of upwelling, thermal fronts and others) (<http://modis.gsfc.nasa.gov> 2007).

Material and Method

Description of the study sites. This research was conducted on a pole and line fishing boat operating in the Flores Sea, East Nusa Tenggara. The satellite image data retrieval was carried out at the Southeast Research and Ocean Observation Center (SeaCORM) Jembrana, Bali. The material used consisted of sea water samples for measuring oceanographic elements. The methods used in data collection are literature study, interview, documentation and observation.

Data analysis method. Regression analysis is a statistical analysis that studies the relationship between two or more variables. In the linear regression analysis it is assumed that this relationship is a linear combination, the simplest case consisting of one independent variable (Pangestu 2016).

Linear regression is a statistical method used to model the relationship between the dependent variable (dependent; response; Y) and one or more independent variables (independent, predictor, X). If there is only one independent variable, it is called simple linear regression, whereas if there is more than one independent variable, it is called multiple linear regression. Regression analysis has at least three use cases, namely for data description, for control and for predictive purposes..

Simple regression is carried out in stages, namely by connecting the independent variable (chlorophyll-a) with the dependent variable (tuna catch), using the data regression analysis function of Microsoft Office Excel (Kurniawan 2008). This simple regression analysis was conducted to determine the relationship between the dependent variable Y (*K. pelamis* catches) and the independent variable X (chlorophyll-a). The regression model used is:

$$Y = a + bX$$

Where:

X - chlorophyll-a;

Y - pole and line catch;

a - the constant (intercept) is the value of Y, when X = 0 which equals the value of a where it intersects the Y axis;

b - regression coefficient.

Chlorophyll-a image analysis. The fishing ground forecast can be predicted from the parameters of sea surface temperature and chlorophyll-a. The analysis of the distribution of sea surface temperature and chlorophyll-a can use level 1b and level 2 Modis images. The Algorithm Theoretical Basic Document Modis (ATBD) 25 is used to determine sea surface temperature and the Morel 4 Algorithm is used to determine the chlorophyll-a. The Modis image levels 1b and 2 use the RMSE test calculation (Chavula 2012).

From the two Root Mean Squared Error (RMSE) tests carried out between the field data and the Modis image data levels 1b and level 2, more accurate values were obtained from the smallest RMSE sum value, namely the Level 2 Modis image, with an RMSE value of chlorophyll-a of 0.090315. So the best fishing ground forecast can be determined through Level 2 Modis Images (Febriani & Sukojo 2016).

Results and Discussion

Fishing ground. In this study we used a pole and line fishing boat (Figure 2) that operates in Savu sea, Flores, East Nusa Tenggara.



Figure 2. Pole and line fishing boat (original).

Composition and comparison of catches with the number of settings.

1. In November 2019 the ship made one trip with a catch of *K. pelamis* of 1,020 kg.
2. In December 2019 the ship made 16 trips with a catch of 28,220 kg. The fishing season generally reaches a turning point in the middle of January and in December, according to the graph, the catch will decrease as the wind season arrives.
3. In January 2020 the catch obtained was of 7,990 kg. January is generally the start of the wind season, so ships rarely go to sea.
4. In February 2020 the catches were dominated by *K. pelamis* and *T. albacares*. According to the fishermen, February was generally the peak of the wind season, so that the catch was not good, until the middle or end of March.
5. In March 2020 the catches were dominated by *K. pelamis* and *T. albacares*, but unlike in the previous month, *C. hippurus* and *E. bipinnulata* were also obtained. In March 2020 the fishermen began to increase the intensity of their trips, but taking into account the strength of the wind and the availability of bait.

The composition of the catch for 5 months can be seen in Table 1.

Table 1

Comparison of catch volumes with the number of settings

Volumes / Period	November	December	January	February	March
Number of setting (times)	2	43	14	15	13
<i>K. pelamis</i> (kg)	1,020	15,300	2,890	6,290	7,480
<i>T. albacares</i> (kg)	0	9,520	2,040	5,440	4,250
<i>C. hippurus</i> (kg)	0	1,870	1,870	0	1,360
<i>E. bipinnulata</i> (kg)	0	1,530	1,190	0	1,870
Total (kg)	1,020	28,220	7,990	11,730	14,960
Catch rate per setting (kg)	510	656,28	570,71	782	1,150

Catch frequencies can also be visualized in Figure 3.




Figure 3. Total catch (November 2019 to March 2020).

Based on the graph above, there is a decrease in the catch volumes during the 4 months of fishing. The lowest catch was obtained in January with a total of 7,990 kg, consisting of: *K. pelamis* (2,890 kg), *T. albacares* (2,040 kg) and others, such as *E. bipinnulata* and *C. hippurus*. The highest catch was obtained in December is 28,220 kg, dominated by *K. pelamis* with 15,300 kg and *T. albacares* with 9,520 kg. In 2016, Kupang Regency was the largest supplier of *K. pelamis* in East Nusa Tenggara (MMAF 2018).

Catch composition. Table 2 and Figure 4 show that *K. pelamis* is the most dominant catch that lands at Tenau port and at the fishing port around Kupang city, East Nusa Tenggara. This is in accordance with the data from the Fisheries Service of the Province of East Nusa Tenggara, suggesting that *K. pelamis* is one of the main commodities produced in Kupang, with 52%, followed by *T. albacares*, with 33%, *C. hippurus*, with 7% and *E. bipinnulata*, with 8%. The catch compositions (in %) can be seen in Figure 4.

Table 2

Catch rated per setting

Name of fish	Amount (kg)	Catch percentage (%)
<i>K. pelamis</i>	32,980	52
<i>T. albacares</i>	21,250	33
<i>C. hippurus</i>	4,590	7
<i>E. bipinnulata</i>	5,100	8
Total catch	63,920	100
Catch rate per setting	734.71	1.14




Figure 4. Catch composition diagram.

Conformity between of the fishing ground and PPDPI. Fishing is carried out on FADs that have been installed, with the position of the FADs stored on the GPS of each

Commented [WU1]: Please use the full name for all the species. Please use English: March

ship. The following is a map of the fishing grounds with the positions of the FADs. Figure 5 below shows the fishing ground for 5 months of fishing.




Figure 5. Fishing ground.

Commented [WU2]: December, not Desember

Forecast map of fishing ground (PPDPI).




Figure 6. PPDPI period in November (1), December (2) (IMRO 2019); January (3), February (4), March (5) (IMRO 2020).

1. In November the PPDPI that can be used correspond to 29 to 30 November 2019. On this map, the potential fishing grounds are around the northwest of Timor Island with an area of 150 NM.
2. In December, 15 files were issued with an interval of 2 days. In the PDPPI 11 to 12 December 2019, the fishing areas that recommended are in the south of Sawu Island, northeast of Sumba Island and north of Rote Island, as well as many other areas with more than 300 NM (IMRO 2019).
3. In January, only 1 file out of 7 uploaded PPDPI files issued corresponded to the time of the fishing trip. On the 31 December to 01 January map, the potential fishing grounds were located around the southeast of Sumba Island, west of Savu Island, with an area of more than 300 NM (IMRO 2020).
4. In February, as many as 10 PPDPI files were uploaded and 3 files corresponded to the time of the fishing trip. One of the suitable map files was issued on the 26 to 27 February recommending fishing zones located north of Sumba Island with an area of 300 NM. Other areas do not seem to be potential fishing grounds (IMRO 2020).
5. In March there were 4 files matching, at the time of the fishing trip. March should be the turning point of the monsoon season that occurs in East Nusa Tenggara, but each region has a different time. One of the maps is the 16 to 17 March map, the areas that are recommended for fishing activities are in the south of Sumba Island and northeast of Sumba Island, with an area of more than 300 NM. There was also one point with an area of 50 NM located on Semau and Kupang islands (IMRO 2020).

Compatibility between the fishing ground and PDPPI. Of all trips with available PDPPI, there were only 17 files that matched, at the time of the fishing trip. For the 4 months period, 17 maps of fishing ground forecasts were used, with a total of 44 settings (Table 3).

The method used to determine the suitability of fishing grounds with PDPPI, issued by IMRO, uses a comparison method between the position of the actual fishing area and the status of the same location on the forecast map, on that date. From 44 setting positions, 6 positions were in accordance with the PDPPI, which corresponds to a match rate of 13.6%, in terms of chlorophyll-a and sea surface temperature. When the daily PDPPI images from the NASA website are not available, the fishing trips cannot be used as a reference in determining their suitability. With a match rate of 13.6%, it can be said that fishermen using FADs have sufficient accuracy in determining fishing grounds.


Table 3

Compatibility of the fishing ground with the PPDPI

No	Setting	Position		Chlorophyll-a (mg m ⁻³)	Information
		Latitude	Longitude		
1	1	10 ⁰ 12' 27" S	121 ⁰ 40' 16" E	0.41	Inappropriate
	2	10 ⁰ 09' 43" S	121 ⁰ 37' 40" E	0.31	Inappropriate
2	1	09 ⁰ 56' 38" S	122 ⁰ 25' 03" E	0.3	Appropriate
	2	10 ⁰ 42' 09" S	121 ⁰ 50' 28" E	0.27	Inappropriate
3	3	10 ⁰ 26' 28" S	122 ⁰ 47' 58" E	0.27	Appropriate
	1	09 ⁰ 32' 11" S	123 ⁰ 20' 42" E	0.24	Inappropriate
4	2	09 ⁰ 45' 37" S	123 ⁰ 13' 20" E	0.22	Inappropriate
	1	10 ⁰ 04' 18" S	122 ⁰ 32' 21" E	0.27	Appropriate
5	2	10 ⁰ 28' 31" S	122 ⁰ 41' 16" E	0.31	Inappropriate
	3	10 ⁰ 12' 31" S	122 ⁰ 05' 10" E	0.33	Appropriate
6	1	10 ⁰ 40' 10" S	122 ⁰ 15' 30" E	0.29	Inappropriate
	2	09 ⁰ 40' 33" S	122 ⁰ 55' 15" E	0.39	Inappropriate
7	1	09 ⁰ 31' 40" S	123 ⁰ 05' 09" E	0.21	Inappropriate
	2	09 ⁰ 05' 45" S	123 ⁰ 00' 41" E	0.28	Inappropriate
7	3	09 ⁰ 47' 16" S	122 ⁰ 58' 34" E	0.13	Inappropriate
	1	09 ⁰ 40' 08" S	123 ⁰ 22' 31" E	0.29	Inappropriate
	2	09 ⁰ 50' 28" S	122 ⁰ 53' 17" E	0.30	Inappropriate

No	Setting	Position		Chlorophyll-a (mg m ⁻³)	Information
		Latitude	Longitude		
8	1	10° 38' 15" S	121° 40' 30" E	0.41	Inappropriate
	2	10° 42' 09" S	121° 50' 28" E	0.38	Inappropriate
	3	10° 49' 25" S	121° 01' 51" E	-	Inappropriate
9	1	09° 32' 11" S	123° 20' 42" E	0.28	Inappropriate
	2	09° 45' 37" S	123° 13' 20" E	0.24	Inappropriate
	3	09° 54' 26" S	123° 01' 51" E	-	Inappropriate
10	1	10° 21' 15" S	122° 35' 07" E	0.45	Inappropriate
	2	10° 10' 35" S	122° 05' 37" E	-	Inappropriate
	3	10° 03' 47" S	122° 45' 10" E	0.54	Appropriate
11	1	09° 40' 36" S	122° 05' 15" E	0.8	Inappropriate
	2	09° 50' 17" S	121° 35' 20" E	1.1	Inappropriate
12	1	09° 30' 14" S	122° 50' 33" E	-	Inappropriate
	2	09° 21' 52" S	123° 00' 16" E	-	Inappropriate
13	1	10° 40' 16" S	123° 41' 40" E	-	Inappropriate
	2	10° 39' 13" S	123° 45' 36" E	-	Inappropriate
	3	10° 40' 41" S	123° 40' 22" E	-	Inappropriate
14	1	09° 50' 20" S	122° 10' 31" E	-	Inappropriate
	2	09° 18' 50" S	122° 46' 15" E	-	Inappropriate
15	1	10° 03' 15" S	122° 45' 17" E	0.31	Inappropriate
	2	10° 03' 40" S	121° 48' 29" E	-	Inappropriate
16	1	10° 08' 36" S	123° 07' 12" E	-	Inappropriate
	2	10° 15' 53" S	122° 46' 41" E	-	Inappropriate
	3	10° 03' 09" S	122° 53' 08" E	-	Inappropriate
17	1	09° 29' 21" S	122° 30' 14" E	0.6	Inappropriate
	2	09° 50' 15" S	122° 30' 35" E	0.54	Inappropriate
	3	09° 55' 23" S	121° 20' 54" E	0.49	Appropriate

Concentration and distribution of chlorophyll-a. Chlorophyll-a concentrations can be obtained by downloading data levels 1, 2 and 3 on the NASA website. Data needs a longer time span such as months and years can use level 3 image data with better quality than others, but level 3 images only contain one specific information depending on what is needed. In this study, the basis used is a level 3 image which only requires chlorophyll-a information with a time span of months. Figure 7 images and Table 4 data correspond to level 3 chlorophyll-a concentration.






Figure 7. Concentration of chlorophyll-a map (www.oceancolour.gsfc.nasa.gov).

The distribution map of chlorophyll-a concentrations (Table 4) from November to March, shows elevated values around the west of Flores Island, Sumba Island and also Larantuka, suggesting that food chain occur in the area.

Table 4
Concentration and distribution of chlorophyll-a

<i>Period</i>	<i>November</i>	<i>December</i>	<i>January</i>	<i>February</i>	<i>March</i>
Concentration of chlorophyll-a (mg m^{-3})					
a. Lowest	0.07	0.08	0.08	0.08	0.08
b. Highest	1.29	1.14	1.15	1.19	1.17
Average value (mg m^{-3})	0.2	0.3	0.2	0.2	0.2
Distribution center	North of Kupang city, Sabu Island, and west of Flores	West of Flores	Kupang city, Sabu Island, Rote Island, and west of Flores	Kupang city, Rote Island, and west of Flores	Kupang city, west of Flores
Compatibility level	Appropriate	Inappropriate	Appropriate	Appropriate	Inappropriate

Conclusions. Based on the results of the study it can be concluded: simultaneously, chlorophyll-a had an effect of 62.2% on the catch. The highest concentration of chlorophyll-a recorded was 1.29 mg m⁻³ and the lowest was 0.07 mg m⁻³; the fishing area used by fishermen has a level of conformity with the forecast map for fishing areas of 13.6%; Not all fishing trips can be used as a reference in determining the suitability of fishing grounds with PDPPI, because not every day PDPPI can be uploaded because it is limited by the availability of images obtained from the NASA website.

Acknowledgments. The authors would like to thank to Director and staff of Institute for Marine Research and Observation (IMRO), Agency for Marine and Fisheries Research and Development, Ministry of Marine Affairs and Fisheries, Indonesia. For the support to sharing and processing of the satellite data related to the distribution of SST and chlorophyll-a in fishing ground and to all other parties for their help and contributions.

Conflict of interest. The authors declare no conflict of interest.

References

- Asagabalda M. A., 2017 Law and the role of Lilifuk Amid Government Regimes: Legal analysis and sustainability of local wisdom in Kuanheun Village, West Kupang Regency, East Nusa Tenggara. *SABDA* 12(2):136-145.
- Behrenfeld M. J., Boss E., Siegel D. A., Shea D. M., 2005 Carbon-based ocean productivity and phytoplankton physiology from space. *Global Biogeochemical Cycles* 19(1):1-14.
- Cayula J. F., Cornillon P., 1992 Edge detection algorithm for SST Images. *Journal of Atmospheric and Oceanic Technology* 9(1):67-80.
- Chavula G., Sungani H., Gondwe K., 2012 Mapping potential fishing grounds in Lake Malawi using AVHRR and MODIS satellite imagery. *International Journal of Geosciences* 3(03):650-658.
- Darsono P., 1999 [Utilization of marine resources and its implications for fishing communities]. *Oceana* 24(4):1-9. [In Indonesian].
- Farda N. M., Jatisworo D., 2019 [Mining space-time patterns on forecast maps of fishing areas in fisheries management areas (FMA) 712, 713, and 573]. *Scientific Magazine GLOBE* 21(2):117-128. [In Indonesian].
- Febriani E. R., Sukojo B. M., 2016 [Comparative analysis of the use of level 1b and level 2 MODIS imagery in determining forecasting fishing areas (Case Study: South Coast of Blitar)]. *ITS Engineering Journal, Surabaya* 5(2):439-442. [In Indonesian].
- Hakim D. M., Wikantika K., Widiadnyana N., Napitu A. M., Darmawan S., 2006 The identification of fishing ground area with MODIS satellite image (Case Study: South Coast of West Java) *Proceeding. ITB Engineering Science* 38B(2):147-158.
- Hasyim S., Zainuddin M., 2008 Predictions of skipjack fishing areas based on oceanographic conditions in the waters of Takalar regency and surrounding areas. *Journal of Science and Technology* 8(2):158-162. [In Indonesian].
- Huot Y., Babin M., Bruyant F., Grob C., Twardowski M. S., Claustre H., 2007 Does chlorophyll-a provide the best index of phytoplankton biomass for primary productivity studies? *Biogeosciences Discussions, Proxies of Biomass For Primary production* 4(2):707-745.
- Jatisworo D., Murdimanto A., 2013 [Identification of thermal fronts in the Makassar Strait and Banda Sea]. *Proceedings of the National Symposium on Geoinformation Science III. Marine Research and Observation Institute*, 8 p. [In Indonesian].
- Kunarso, Zainuri M., Sugianto D. N., Marwoto J., Hariyadi, Muslim, 2019 Prediction of demersal fishing ground associated with coral reefs in the Coastal Jepara Regency, Central Java, Indonesia based on Sentinel 2a imagery. *Advances in Science, Technology and Engineering Systems (ASTES) Journal* 4(6):263-269.

- Kurniawan D., 2008 [Linear regression]. Academia accelerating the world's research. Research Development Core Team R: A language and environment for statistical computing, 14 p. [In Indonesian].
- Kusumastanto T., 1996 [An evaluation of government policies on the development of fisheries in Indonesia]. AGRIS, Fisheries Economy Bulletin 2(3):27-41. [In Indonesian].
- Latif B., Noor S. M., Sumardi J., Irwansyah, 2019 Sustainable development and sea protection: Trade on fish and fishery product. ICROEST. IOP Conference Series: Earth and Environmental Science 343(1):1-6.
- Laurs R. M., Fiedler P. C., Montgomery D. R., 1984 Albacore tuna catch distribution relative to environmental features observed from satellites. Deep-Sea Research 31:1085-1099.
- Lee Y. J., Matrai P. A., Friedrichs M. A. M., Saba V. S., Aumont O., Babin M., Buitenhuis E. T., Chevallier M., Mora L., Dessert D., Dunne J. P., Ellingsen I. H., Feldman D., Frouin R., Gehlen M., Gorgues T., Ilyina T., Jin M., John J. G., Lawrence J., Manizza M., Menkes C. E., Perruche C., Fouest V. L., Popova E. E., Romanou A., Samuelsen A., Schwinger J., Séférian R., Stock C. A., Tjiputra J., Tremblay L. B., Ueyoshi K., Vichi M., Yool A., Zhang J., 2016 Net primary productivity estimates and environmental variables in the Arctic Ocean: An assessment of coupled physical-biogeochemical models. Journal of Geophysical Research: Oceans 121(12):8635-8669.
- Nainggolan C., 2012 [Fishing methods]. Open University, Jakarta, 61 p. [In Indonesian].
- Nugraha E., Gunawan R., Danapraja S., Yusrizal, Kusdinar A., Waluyo A. S., Hutajulu J., Prayitno H., Halim S., Sutisna D. H., 2020b The sea surface temperature effect on the length and size of skipjack tuna (*Katsuwonus pelamis*) catches in the Banda Sea, Indonesia. AACL Bioflux 13(1):1-18.
- Nugraha E., Yudho G. S., Jaenudin A., Yusrizal, Kusmedy B., Kusnidar A., Husen E. S., 2020a Relationship between length and weight of skipjack tuna (*Katsuwonus Pelamis*) purse seine catching in the Maluku Sea, Indonesia. AACL Bioflux 13(1):330-345.
- Nurman A., 2010 [Utilization of fashionable data to detect fishing grounds on the east and west coasts of North Sumatra]. Medan State University. Geography Journal 2(2):17-30. [In Indonesian].
- Nusantara R. A., Rosyid A., Boesono H., 2014 [Analysis of differences in depth of fishing area towards fish composition on Danish seine in Rembang waters]. Journal of Fisheries Resources Utilization Management and Technology 3(3):96-104. [In Indonesian].
- Nuzapril M., Susilo S. B., Panjaitan J. P., 2017 Relationship between chlorophyll-a concentration with primary productivity rate using Landsat 8 imagery. Journal of Fisheries and Marine Technology 8(1):105-114.
- Pangestu S., 2016 [Simple linear regression. Applied linear model]. Digital Library of a Open University of Indonesia, 398 p. [In Indonesian].
- Sato K., Matsuno K., Arima D., Abe Y., Yamaguchi A., 2015 Spatial and temporal changes in zooplankton abundance, biovolume, and size spectra in the neighboring waters of Japan: Analyses using an optical plankton counter. Zoological Studies. A Springer Open Journal 54(18):1-15.
- Seki M. P., Polovina J. J., Kobayashi D. R., Bidigare R. R., Mitchum G. T., 2002 An oceanographic characterization of swordfish (*Xiphias gladius*) longline fishing grounds in the springtime subtropical North Pacific. Fisheries Oceanography 11(5):251-266.
- Sudirman, Mallawa A., 2012 [Fishing techniques]. Rineka Cipta, Jakarta, Indonesia, 168 p. [In Indonesian].
- Sugiari, Anshori M. A., Kusumawardani M., 2015 [Coordinate information system for estimated tuna fishing grounds for fishermen]. Journal of JARTEL State Polytechnic of Malang 1(1):74-84. [In Indonesian].

- Suniada K. I., Susilo E., Hastuti A. W., 2015 [Validation of forecast map of fishing grounds (PPDPI) in Java Sea Waters (FMA-RI 712)]. Proceedings of the National Marine and Fisheries Science and Technology Forum 87-96. [In Indonesian].
- Tambun R., Simbolon D., Wahju R., Supartono, 2018 [Potential zones of fishing ground based on season at FMA RI 718]. Journal of Tropical Marine Science and Technology 10(3):757-768. [In Indonesian].
- Tuli M., Menofatria B., Adrianto L., 2015 [Analysis of skipjack fish (*Katsuwonus pelamis*) resources in the waters of Pohuwato Regency, Gorontalo Province]. Marine Fisheries Journal 6(2):109-117. [In Indonesian].
- Wibawa T. A., 2011 [Utilization of oceanographic satellite data for prediction of potential fishing grounds for bigeye tuna (*T. obesus*) in the South Indian Ocean, Java and Bali]. SEGARA Journal 7(1):29-41. [In Indonesian].
- Zainuddin M., 2011 Skipjack tuna in relation to sea surface temperature and chlorophyll-a concentration of Bone Bay using remotely sensed satellite data. Journal of Tropical Marine Science and Technology 3(1):82-90.
- Zainuddin M., Safruddin, Farhum S. A., Nelwan A., Selamat M. B., Hidayat S., Sudirman, 2015 [Characteristics of potential areas for catching skipjack tuna in the Bone-Sea Bay of Flores based on satellite data on sea surface temperature and chlorophyll-a in the period January-June 2014 Makassar]. Journal of Science and Technology Utilization of fisheries resources. Hasanuddin University 2(3):228-237. [In Indonesian].
- Zhang C., Han M., 2015 Mapping chlorophyll-a concentration in Laizhou Bay using Landsat 8 OLI data. E-proceedings of the 36th IAHR World Congress, The Hague, The Netherlands, pp. 12-17.
- *** Institute for Marine Research and Observation, IMRO, 2019 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian].
- *** Institute for Marine Research and Observation, IMRO, 2020 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2016 [East Nusa Tenggara Provincial annual report 2016]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2018 [East Nusa Tenggara Provincial annual report 2018]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** <http://modis.gsfc.nasa.gov> 2007
- *** www.oceancolour.gsfc.nasa.gov

Received: 24 May 2021. Accepted: July 2021. Published online: July 2021.

Authors:

Maman Hermawan, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: mhermawan60@gmail.com
 I Nyoman Radiarta, Institute for Marine Research and Observation, Agency for Marine and Fisheries Research and Development Baru Street, Perancak, Negara, Jembrana 82251 Bali, Indonesia, e-mail: radiarta@kkp.go.id
 I Gusti Bagus Ega Krisna Bayu, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: egabayu91@gmail.com
 Erick Nugraha, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: nugraha_eriq1@yahoo.co.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Hermawan M., Radiarta I. N., Bayu I. G. B. E. K., Nugraha E., 2021 Identifying fishing grounds in the Savu Sea, Indonesia: fishermen's experience compared with the chlorophyll-a forecast maps. AACL Bioflux 14(4):

← Kembali ↶ ↷ →

📁 Arsipkan 📁 Pindahkan 🗑️ Hapus 🛡️ Spam ⋮

☰ ▲ ▼ ✕

• Revision 4

Yahoo/Terkirim ☆

 • **Eni Kovacs** <ek.bioflux@gmail.com>
Kepada: erick nugraha

🖨️ 🔄 Min, 18 Jul 2021 jam 00.21 ☆

Dear Dr. Nugraha,

Regarding your manuscript submitted to AACL Bioflux, the editorial team has some requests prior to acceptance (please see the attachment).

Please note: Always operate corrections/additions (or deletions) in the manuscript we are sending you, by **highlighting with a bright color** (for an easy identification). We never work on the manuscript you send back, just identify the corrections and operate them on our document (to avoid any undesirable accidental operations like changed page set up, otherwise the editors have to start all the work from the beginning, and we cannot ask them to re-check every manuscript word by word to identify unmarked modifications).


Thank you for understanding!

—
Kind regards,
Editor AACL Bioflux
Eniko Kovacs



Nugraha_PP....doc
1.6MB

↶ ↷ → ⋮



Identifying fishing grounds in the Savu Sea, Indonesia: fishermen's experience compared with the chlorophyll-a forecast maps

¹Maman Hermawan, ²I Nyoman Radiarta, ¹I Gusti B. E. K. Bayu, ¹Erick Nugraha

¹ Faculty of Fishing Technology, Jakarta Technical University of Fisheries, Pasarminggu, South Jakarta, Indonesia; ² Institute for Marine Research and Observation, Negara, Jembrana, Bali, Indonesia. Corresponding author: E. Nugraha, nugraha_eriq1@yahoo.co.id

Abstract. The Savu Sea in East Nusa Tenggara Province is the location of fishing activities for most fishermen in East Nusa Tenggara. Despite its potential, the knowledge of the fishing ground is still conventional. Currently, predictive methods for trips planning are available, such as the fishing ground forecast maps based on the chlorophyll-a, sea surface temperature and salinity. The related data are provided by NASA, being processed by the Marine Research and Observation Center to display the potential fishing ground as a box view with an accuracy level of 5 square miles. Eventually, the forecast map of the fishing area will be compared with the fishing locations known by the fishermen, for validation. The data collection used a survey method and the data analysis used both a descriptive analysis and the linear regression. The forecast maps matched with the observed fishing locations at only 13.6%. The highest recorded chlorophyll-a concentration was of 1.29 mg m⁻³ (constantly observed in the West Flores) and the lowest was of 0.07 mg m⁻³. The chlorophyll-a has an effect on 62.2% of the catch and 52% of the catch is composed of *Katsuwonus pelamis*

Key Words: MODIS, sea surface temperature, catch composition, *Katsuwonus pelamis*.

Introduction. Indonesia is the largest archipelagic country in the world, having a coastline length of 81,000 km² and a sea area of about 3,100,000 km² (Kusumastanto 1996; Darsono 1999; Latif et al 2019). Managing the fisheries resource requires determining the fish growth patterns and the catch period, in order to avoid over-exploitation (Nugraha et al 2020b). The province of East Nusa Tenggara has a sea area of 200,000 km² with a coastline of 5,700 km² (MMAF 2016). Savu Sea has a good potential for fisheries and the whole marine sector (Asagabaldan 2017). A fishing area is an area where fish are caught in maximum amount and fishing gear can be operated economically. Assessment and mapping of fishing areas are needed (Nusantara et al 2014). The *Katsuwonus pelamis* is a fast swimming fish and has the greedy character of an opportunistic predator, with food types ranging from small fish (Clupeidae and Engraulidae), squids, crustaceans and zooplankton (Tuli et al 2015). *K. pelamis* is a pelagic fish living in groups, from the coastal areas to the open sea (Nainggolan 2012). Schooling fish should be kept as close to the boat as possible so that they can be easily caught using pole and line fishing rods (Sudirman & Mallawa 2012). The fishing grounds are determined by the waters' condition in the habitat of a species. Water conditions are usually described by oceanographic parameters (Nugraha et al 2020a).

The Sea Surface Temperature (SST) of the *K. pelamis* fishing ground ranges from 29.75 to 30.25°C (Zainuddin et al 2015; Nugraha et al 2020b; Zainuddin 2011). Meanwhile, the potential area has a characteristic chlorophyll-a concentration of around 0.125 mg m⁻³. This condition is mainly seen in May. Outside of this range, the catch of *K. pelamis* tends to decline, as it happens in April (when the SST values in the Bone-Sea Flores bay are the highest). Likewise, when the chlorophyll-a concentration in the fishing ground significantly increase and the catch tends to decrease. This means that there is a

certain range of values for the two oceanographic factors that are very suitable for the presence and abundance of *K. pelamis*. The distribution of chlorophyll-a can be a parameter used for identifying the presence of fish in a water area (Hasyim & Zainuddin 2008; Laurs et al 1984).

The Forecast Maps of Fishing Ground (PPDPI) is the result of a research conducted by the Institute for Marine Research and Observation (IMRO) in order to identify fishing grounds, based on an internet service of information on the fishing areas' characteristics (Sugiari 2015; Panjaitan 2014). PPDPI have been produced for and distributed to the fishing communities of Indonesia, since 2000 (Tambun et al 2018; Farda & Jatisworo 2019). PPDPI was compiled using remote sensing data which is also widely used to detect water conditions such as the thermal front (Cayula & Cornillon 1992; Wibawa 2011; Jatisworo & Murdimanto 2013) and the concentration of chlorophyll-a (Seki et al 2002) which is one of the indicators of water fertility and school of fish. PPDPI preparation was carried out using SST and water fertility data (chlorophyll-a) generated from MODIS Aqua level 3 daily composites, in the Hierarchical Data Format (HDF) (Suniada et al 2015).

The predictive maps of fishing grounds that already exists in Indonesia still have many weaknesses, including the spatial aspect of the points of forecasting located far from the coast and the uncertainty of the predicted type of fish. Thus, if inappropriate gear is selected for the fishing trip, the catch productivity is negatively affected, even in presence of abundant fish populations (Kunarso et al 2019).

Chlorophyll-a is an indicator of the abundance of phytoplankton in waters, playing a role in the photosynthesis process (Huot et al 2007; Zhang & Han 2015). Phytoplankton contributes to a large extent to determine primary productivity in waters. Organic carbon production during photosynthesis is defined as primary productivity or Net Primary Productivity (Lee et al 2016).

Linear regression analysis was carried out on the model of the relationship between chlorophyll-a concentration and primary productivity. The results of this equation can be applied to the satellite imagery so that it can help to monitor the water quality conditions (Nuzapril et al 2017). This abundance of phytoplankton will result in the creation of food chain clusters from phytoplankton to large pelagic fish schools (Sato et al 2015; Behrenfeld et al 2005).




Figure 1. Map of chlorophyll-a concentration (IMRO 2020).

Remote sensing data can show the variables used in determining fish catchment areas. Certain remote sensing images used to extract SST and chlorophyll-a information are provided by the MODIS program (Nurman 2010; Hakim et al 2006).

Satellite imagery is one of the applications of the remote sensing technology. The Moderate-resolution Imaging Spectroradiometer (MODIS) technology uses sensors belonging to the Earth Observing System (EOS), carried by two satellites produced by the National Aeronautics and Space Administration (NASA), namely Terra and Aqua (<http://modis.gsfc.nasa.gov> 2007).

The MODIS sensor is a derivative of the Advanced Very High Resolution Radiometer (AVHRR) sensor, Sea-viewing Wide Field-of-View Sensor (SeaWiFS) and High Resolution Imaging Spectrometer (HRS) owned by EOS, which has previously orbited, so it can be used to measure sea and atmosphere-level parameters (such as the sea surface temperature, chlorophyll-a concentration and water vapor content) and marine phenomena (such as the occurrence of upwelling, thermal fronts and others) (<http://modis.gsfc.nasa.gov> 2007).

Material and Method

Description of the study sites. This research was conducted on a pole and line fishing boat operating in the Flores Sea, East Nusa Tenggara. The satellite image data retrieval was carried out at the Southeast Research and Ocean Observation Center (SeaCORM) Jembrana, Bali. The material used consisted of sea water samples for measuring oceanographic elements. The methods used in data collection are literature study, interview, documentation and observation.

Data analysis method. Regression analysis is a statistical analysis that studies the relationship between two or more variables. In the linear regression analysis it is assumed that this relationship is a linear combination, the simplest case consisting of one independent variable (Pangestu 2016).

Linear regression is a statistical method used to model the relationship between the dependent variable (dependent; response; Y) and one or more independent variables (independent, predictor, X). If there is only one independent variable, it is called simple linear regression, whereas if there is more than one independent variable, it is called multiple linear regression. Regression analysis has at least three use cases, namely for data description, for control and for predictive purposes..

Simple regression is carried out in stages, namely by connecting the independent variable (chlorophyll-a) with the dependent variable (tuna catch), using the data regression analysis function of Microsoft Office Excel (Kurniawan 2008). This simple regression analysis was conducted to determine the relationship between the dependent variable Y (*K. pelamis* catches) and the independent variable X (chlorophyll-a). The regression model used is:

$$Y = a + bX$$

Where:

X - chlorophyll-a;

Y - pole and line catch;

a - the constant (intercept) is the value of Y, when X = 0 which equals the value of a where it intersects the Y axis;

b - regression coefficient.

Chlorophyll-a image analysis. The fishing ground forecast can be predicted from the parameters of sea surface temperature and chlorophyll-a. The analysis of the distribution of sea surface temperature and chlorophyll-a can use level 1b and level 2 Modis images. The Algorithm Theoretical Basic Document Modis (ATBD) 25 is used to determine sea surface temperature and the Morel 4 Algorithm is used to determine the chlorophyll-a. The Modis image levels 1b and 2 use the RMSE test calculation (Chavula 2012).

From the two Root Mean Squared Error (RMSE) tests carried out between the field data and the Modis image data levels 1b and level 2, more accurate values were obtained from the smallest RMSE sum value, namely the Level 2 Modis image, with an RMSE value of chlorophyll-a of 0.090315. So the best fishing ground forecast can be determined through Level 2 Modis Images (Febriani & Sukojo 2016).

Results and Discussion

Fishing ground. In this study we used a pole and line fishing boat (Figure 2) that operates in Savu sea, Flores, East Nusa Tenggara.



Figure 2. Pole and line fishing boat (original).

Composition and comparison of catches with the number of settings.

1. In November 2019 the ship made one trip with a catch of *K. pelamis* of 1,020 kg.
2. In December 2019 the ship made 16 trips with a catch of 28,220 kg. The fishing season generally reaches a turning point in the middle of January and in December, according to the graph, the catch will decrease as the wind season arrives.
3. In January 2020 the catch obtained was of 7,990 kg. January is generally the start of the wind season, so ships rarely go to sea.
4. In February 2020 the catches were dominated by *K. pelamis* and *T. albacares*. According to the fishermen, February was generally the peak of the wind season, so that the catch was not good, until the middle or end of March.
5. In March 2020 the catches were dominated by *K. pelamis* and *T. albacares*, but unlike in the previous month, *C. hippurus* and *E. bipinnulata* were also obtained. In March 2020 the fishermen began to increase the intensity of their trips, but taking into account the strength of the wind and the availability of bait.

The composition of the catch for 5 months can be seen in Table 1.

Table 1

Comparison of catch volumes with the number of settings

Volumes / Period	November	December	January	February	March
Number of setting (times)	2	43	14	15	13
<i>K. pelamis</i> (kg)	1,020	15,300	2,890	6,290	7,480
<i>T. albacares</i> (kg)	0	9,520	2,040	5,440	4,250
<i>C. hippurus</i> (kg)	0	1,870	1,870	0	1,360
<i>E. bipinnulata</i> (kg)	0	1,530	1,190	0	1,870
Total (kg)	1,020	28,220	7,990	11,730	14,960
Catch rate per setting (kg)	510	656,28	570,71	782	1,150

Catch frequencies can also be visualized in Figure 3.




Figure 3. Total catch (November 2019 to March 2020).

Based on the graph above, there is a decrease in the catch volumes during the 4 months of fishing. The lowest catch was obtained in January with a total of 7,990 kg, consisting of: *K. pelamis* (2,890 kg), *T. albacares* (2,040 kg) and others, such as *E. bipinnulata* and *C. hippurus*. The highest catch was obtained in December is 28,220 kg, dominated by *K. pelamis* with 15,300 kg and *T. albacares* with 9,520 kg. In 2016, Kupang Regency was the largest supplier of *K. pelamis* in East Nusa Tenggara (MMAF 2018).

Catch composition. Table 2 and Figure 4 show that *K. pelamis* is the most dominant catch that lands at Tenau port and at the fishing port around Kupang city, East Nusa Tenggara. This is in accordance with the data from the Fisheries Service of the Province of East Nusa Tenggara, suggesting that *K. pelamis* is one of the main commodities produced in Kupang, with 52%, followed by *T. albacares*, with 33%, *C. hippurus*, with 7% and *E. bipinnulata*, with 8%. The catch compositions (in %) can be seen in Figure 4.

Table 2

Catch rated per setting

Name of fish	Amount (kg)	Catch percentage (%)
<i>K. pelamis</i>	32,980	52
<i>T. albacares</i>	21,250	33
<i>C. hippurus</i>	4,590	7
<i>E. bipinnulata</i>	5,100	8
Total catch	63,920	100
Catch rate per setting	734.71	1.14




Figure 4. Catch composition diagram.

Conformity between of the fishing ground and PPDPI. Fishing is carried out on FADs that have been installed, with the position of the FADs stored on the GPS of each

Commented [WU1]: Please use the full name for all the species. Please use English: March

ship. The following is a map of the fishing grounds with the positions of the FADs. Figure 5 below shows the fishing ground for 5 months of fishing.




Figure 5. Fishing ground.

Commented [WU2]: December, not Desember

Forecast map of fishing ground (PPDPI).




Figure 6. PPDPI period in November (1), December (2) (IMRO 2019); January (3), February (4), March (5) (IMRO 2020).

1. In November the PPDPI that can be used correspond to 29 to 30 November 2019. On this map, the potential fishing grounds are around the northwest of Timor Island with an area of 150 NM.
2. In December, 15 files were issued with an interval of 2 days. In the PDPPI 11 to 12 December 2019, the fishing areas that recommended are in the south of Sawu Island, northeast of Sumba Island and north of Rote Island, as well as many other areas with more than 300 NM (IMRO 2019).
3. In January, only 1 file out of 7 uploaded PPDPI files issued corresponded to the time of the fishing trip. On the 31 December to 01 January map, the potential fishing grounds were located around the southeast of Sumba Island, west of Savu Island, with an area of more than 300 NM (IMRO 2020).
4. In February, as many as 10 PPDPI files were uploaded and 3 files corresponded to the time of the fishing trip. One of the suitable map files was issued on the 26 to 27 February recommending fishing zones located north of Sumba Island with an area of 300 NM. Other areas do not seem to be potential fishing grounds (IMRO 2020).
5. In March there were 4 files matching, at the time of the fishing trip. March should be the turning point of the monsoon season that occurs in East Nusa Tenggara, but each region has a different time. One of the maps is the 16 to 17 March map, the areas that are recommended for fishing activities are in the south of Sumba Island and northeast of Sumba Island, with an area of more than 300 NM. There was also one point with an area of 50 NM located on Semau and Kupang islands (IMRO 2020).

Compatibility between the fishing ground and PDPPI. Of all trips with available PDPPI, there were only 17 files that matched, at the time of the fishing trip. For the 4 months period, 17 maps of fishing ground forecasts were used, with a total of 44 settings (Table 3).

The method used to determine the suitability of fishing grounds with PDPPI, issued by IMRO, uses a comparison method between the position of the actual fishing area and the status of the same location on the forecast map, on that date. From 44 setting positions, 6 positions were in accordance with the PDPPI, which corresponds to a match rate of 13.6%, in terms of chlorophyll-a and sea surface temperature. When the daily PDPPI images from the NASA website are not available, the fishing trips cannot be used as a reference in determining their suitability. With a match rate of 13.6%, it can be said that fishermen using FADs have sufficient accuracy in determining fishing grounds.


Table 3

Compatibility of the fishing ground with the PPDPI

No	Setting	Position		Chlorophyll-a (mg m ⁻³)	Information
		Latitude	Longitude		
1	1	10 ⁰ 12' 27" S	121 ⁰ 40' 16" E	0.41	Inappropriate
	2	10 ⁰ 09' 43" S	121 ⁰ 37' 40" E	0.31	Inappropriate
2	1	09 ⁰ 56' 38" S	122 ⁰ 25' 03" E	0.3	Appropriate
	2	10 ⁰ 42' 09" S	121 ⁰ 50' 28" E	0.27	Inappropriate
3	3	10 ⁰ 26' 28" S	122 ⁰ 47' 58" E	0.27	Appropriate
	1	09 ⁰ 32' 11" S	123 ⁰ 20' 42" E	0.24	Inappropriate
4	2	09 ⁰ 45' 37" S	123 ⁰ 13' 20" E	0.22	Inappropriate
	1	10 ⁰ 04' 18" S	122 ⁰ 32' 21" E	0.27	Appropriate
5	2	10 ⁰ 28' 31" S	122 ⁰ 41' 16" E	0.31	Inappropriate
	3	10 ⁰ 12' 31" S	122 ⁰ 05' 10" E	0.33	Appropriate
6	1	10 ⁰ 40' 10" S	122 ⁰ 15' 30" E	0.29	Inappropriate
	2	09 ⁰ 40' 33" S	122 ⁰ 55' 15" E	0.39	Inappropriate
7	1	09 ⁰ 31' 40" S	123 ⁰ 05' 09" E	0.21	Inappropriate
	2	09 ⁰ 05' 45" S	123 ⁰ 00' 41" E	0.28	Inappropriate
7	3	09 ⁰ 47' 16" S	122 ⁰ 58' 34" E	0.13	Inappropriate
	1	09 ⁰ 40' 08" S	123 ⁰ 22' 31" E	0.29	Inappropriate
	2	09 ⁰ 50' 28" S	122 ⁰ 53' 17" E	0.30	Inappropriate

No	Setting	Position		Chlorophyll-a (mg m ⁻³)	Information
		Latitude	Longitude		
8	1	10° 38' 15" S	121° 40' 30" E	0.41	Inappropriate
	2	10° 42' 09" S	121° 50' 28" E	0.38	Inappropriate
	3	10° 49' 25" S	121° 01' 51" E	-	Inappropriate
9	1	09° 32' 11" S	123° 20' 42" E	0.28	Inappropriate
	2	09° 45' 37" S	123° 13' 20" E	0.24	Inappropriate
	3	09° 54' 26" S	123° 01' 51" E	-	Inappropriate
10	1	10° 21' 15" S	122° 35' 07" E	0.45	Inappropriate
	2	10° 10' 35" S	122° 05' 37" E	-	Inappropriate
	3	10° 03' 47" S	122° 45' 10" E	0.54	Appropriate
11	1	09° 40' 36" S	122° 05' 15" E	0.8	Inappropriate
	2	09° 50' 17" S	121° 35' 20" E	1.1	Inappropriate
12	1	09° 30' 14" S	122° 50' 33" E	-	Inappropriate
	2	09° 21' 52" S	123° 00' 16" E	-	Inappropriate
13	1	10° 40' 16" S	123° 41' 40" E	-	Inappropriate
	2	10° 39' 13" S	123° 45' 36" E	-	Inappropriate
	3	10° 40' 41" S	123° 40' 22" E	-	Inappropriate
14	1	09° 50' 20" S	122° 10' 31" E	-	Inappropriate
	2	09° 18' 50" S	122° 46' 15" E	-	Inappropriate
15	1	09° 30' 16" S	123° 01' 51" E	-	Inappropriate
	2	10° 03' 15" S	122° 45' 17" E	0.31	Inappropriate
16	1	10° 03' 40" S	121° 48' 29" E	-	Inappropriate
	2	10° 08' 36" S	123° 07' 12" E	-	Inappropriate
	3	10° 15' 53" S	122° 46' 41" E	-	Inappropriate
17	1	10° 03' 09" S	122° 53' 08" E	-	Inappropriate
	2	09° 29' 21" S	122° 30' 14" E	0.6	Inappropriate
	3	09° 50' 15" S	122° 30' 35" E	0.54	Inappropriate
				0.49	Appropriate

Concentration and distribution of chlorophyll-a. Chlorophyll-a concentrations can be obtained by downloading data levels 1, 2 and 3 on the NASA website. Data needs a longer time span such as months and years can use level 3 image data with better quality than others, but level 3 images only contain one specific information depending on what is needed. In this study, the basis used is a level 3 image which only requires chlorophyll-a information with a time span of months. Figure 7 images and Table 4 data correspond to level 3 chlorophyll-a concentration.






Figure 7. Concentration of chlorophyll-a map (www.oceancolour.gsfc.nasa.gov).

The distribution map of chlorophyll-a concentrations (Table 4) from November to March, shows elevated values around the west of Flores Island, Sumba Island and also Larantuka, suggesting that food chain occur in the area.

Table 4
Concentration and distribution of chlorophyll-a

<i>Period</i>	<i>November</i>	<i>December</i>	<i>January</i>	<i>February</i>	<i>March</i>
Concentration of chlorophyll-a (mg m^{-3})					
a. Lowest	0.07	0.08	0.08	0.08	0.08
b. Highest	1.29	1.14	1.15	1.19	1.17
Average value (mg m^{-3})	0.2	0.3	0.2	0.2	0.2
Distribution center	North of Kupang city, Sabu Island, and west of Flores	West of Flores	Kupang city, Sabu Island, Rote Island, and west of Flores	Kupang city, Rote Island, and west of Flores	Kupang city, west of Flores
Compatibility level	Appropriate	Inappropriate	Appropriate	Appropriate	Inappropriate

Conclusions. Based on the results of the study it can be concluded: simultaneously, chlorophyll-a had an effect of 62.2% on the catch. The highest concentration of chlorophyll-a recorded was 1.29 mg m⁻³ and the lowest was 0.07 mg m⁻³; the fishing area used by fishermen has a level of conformity with the forecast map for fishing areas of 13.6%; Not all fishing trips can be used as a reference in determining the suitability of fishing grounds with PDPPI, because not every day PDPPI can be uploaded because it is limited by the availability of images obtained from the NASA website.

Acknowledgments. The authors would like to thank to Director and staff of Institute for Marine Research and Observation (IMRO), Agency for Marine and Fisheries Research and Development, Ministry of Marine Affairs and Fisheries, Indonesia. For the support to sharing and processing of the satellite data related to the distribution of SST and chlorophyll-a in fishing ground and to all other parties for their help and contributions.

Conflict of interest. The authors declare no conflict of interest.

References

- Asagabalda M. A., 2017 Law and the role of Lilifuk Amid Government Regimes: Legal analysis and sustainability of local wisdom in Kuanheun Village, West Kupang Regency, East Nusa Tenggara. *SABDA* 12(2):136-145.
- Behrenfeld M. J., Boss E., Siegel D. A., Shea D. M., 2005 Carbon-based ocean productivity and phytoplankton physiology from space. *Global Biogeochemical Cycles* 19(1):1-14.
- Cayula J. F., Cornillon P., 1992 Edge detection algorithm for SST Images. *Journal of Atmospheric and Oceanic Technology* 9(1):67-80.
- Chavula G., Sungani H., Gondwe K., 2012 Mapping potential fishing grounds in Lake Malawi using AVHRR and MODIS satellite imagery. *International Journal of Geosciences* 3(03):650-658.
- Darsono P., 1999 [Utilization of marine resources and its implications for fishing communities]. *Oceana* 24(4):1-9. [In Indonesian].
- Farda N. M., Jatisworo D., 2019 [Mining space-time patterns on forecast maps of fishing areas in fisheries management areas (FMA) 712, 713, and 573]. *Scientific Magazine GLOBE* 21(2):117-128. [In Indonesian].
- Febriani E. R., Sukojo B. M., 2016 [Comparative analysis of the use of level 1b and level 2 MODIS imagery in determining forecasting fishing areas (Case Study: South Coast of Blitar)]. *ITS Engineering Journal, Surabaya* 5(2):439-442. [In Indonesian].
- Hakim D. M., Wikantika K., Widiadnyana N., Napitu A. M., Darmawan S., 2006 The identification of fishing ground area with MODIS satellite image (Case Study: South Coast of West Java) *Proceeding. ITB Engineering Science* 38B(2):147-158.
- Hasyim S., Zainuddin M., 2008 Predictions of skipjack fishing areas based on oceanographic conditions in the waters of Takalar regency and surrounding areas. *Journal of Science and Technology* 8(2):158-162. [In Indonesian].
- Huot Y., Babin M., Bruyant F., Grob C., Twardowski M. S., Claustre H., 2007 Does chlorophyll-a provide the best index of phytoplankton biomass for primary productivity studies? *Biogeosciences Discussions, Proxies of Biomass For Primary production* 4(2):707-745.
- Jatisworo D., Murdimanto A., 2013 [Identification of thermal fronts in the Makassar Strait and Banda Sea]. *Proceedings of the National Symposium on Geoinformation Science III. Marine Research and Observation Institute*, 8 p. [In Indonesian].
- Kunarso, Zainuri M., Sugianto D. N., Marwoto J., Hariyadi, Muslim, 2019 Prediction of demersal fishing ground associated with coral reefs in the Coastal Jepara Regency, Central Java, Indonesia based on Sentinel 2a imagery. *Advances in Science, Technology and Engineering Systems (ASTES) Journal* 4(6):263-269.

- Kurniawan D., 2008 [Linear regression]. Academia accelerating the world's research. Research Development Core Team R: A language and environment for statistical computing, 14 p. [In Indonesian].
- Kusumastanto T., 1996 [An evaluation of government policies on the development of fisheries in Indonesia]. AGRIS, Fisheries Economy Bulletin 2(3):27-41. [In Indonesian].
- Latif B., Noor S. M., Sumardi J., Irwansyah, 2019 Sustainable development and sea protection: Trade on fish and fishery product. ICROEST. IOP Conference Series: Earth and Environmental Science 343(1):1-6.
- Laurs R. M., Fiedler P. C., Montgomery D. R., 1984 Albacore tuna catch distribution relative to environmental features observed from satellites. Deep-Sea Research 31:1085-1099.
- Lee Y. J., Matrai P. A., Friedrichs M. A. M., Saba V. S., Aumont O., Babin M., Buitenhuis E. T., Chevallier M., Mora L., Dessert D., Dunne J. P., Ellingsen I. H., Feldman D., Frouin R., Gehlen M., Gorgues T., Ilyina T., Jin M., John J. G., Lawrence J., Manizza M., Menkes C. E., Perruche C., Fouest V. L., Popova E. E., Romanou A., Samuelsen A., Schwinger J., Séférian R., Stock C. A., Tjiputra J., Tremblay L. B., Ueyoshi K., Vichi M., Yool A., Zhang J., 2016 Net primary productivity estimates and environmental variables in the Arctic Ocean: An assessment of coupled physical-biogeochemical models. Journal of Geophysical Research: Oceans 121(12):8635-8669.
- Nainggolan C., 2012 [Fishing methods]. Open University, Jakarta, 61 p. [In Indonesian].
- Nugraha E., Gunawan R., Danapraja S., Yusrizal, Kusdinar A., Waluyo A. S., Hutajulu J., Prayitno H., Halim S., Sutisna D. H., 2020b The sea surface temperature effect on the length and size of skipjack tuna (*Katsuwonus pelamis*) catches in the Banda Sea, Indonesia. AACL Bioflux 13(1):1-18.
- Nugraha E., Yudho G. S., Jaenudin A., Yusrizal, Kusmedy B., Kusnidar A., Husen E. S., 2020a Relationship between length and weight of skipjack tuna (*Katsuwonus Pelamis*) purse seine catching in the Maluku Sea, Indonesia. AACL Bioflux 13(1):330-345.
- Nurman A., 2010 [Utilization of fashionable data to detect fishing grounds on the east and west coasts of North Sumatra]. Medan State University. Geography Journal 2(2):17-30. [In Indonesian].
- Nusantara R. A., Rosyid A., Boesono H., 2014 [Analysis of differences in depth of fishing area towards fish composition on Danish seine in Rembang waters]. Journal of Fisheries Resources Utilization Management and Technology 3(3):96-104. [In Indonesian].
- Nuzapril M., Susilo S. B., Panjaitan J. P., 2017 Relationship between chlorophyll-a concentration with primary productivity rate using Landsat 8 imagery. Journal of Fisheries and Marine Technology 8(1):105-114.
- Pangestu S., 2016 [Simple linear regression. Applied linear model]. Digital Library of a Open University of Indonesia, 398 p. [In Indonesian].
- Sato K., Matsuno K., Arima D., Abe Y., Yamaguchi A., 2015 Spatial and temporal changes in zooplankton abundance, biovolume, and size spectra in the neighboring waters of Japan: Analyses using an optical plankton counter. Zoological Studies. A Springer Open Journal 54(18):1-15.
- Seki M. P., Polovina J. J., Kobayashi D. R., Bidigare R. R., Mitchum G. T., 2002 An oceanographic characterization of swordfish (*Xiphias gladius*) longline fishing grounds in the springtime subtropical North Pacific. Fisheries Oceanography 11(5):251-266.
- Sudirman, Mallawa A., 2012 [Fishing techniques]. Rineka Cipta, Jakarta, Indonesia, 168 p. [In Indonesian].
- Sugiari, Anshori M. A., Kusumawardani M., 2015 [Coordinate information system for estimated tuna fishing grounds for fishermen]. Journal of JARTEL State Polytechnic of Malang 1(1):74-84. [In Indonesian].

- Suniada K. I., Susilo E., Hastuti A. W., 2015 [Validation of forecast map of fishing grounds (PPDPI) in Java Sea Waters (FMA-RI 712)]. Proceedings of the National Marine and Fisheries Science and Technology Forum 87-96. [In Indonesian].
- Tambun R., Simbolon D., Wahju R., Supartono, 2018 [Potential zones of fishing ground based on season at FMA RI 718]. Journal of Tropical Marine Science and Technology 10(3):757-768. [In Indonesian].
- Tuli M., Menofatria B., Adrianto L., 2015 [Analysis of skipjack fish (*Katsuwonus pelamis*) resources in the waters of Pohuwato Regency, Gorontalo Province]. Marine Fisheries Journal 6(2):109-117. [In Indonesian].
- Wibawa T. A., 2011 [Utilization of oceanographic satellite data for prediction of potential fishing grounds for bigeye tuna (*T. obesus*) in the South Indian Ocean, Java and Bali]. SEGARA Journal 7(1):29-41. [In Indonesian].
- Zainuddin M., 2011 Skipjack tuna in relation to sea surface temperature and chlorophyll-a concentration of Bone Bay using remotely sensed satellite data. Journal of Tropical Marine Science and Technology 3(1):82-90.
- Zainuddin M., Safruddin, Farhum S. A., Nelwan A., Selamat M. B., Hidayat S., Sudirman, 2015 [Characteristics of potential areas for catching skipjack tuna in the Bone-Sea Bay of Flores based on satellite data on sea surface temperature and chlorophyll-a in the period January-June 2014 Makassar]. Journal of Science and Technology Utilization of fisheries resources. Hasanuddin University 2(3):228-237. [In Indonesian].
- Zhang C., Han M., 2015 Mapping chlorophyll-a concentration in Laizhou Bay using Landsat 8 OLI data. E-proceedings of the 36th IAHR World Congress, The Hague, The Netherlands, pp. 12-17.
- *** Institute for Marine Research and Observation, IMRO, 2019 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian].
- *** Institute for Marine Research and Observation, IMRO, 2020 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2016 [East Nusa Tenggara Provincial annual report 2016]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2018 [East Nusa Tenggara Provincial annual report 2018]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** <http://modis.gsfc.nasa.gov> 2007
- *** www.oceancolour.gsfc.nasa.gov

Received: 24 May 2021. Accepted: July 2021. Published online: July 2021.

Authors:

Maman Hermawan, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: mhermawan60@gmail.com
 I Nyoman Radiarta, Institute for Marine Research and Observation, Agency for Marine and Fisheries Research and Development Baru Street, Perancak, Negara, Jembrana 82251 Bali, Indonesia, e-mail: radiarta@kkp.go.id
 I Gusti Bagus Ega Krisna Bayu, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: egabayu91@gmail.com
 Erick Nugraha, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: nugraha_eriq1@yahoo.co.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Hermawan M., Radiarta I. N., Bayu I. G. B. E. K., Nugraha E., 2021 Identifying fishing grounds in the Savu Sea, Indonesia: fishermen's experience compared with the chlorophyll-a forecast maps. AACL Bioflux 14(4):

← Kembali ↩️ ↶️ ↷️

📁 Arsipkan 📁 Pindahkan 🗑️ Hapus 🛡️ Spam ⋮

🔍 ▲ ▼ ✕



• **Eni Kovacs** <ek.bioflux@gmail.com>
Kepada: erick nugraha

🖨️ 🔗 Min, 18 Jul 2021 jam 20.22 ☆

Dear Dr. Nugraha,

Could you please check the final form of the article attached, some adjustments were operated. We also changed the title, please see if you agree and make the necessary adjustments according tot he comments.

If you have any final observations and instructions prior to the article publishing, please inform us, any changes should be clearly highlighted. Change requests cannot be considered after publishing the material.


If all the authors agree with the final form of the article, could you please send us your acceptance for publishing, signed by all the authors.

> Tampilkan pesan asli

--
Kind regards,
Editor AACL Bioflux
Eniko Kovacs



Nugraha_PP....doc
1.6MB



Identifying fishing grounds in the Savu Sea, Indonesia: fishermen's experience compared with the chlorophyll-a forecast maps

¹Maman Hermawan, ²I Nyoman Radiarta, ¹I Gusti B. E. K. Bayu, ¹Erick Nugraha

¹ Faculty of Fishing Technology, Jakarta Technical University of Fisheries, Pasarminggu, South Jakarta, Indonesia; ² Institute for Marine Research and Observation, Negara, Jembrana, Bali, Indonesia. Corresponding author: E. Nugraha, nugraha_eriq1@yahoo.co.id

Abstract. The Savu Sea in East Nusa Tenggara Province is the location of fishing activities for most fishermen in East Nusa Tenggara. Despite its potential, the knowledge of the fishing ground is still conventional. Currently, predictive methods for trips planning are available, such as the fishing ground forecast maps based on the chlorophyll-a, sea surface temperature and salinity. The related data are provided by NASA, being processed by the Marine Research and Observation Center to display the potential fishing ground as a box view with an accuracy level of 5 square miles. Eventually, the forecast map of the fishing area will be compared with the fishing locations known by the fishermen, for validation. The data collection used a survey method and the data analysis used both a descriptive analysis and the linear regression. The forecast maps matched with the observed fishing locations at only 13.6%. The highest recorded chlorophyll-a concentration was of 1.29 mg m^{-3} (constantly observed in the West Flores) and the lowest was of 0.07 mg m^{-3} . The chlorophyll-a has an effect on 62.2% of the catch and 52% of the catch is composed of *Katsuwonus pelamis*.

Key Words: MODIS, sea surface temperature, catch composition, *Katsuwonus pelamis*, PPDPI.

Introduction. Indonesia is the largest archipelagic country in the world, having a coastline length of 81,000 km² and a sea area of about 3,100,000 km² (Kusumastanto 1996; Darsono 1999; Latif et al 2019). Managing the fisheries resource requires determining the fish growth patterns and the catch period, in order to avoid over-exploitation (Nugraha et al 2020b). The province of East Nusa Tenggara has a sea area of 200,000 km² with a coastline of 5,700 km² (MMAF 2016). Savu Sea has a good potential for fisheries and the whole marine sector (Asagabaldan 2017). A fishing area is an area where fish are caught in maximum amount and fishing gear can be operated economically. Assessment and mapping of fishing areas are needed (Nusantara et al 2014). The *Katsuwonus pelamis* is a fast swimming fish and has the greedy character of an opportunistic predator, with food types ranging from small fish (Clupeidae and Engraulidae), squids, crustaceans and zooplankton (Tuli et al 2015). *K. pelamis* is a pelagic fish living in groups, from the coastal areas to the open sea (Nainggolan 2012). Schooling fish should be kept as close to the boat as possible so that they can be easily caught using pole and line fishing rods (Sudirman & Mallawa 2012). The fishing grounds are determined by the waters' condition in the habitat of a species. Water conditions are usually described by oceanographic parameters (Nugraha et al 2020a).

The Sea Surface Temperature (SST) of the *K. pelamis* fishing ground ranges from 29.75 to 30.25°C (Zainuddin et al 2015; Nugraha et al 2020b; Zainuddin 2011). Meanwhile, the potential area has a characteristic chlorophyll-a concentration of around 0.125 mg m^{-3} . This condition is mainly seen in May. Outside of this range, the catch of *K. pelamis* tends to decline, as it happens in April (when the SST values in the Bone-Sea Flores bay are the highest). Likewise, when the chlorophyll-a concentration in the fishing ground significantly increase and the catch tends to decrease. This means that there is a

certain range of values for the two oceanographic factors that are very suitable for the presence and abundance of *K. pelamis*. The distribution of chlorophyll-a can be a parameter used for identifying the presence of fish in a water area (Hasyim & Zainuddin 2008; Laurs et al 1984).

The Forecast Maps of Fishing Ground (PPDPI) is the result of a research conducted by the Institute for Marine Research and Observation (IMRO) in order to identify fishing grounds, based on an internet service of information on the fishing areas' characteristics (Sugiari 2015; Panjaitan 2014). PPDPI have been produced for and distributed to the fishing communities of Indonesia, since 2000 (Tambun et al 2018; Farda & Jatisworo 2019). PPDPI was compiled using remote sensing data which is also widely used to detect water conditions such as the thermal front (Cayula & Cornillon 1992; Wibawa 2011; Jatisworo & Murdimanto 2013) and the concentration of chlorophyll-a (Seki et al 2002) which is one of the indicators of water fertility and school of fish. PPDPI preparation was carried out using SST and water fertility data (chlorophyll-a) generated from MODIS Aqua level 3 daily composites, in the Hierarchical Data Format (HDF) (Suniada et al 2015).

The predictive maps of fishing grounds that already exists in Indonesia still have many weaknesses, including the spatial aspect of the points of forecasting located far from the coast and the uncertainty of the predicted type of fish. Thus, if inappropriate gear is selected for the fishing trip, the catch productivity is negatively affected, even in presence of abundant fish populations (Kunarso et al 2019).

Chlorophyll-a is an indicator of the abundance of phytoplankton in waters, playing a role in the photosynthesis process (Huot et al 2007; Zhang & Han 2015). Phytoplankton contributes to a large extent to determine primary productivity in waters. Organic carbon production during photosynthesis is defined as primary productivity or Net Primary Productivity (Lee et al 2016).

Linear regression analysis was carried out on the model of the relationship between chlorophyll-a concentration and primary productivity. The results of this equation can be applied to the satellite imagery so that it can help to monitor the water quality conditions (Nuzapril et al 2017). This abundance of phytoplankton will result in the creation of food chain clusters from phytoplankton to large pelagic fish schools (Sato et al 2015; Behrenfald et al 2005).




Figure 1. Map of chlorophyll-a concentration (IMRO 2020).

Remote sensing data can show the variables used in determining fish catchment areas. Certain remote sensing images used to extract SST and chlorophyll-a information are provided by the MODIS program (Nurman 2010; Hakim et al 2006).

Satellite imagery is one of the applications of the remote sensing technology. The Moderate-resolution Imaging Spectroradiometer (MODIS) technology uses sensors belonging to the Earth Observing System (EOS), carried by two satellites produced by the National Aeronautics and Space Administration (NASA), namely Terra and Aqua (<http://modis.gsfc.nasa.gov> 2007).

The MODIS sensor is a derivative of the Advanced Very High Resolution Radiometer (AVHRR) sensor, Sea-viewing Wide Field-of-View Sensor (SeaWiFS) and High Resolution Imaging Spectrometer (HRS) owned by EOS, which has previously orbited, so it can be used to measure sea and atmosphere-level parameters (such as the sea surface temperature, chlorophyll-a concentration and water vapor content) and marine phenomena (such as the occurrence of upwelling, thermal fronts and others) (<http://modis.gsfc.nasa.gov> 2007).

Material and Method

Description of the study sites. This research was conducted on a pole and line fishing boat operating in the Flores Sea, East Nusa Tenggara. The satellite image data retrieval was carried out at the Southeast Research and Ocean Observation Center (SeaCORM) Jembrana, Bali. The material used consisted of sea water samples for measuring oceanographic elements. The methods used in data collection are literature study, interview, documentation and observation.

Data analysis method. Regression analysis is a statistical analysis that studies the relationship between two or more variables. In the linear regression analysis it is assumed that this relationship is a linear combination, the simplest case consisting of one independent variable (Pangestu 2016).

Linear regression is a statistical method used to model the relationship between the dependent variable (dependent; response; Y) and one or more independent variables (independent, predictor, X). If there is only one independent variable, it is called simple linear regression, whereas if there is more than one independent variable, it is called multiple linear regression. Regression analysis has at least three use cases, namely for data description, for control and for predictive purposes..

Simple regression is carried out in stages, namely by connecting the independent variable (chlorophyll-a) with the dependent variable (tuna catch, using the data regression analysis function of Microsoft Office Excel (Kurniawan 2008). This simple regression analysis was conducted to determine the relationship between the dependent variable Y (*K. pelamis* catches) and the independent variable X (chlorophyll-a). The regression model used is:

$$Y = a + bX$$

Where:

X - chlorophyll-a;

Y - pole and line catch;

a - the constant (intercept) is the value of Y, when X = 0 which equals the value of a where it intersects the Y axis;

b - regression coefficient.

Chlorophyll-a image analysis. The fishing ground forecast can be predicted from the parameters of sea surface temperature and chlorophyll-a. The analysis of the distribution of sea surface temperature and chlorophyll-a can use level 1b and level 2 Modis images. The Algorithm Theoretical Basic Document Modis (ATBD) 25 is used to determine sea surface temperature and the Morel 4 Algorithm is used to determine the chlorophyll-a. The Modis image levels 1b and 2 use the RMSE test calculation (Chavula 2012).

From the two Root Mean Squared Error (RMSE) tests carried out between the field data and the Modis image data levels 1b and level 2, more accurate values were obtained from the smallest RMSE sum value, namely the Level 2 Modis image, with an RMSE value of chlorophyll-a of 0.090315. So the best fishing ground forecast can be determined through Level 2 Modis Images (Febriani & Sukojo 2016).

Results and Discussion

Fishing ground. In this study we used a pole and line fishing boat (Figure 2) that operates in Savu sea, Flores, East Nusa Tenggara.



Figure 2. Pole and line fishing boat (original).

Composition and comparison of catches with the number of settings.

1. In November 2019 the ship made one trip with a catch of *K. pelamis* of 1,020 kg.
2. In December 2019 the ship made 16 trips with a catch of 28,220 kg. The fishing season generally reaches a turning point in the middle of January and in December, according to the graph, the catch will decrease as the wind season arrives.
3. In January 2020 the catch obtained was of 7,990 kg. January is generally the start of the wind season, so ships rarely go to sea.
4. In February 2020 the catches were dominated by *K. pelamis* and *T. albacares*. According to the fishermen, February was generally the peak of the wind season, so that the catch was not good, until the middle or end of March.
5. In March 2020 the catches were dominated by *K. pelamis* and *T. albacares*, but unlike in the previous month, *C. hippurus* and *E. bipinnulata* were also obtained. In March 2020 the fishermen began to increase the intensity of their trips, but taking into account the strength of the wind and the availability of bait.

The composition of the catch for 5 months can be seen in Table 1.

Table 1
Comparison of catch volumes with the number of settings

Volumes / Period	November	December	January	February	March
Number of setting (times)	2	43	14	15	13
<i>K. pelamis</i> (kg)	1,020	15,300	2,890	6,290	7,480
<i>T. albacares</i> (kg)	0	9,520	2,040	5,440	4,250
<i>C. hippurus</i> (kg)	0	1,870	1,870	0	1,360
<i>E. bipinnulata</i> (kg)	0	1,530	1,190	0	1,870
Total (kg)	1,020	28,220	7,990	11,730	14,960
Catch rate per setting (kg)	510	656,28	570,71	782	1,150

Catch frequencies can also be visualized in Figure 3.




Figure 3. Total catch (November 2019 to March 2020).

Based on the graph above, there is a decrease in the catch volumes during the 4 months of fishing. The lowest catch was obtained in January with a total of 7,990 kg, consisting of: *K. pelamis* (2,890 kg), *T. albacares* (2,040 kg) and others, such as *E. bipinnulata* and *C. hippurus*. The highest catch was obtained in December is 28,220 kg, dominated by *K. pelamis* with 15,300 kg and *T. albacares* with 9,520 kg. In 2016, Kupang Regency was the largest supplier of *K. pelamis* in East Nusa Tenggara (MMAF 2018).

Catch composition. Table 2 and Figure 4 show that *K. pelamis* is the most dominant catch that lands at Tenau port and at the fishing port around Kupang city, East Nusa Tenggara. This is in accordance with the data from the Fisheries Service of the Province of East Nusa Tenggara, suggesting that *K. pelamis* is one of the main commodities produced in Kupang, with 52%, followed by *T. albacares*, with 33%, *C. hippurus*, with 7% and *E. bipinnulata*, with 8%. The catch compositions (in %) can be seen in Figure 4.

Table 2

Catch rated per setting

Name of fish	Amount (kg)	Catch percentage (%)
<i>K. pelamis</i>	32,980	52
<i>T. albacares</i>	21,250	33
<i>C. hippurus</i>	4,590	7
<i>E. bipinnulata</i>	5,100	8
Total catch	63,920	100
Catch rate per setting	734.71	1.14




Figure 4. Catch composition diagram.

Conformity between of the fishing ground and PPDPI. Fishing is carried out on FADs that have been installed, with the position of the FADs stored on the GPS of each ship. The following is a map of the fishing grounds with the positions of the FADs. Figure 5 below shows the fishing ground for 5 months of fishing.




Figure 5. Fishing ground.

Forecast map of fishing ground (PPDPI).




Figure 6. PPDPI period in November (1), December (2) (IMRO 2019); January (3), February (4), March (5) (IMRO 2020).

1. In November the PPDPI that can be used correspond to 29 to 30 November 2019. On this map, the potential fishing grounds are around the northwest of Timor Island with an area of 150 NM.
2. In December, 15 files were issued with an interval of 2 days. In the PDPPI 11 to 12 December 2019, the fishing areas that recommended are in the south of Sawu Island, northeast of Sumba Island and north of Rote Island, as well as many other areas with more than 300 NM (IMRO 2019).
3. In January, only 1 file out of 7 uploaded PPDPI files issued corresponded to the time of the fishing trip. On the 31 December to 01 January map, the potential fishing grounds were located around the southeast of Sumba Island, west of Savu Island, with an area of more than 300 NM (IMRO 2020).
4. In February, as many as 10 PPDPI files were uploaded and 3 files corresponded to the time of the fishing trip. One of the suitable map files was issued on the 26 to 27 February recommending fishing zones located north of Sumba Island with an area of 300 NM. Other areas do not seem to be potential fishing grounds (IMRO 2020).
5. In March there were 4 files matching, at the time of the fishing trip. March should be the turning point of the monsoon season that occurs in East Nusa Tenggara, but each region has a different time. One of the maps is the 16 to 17 March map, the areas that are recommended for fishing activities are in the south of Sumba Island and northeast of Sumba Island, with an area of more than 300 NM. There was also one point with an area of 50 NM located on Semau and Kupang islands (IMRO 2020).

Compatibility between the fishing ground and PDPPI. Of all trips with available PDPPI, there were only 17 files that matched, at the time of the fishing trip. For the 4 months period, 17 maps of fishing ground forecasts were used, with a total of 44 settings (Table 3).

The method used to determine the suitability of fishing grounds with PDPPI, issued by IMRO, uses a comparison method between the position of the actual fishing area and the status of the same location on the forecast map, on that date. From 44 setting positions, 6 positions were in accordance with the PDPPI, which corresponds to a match rate of 13.6%, in terms of chlorophyll-a and sea surface temperature. When the daily PDPPI images from the NASA website are not available, the fishing trips cannot be used as a reference in determining their suitability. With a match rate of 13.6%, it can be said that fishermen using FADs have sufficient accuracy in determining fishing grounds.


Table 3

Compatibility of the fishing ground with the PPDPI

No	Setting	Position		Chlorophyll-a (mg m ⁻³)	Information
		Latitude	Longitude		
1	1	10° 12' 27" S	121° 40' 16" E	0.41	Inappropriate
	2	10° 09' 43" S	121° 37' 40" E	0.31	Inappropriate
2	1	09° 56' 38" S	122° 25' 03" E	0.3	Appropriate
	2	10° 42' 09" S	121° 50' 28" E	0.27	Inappropriate
	3	10° 26' 28" S	122° 47' 58" E	0.27	Appropriate
3	1	09° 32' 11" S	123° 20' 42" E	0.24	Inappropriate
	2	09° 45' 37" S	123° 13' 20" E	0.22	Inappropriate
4	1	10° 04' 18" S	122° 32' 21" E	0.27	Appropriate
	2	10° 28' 31" S	122° 41' 16" E	0.31	Inappropriate
	3	10° 12' 31" S	122° 05' 10" E	0.33	Appropriate
5	1	10° 40' 10" S	122° 15' 30" E	0.29	Inappropriate
	2	09° 40' 33" S	122° 55' 15" E	0.39	Inappropriate
6	1	09° 31' 40" S	123° 05' 09" E	0.21	Inappropriate
	2	09° 05' 45" S	123° 00' 41" E	0.28	Inappropriate
	3	09° 47' 16" S	122° 58' 34" E	0.13	Inappropriate
7	1	09° 40' 08" S	123° 22' 31" E	0.29	Inappropriate
	2	09° 50' 28" S	122° 53' 17" E	0.30	Inappropriate

No	Setting	Position		Chllorophyll-a (mg m ⁻³)	Information
		Latitude	Longitude		
8	1	10 ⁰ 38' 15" S	121 ⁰ 40' 30" E	0.41	Inappropriate
	2	10 ⁰ 42' 09" S	121 ⁰ 50' 28" E	0.38	Inappropriate
	3	10 ⁰ 49' 25" S	121 ⁰ 01' 51" E	-	Inappropriate
9	1	09 ⁰ 32' 11" S	123 ⁰ 20' 42" E	0.28	Inappropriate
	2	09 ⁰ 45' 37" S	123 ⁰ 13' 20" E	0.24	Inappropriate
	3	09 ⁰ 54' 26" S	123 ⁰ 01' 51" E	-	Inappropriate
10	1	10 ⁰ 21' 15" S	122 ⁰ 35' 07" E	0.45	Inappropriate
	2	10 ⁰ 10' 35" S	122 ⁰ 05' 37" E	-	Inappropriate
	3	10 ⁰ 03' 47" S	122 ⁰ 45' 10" E	0.54	Appropriate
11	1	09 ⁰ 40' 36" S	122 ⁰ 05' 15" E	0.8	Inappropriate
	2	09 ⁰ 50' 17" S	121 ⁰ 35' 20" E	1.1	Inappropriate
12	1	09 ⁰ 30' 14" S	122 ⁰ 50' 33" E	-	Inappropriate
	2	09 ⁰ 21' 52" S	123 ⁰ 00' 16" E	-	Inappropriate
13	1	10 ⁰ 40' 16" S	123 ⁰ 41' 40" E	-	Inappropriate
	2	10 ⁰ 39' 13" S	123 ⁰ 45' 36" E	-	Inappropriate
	3	10 ⁰ 40' 41" S	123 ⁰ 40' 22" E	-	Inappropriate
14	1	09 ⁰ 50' 20" S	122 ⁰ 10' 31" E	-	Inappropriate
	2	09 ⁰ 18' 50" S	122 ⁰ 46' 15" E	-	Inappropriate
	3	09 ⁰ 30' 16" S	123 ⁰ 01' 51" E	-	Inappropriate
15	1	10 ⁰ 03' 15" S	122 ⁰ 45' 17" E	0.31	Inappropriate
	2	10 ⁰ 03' 40" S	121 ⁰ 48' 29" E	-	Inappropriate
16	1	10 ⁰ 08' 36" S	123 ⁰ 07' 12" E	-	Inappropriate
	2	10 ⁰ 15' 53" S	122 ⁰ 46' 41" E	-	Inappropriate
	3	10 ⁰ 03' 09" S	122 ⁰ 53' 08" E	-	Inappropriate
17	1	09 ⁰ 29' 21" S	122 ⁰ 30' 14" E	0.6	Inappropriate
	2	09 ⁰ 50' 15" S	122 ⁰ 30' 35" E	0.54	Inappropriate
	3	09 ⁰ 55' 23" S	121 ⁰ 20' 54" E	0.49	Appropriate

Concentration and distribution of chlorophyll-a. Chlorophyll-a concentrations can be obtained by downloading data levels 1, 2 and 3 on the NASA website. Data needs a longer time span such as months and years can use level 3 image data with better quality than others, but level 3 images only contain one specific information depending on what is needed. In this study, the basis used is a level 3 image which only requires chlorophyll-a information with a time span of months. Figure 7 images and Table 4 data correspond to level 3 chlorophyll-a concentration.






Figure 7. Concentration of chlorophyll-a map (www.oceancolour.gsfc.nasa.gov).

The distribution map of chlorophyll-a concentrations (Table 4) from November to March, shows elevated values around the west of Flores Island, Sumba Island and also Larantuka, suggesting that food chain occur in the area.

Table 4
Concentration and distribution of chlorophyll-a

<i>Period</i>	<i>November</i>	<i>December</i>	<i>January</i>	<i>February</i>	<i>March</i>
Concentration of chlorophyll-a (mg m^{-3})					
a. Lowest	0.07	0.08	0.08	0.08	0.08
b. Highest	1.29	1.14	1.15	1.19	1.17
Average value (mg m^{-3})	0.2	0.3	0.2	0.2	0.2
Distribution center	North of Kupang city, Sabu Island, and west of Flores	West of Flores	Kupang city, Sabu Island, Rote Island, and west of Flores	Kupang city, Rote Island, and west of Flores	Kupang city, west of Flores
Compatibility level	Appropriate	Inappropriate	Appropriate	Appropriate	Inappropriate

Conclusions. Based on the results of the study it can be concluded: simultaneously, chlorophyll-a had an effect of 62.2% on the catch. The highest concentration of chlorophyll-a recorded was 1.29 mg m⁻³ and the lowest was 0.07 mg m⁻³; the fishing area used by fishermen has a level of conformity with the forecast map for fishing areas of 13.6%; Not all fishing trips can be used as a reference in determining the suitability of fishing grounds with PDPPI, because not every day PDPPI can be uploaded because it is limited by the availability of images obtained from the NASA website.

Acknowledgments. The authors would like to thank to Director and staff of Institute for Marine Research and Observation (IMRO), Agency for Marine and Fisheries Research and Development, Ministry of Marine Affairs and Fisheries, Indonesia. For the support to sharing and processing of the satellite data related to the distribution of SST and chlorophyll-a in fishing ground and to all other parties for their help and contributions.

Conflict of interest. The authors declare no conflict of interest.

References

- Asagabaldan M. A., 2017 Law and the role of Lilifuk Amid Government Regimes: Legal analysis and sustainability of local wisdom in Kuanheun Village, West Kupang Regency, East Nusa Tenggara. *SABDA* 12(2):136-145.
- Behrenfeld M. J., Boss E., Siegel D. A., Shea D. M., 2005 Carbon-based ocean productivity and phytoplankton physiology from space. *Global Biogeochemical Cycles* 19(1):1-14.
- Cayula J. F., Cornillon P., 1992 Edge detection algorithm for SST Images. *Journal of Atmospheric and Oceanic Technology* 9(1):67-80.
- Chavula G., Sungani H., Gondwe K., 2012 Mapping potential fishing grounds in Lake Malawi using AVHRR and MODIS satellite imagery. *International Journal of Geosciences* 3(03):650-658.
- Darsono P., 1999 [Utilization of marine resources and its implications for fishing communities]. *Oceana* 24(4):1-9. [In Indonesian].
- Farda N. M., Jatisworo D., 2019 [Mining space-time patterns on forecast maps of fishing areas in fisheries management areas (FMA) 712, 713, and 573]. *Scientific Magazine GLOBE* 21(2):117-128. [In Indonesian].
- Febriani E. R., Sukojo B. M., 2016 [Comparative analysis of the use of level 1b and level 2 MODIS imagery in determining forecasting fishing areas (Case Study: South Coast of Blitar)]. *ITS Engineering Journal, Surabaya* 5(2):439-442. [In Indonesian].
- Hakim D. M., Wikantika K., Widiadnyana N., Napitu A. M., Darmawan S., 2006 The identification of fishing ground area with MODIS satellite image (Case Study: South Coast of West Java) *Proceeding. ITB Engineering Science* 38B(2):147-158.
- Hasyim S., Zainuddin M., 2008 Predictions of skipjack fishing areas based on oceanographic conditions in the waters of Takalar regency and surrounding areas. *Journal of Science and Technology* 8(2):158-162. [In Indonesian].
- Huot Y., Babin M., Bruyant F., Grob C., Twardowski M. S., Claustre H., 2007 Does chlorophyll-a provide the best index of phytoplankton biomass for primary productivity studies? *Biogeosciences Discussions, Proxies of Biomass For Primary production* 4(2):707-745.
- Jatisworo D., Murdimanto A., 2013 [Identification of thermal fronts in the Makassar Strait and Banda Sea]. *Proceedings of the National Symposium on Geoinformation Science III. Marine Research and Observation Institute*, 8 p. [In Indonesian].
- Kunarso, Zainuri M., Sugianto D. N., Marwoto J., Hariyadi, Muslim, 2019 Prediction of demersal fishing ground associated with coral reefs in the Coastal Jepara Regency, Central Java, Indonesia based on Sentinel 2a imagery. *Advances in Science, Technology and Engineering Systems (ASTES) Journal* 4(6):263-269.

- Kurniawan D., 2008 [Linear regression]. Academia accelerating the world's research. Research Development Core Team R: A language and environment for statistical computing, 14 p. [In Indonesian].
- Kusumastanto T., 1996 [An evaluation of government policies on the development of fisheries in Indonesia]. *AGRIS, Fisheries Economy Bulletin* 2(3):27-41. [In Indonesian].
- Latif B., Noor S. M., Sumardi J., Irwansyah, 2019 Sustainable development and sea protection: Trade on fish and fishery product. ICROEST. IOP Conference Series: Earth and Environmental Science 343(1):1-6.
- Laurs R. M., Fiedler P. C., Montgomery D. R., 1984 Albacore tuna catch distribution relative to environmental features observed from satellites. *Deep-Sea Research* 31:1085-1099.
- Lee Y. J., Matrai P. A., Friedrichs M. A. M., Saba V. S., Aumont O., Babin M., Buitenhuis E. T., Chevallier M., Mora L., Dessert D., Dunne J. P., Ellingsen I. H., Feldman D., Frouin R., Gehlen M., Gorgues T., Ilyina T., Jin M., John J. G., Lawrence J., Manizza M., Menkes C. E., Perruche C., Fouest V. L., Popova E. E., Romanou A., Samuelsen A., Schwinger J., Séférian R., Stock C. A., Tjiputra J., Tremblay L. B., Ueyoshi K., Vichi M., Yool A., Zhang J., 2016 Net primary productivity estimates and environmental variables in the Arctic Ocean: An assessment of coupled physical-biogeochemical models. *Journal of Geophysical Research: Oceans* 121(12):8635-8669.
- Nainggolan C., 2012 [Fishing methods]. Open University, Jakarta, 61 p. [In Indonesian].
- Nugraha E., Gunawan R., Danapraja S., Yusrizal, Kusdinar A., Waluyo A. S., Hutajulu J., Prayitno H., Halim S., Sutisna D. H., 2020b The sea surface temperature effect on the length and size of skipjack tuna (*Katsuwonus pelamis*) catches in the Banda Sea, Indonesia. *AACL Bioflux* 13(1):1-18.
- Nugraha E., Yudho G. S., Jaenudin A., Yusrizal, Kusmedy B., Kusnidar A., Husen E. S., 2020a Relationship between length and weight of skipjack tuna (*Katsuwonus Pelamis*) purse seine catching in the Maluku Sea, Indonesia. *AACL Bioflux* 13(1):330-345.
- Nurman A., 2010 [Utilization of fashionable data to detect fishing grounds on the east and west coasts of North Sumatra]. Medan State University. *Geography Journal* 2(2):17-30. [In Indonesian].
- Nusantara R. A., Rosyid A., Boesono H., 2014 [Analysis of differences in depth of fishing area towards fish composition on Danish seine in Rembang waters]. *Journal of Fisheries Resources Utilization Management and Technology* 3(3):96-104. [In Indonesian].
- Nuzapril M., Susilo S. B., Panjaitan J. P., 2017 Relationship between chlorophyll-a concentration with primary productivity rate using Landsat 8 imagery. *Journal of Fisheries and Marine Technology* 8(1):105-114.
- Pangestu S., 2016 [Simple linear regression. Applied linear model]. Digital Library of a Open University of Indonesia, 398 p. [In Indonesian].
- Sato K., Matsuno K., Arima D., Abe Y., Yamaguchi A., 2015 Spatial and temporal changes in zooplankton abundance, biovolume, and size spectra in the neighboring waters of Japan: Analyses using an optical plankton counter. *Zoological Studies. A Springer Open Journal* 54(18):1-15.
- Seki M. P., Polovina J. J., Kobayashi D. R., Bidigare R. R., Mitchum G. T., 2002 An oceanographic characterization of swordfish (*Xiphias gladius*) longline fishing grounds in the springtime subtropical North Pacific. *Fisheries Oceanography* 11(5):251-266.
- Sudirman, Mallawa A., 2012 [Fishing techniques]. Rineka Cipta, Jakarta, Indonesia, 168 p. [In Indonesian].
- Sugiari, Anshori M. A., Kusumawardani M., 2015 [Coordinate information system for estimated tuna fishing grounds for fishermen]. *Journal of JARTEL State Polytechnic of Malang* 1(1):74-84. [In Indonesian].

- Suniada K. I., Susilo E., Hastuti A. W., 2015 [Validation of forecast map of fishing grounds (PPDPI) in Java Sea Waters (FMA-RI 712)]. Proceedings of the National Marine and Fisheries Science and Technology Forum 87-96. [In Indonesian].
- Tambun R., Simbolon D., Wahju R., Supartono, 2018 [Potential zones of fishing ground based on season at FMA RI 718]. Journal of Tropical Marine Science and Technology 10(3):757-768. [In Indonesian].
- Tuli M., Menofatria B., Adrianto L., 2015 [Analysis of skipjack fish (*Katsuwonus pelamis*) resources in the waters of Pohuwato Regency, Gorontalo Province]. Marine Fisheries Journal 6(2):109-117. [In Indonesian].
- Wibawa T. A., 2011 [Utilization of oceanographic satellite data for prediction of potential fishing grounds for bigeye tuna (*T. obesus*) in the South Indian Ocean, Java and Bali]. SEGARA Journal 7(1):29-41. [In Indonesian].
- Zainuddin M., 2011 Skipjack tuna in relation to sea surface temperature and chlorophyll-a concentration of Bone Bay using remotely sensed satellite data. Journal of Tropical Marine Science and Technology 3(1):82-90.
- Zainuddin M., Safruddin, Farhum S. A., Nelwan A., Selamat M. B., Hidayat S., Sudirman, 2015 [Characteristics of potential areas for catching skipjack tuna in the Bone-Sea Bay of Flores based on satellite data on sea surface temperature and chlorophyll-a in the period January-June 2014 Makassar]. Journal of Science and Technology Utilization of fisheries resources. Hasanuddin University 2(3):228-237. [In Indonesian].
- Zhang C., Han M., 2015 Mapping chlorophyll-a concentration in Laizhou Bay using Landsat 8 OLI data. E-proceedings of the 36th IAHR World Congress, The Hague, The Netherlands, pp. 12-17.
- *** Institute for Marine Research and Observation, IMRO, 2019 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian].
- *** Institute for Marine Research and Observation, IMRO, 2020 [Forecast map of fishing ground]. Bali. <http://bpol.litbang.kkp.go.id/> [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2016 [East Nusa Tenggara Provincial annual report 2016]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** Ministry of Marine Affairs and Fisheries, MMAF, 2018 [East Nusa Tenggara Provincial annual report 2018]. East Nusa Tenggara Provincial Marine and Fisheries Service. [In Indonesian].
- *** <http://modis.gsfc.nasa.gov> 2007
- *** www.oceancolour.gsfc.nasa.gov

Received: 24 May 2021. Accepted: July 2021. Published online: July 2021.

Authors:

Maman Hermawan, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: mhermawan60@gmail.com

I Nyoman Radiarta, Institute for Marine Research and Observation, Agency for Marine and Fisheries Research and Development Baru Street, Perancak, Negara, Jembrana 82251 Bali, Indonesia, e-mail: radiarta@kkp.go.id

I Gusti Bagus Ega Krisna Bayu, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: egabayu91@gmail.com

Erick Nugraha, Jakarta Technical University of Fisheries, Faculty of Fishing Technology, AUP Street Pasarminggu 12520, South Jakarta, Indonesia, e-mail: nugraha_eriq1@yahoo.co.id

This is an open-access article distributed under the terms of the Creative Commons Attribution License, which permits unrestricted use, distribution and reproduction in any medium, provided the original author and source are credited.

How to cite this article:

Hermawan M., Radiarta I. N., Bayu I. G. B. E. K., Nugraha E., 2021 Identifying fishing grounds in the Savu Sea, Indonesia: fishermen's experience compared with the chlorophyll-a forecast maps. AACL Bioflux 14(4):



erick nugraha <nugraha_eriq1@yahoo.co.id>
Kepada: Eni Kovacs

Min, 18 Jul 2021 jam 21.42 ☆

Dear Dr. Eni,

I send you edited manuscript journal.
thank for your cooperation.

Regards,

Erick Nugraha

> Tampilkan pesan asli

↓ Unduh semua lampiran sebagai file zip




Acceptancepdf
148,5kB



Nugraha_PP....doc
1,6MB





ACCEPTANCE FOR PUBLISHING

Hereby we declare our article with the title:

Identifying fishing grounds in the Savu Sea, Indonesia: fishermen's experience compared with the chlorophyll-a forecast maps

It has gone through several editing processes and we agreed to publish it.

Thank you.

Name of the authors:

Maman Hermawan

I Nyoman Radiarta

I Gusti Bagus Ega Krisna Bayu

Erick Nugraha

July 18, 2021

← Kembali ↩ ⏪ ⏩

📁 Arsipkan

📁 Pindahkan

🗑 Hapus

🛡 Spam



• Published manuscript

Yahoo/Email M... ☆



• **Eni Kovacs** <ek.bioflux@gmail.com>

Kepada: erick nugraha



Rab, 21 Jul 2021 jam 02.55 ☆

Dear Dr. Nugraha,

Your manuscript has been published:

<http://www.bioflux.com.ro/docs/2021.1943-1954.pdf>

Thank you for your collaboration!

--

Kind regards,
Editor AAAL Bioflux
Eniko Kovacs

